




Do not use if aluminum seal is broken or removed before use.

KEEP THIS END UP


**SANDIMMUNE®**  
**Oral Solution**  
**(cyclosporine oral solution, USP)**

300125665-  
US


Precision “pump” allows accurate dosing, less waste, less mess. Do not rinse syringe or tube before use.




1. To open, flip-up cap and pull to unwrap aluminum seal. Remove rubber stopper and discard.




2. Insert tube with attached white stopper (not syringe) into Sandimmune® (cyclosporine) bottle until it snaps into place.



3. Insert tip of syringe into opening at top of white stopper. Note, do not rinse syringe before use. If water gets into the product through the syringe or any other means, it will cause variation in dose. Draw up the prescribed amount of Sandimmune® Oral Solution.



4. If large bubbles form in syringe during withdrawal, empty the Sandimmune® solution back into the bottle and repeat withdrawal procedure.



5. After use, wipe syringe with dry tissue, **do not rinse**. Store syringe in clear case. Note, the tube with attached white stopper should remain in the bottle. Reseal bottle with screw-on cap.

NDC 0078-0110-22

50 mL

**SANDIMMUNE®**  
**Oral Solution**  
**(cyclosporine oral solution, USP)**

**100mg/mL**

This bottle is not child-resistant (CR) until white CR cap is put on.

**Dosage:** See package insert for dosage information.

**WARNING:** Sandimmune® (cyclosporine oral solution, USP) is NOT BIOEQUIVALENT to Neoral® (cyclosporine oral solution, USP) MODIFIED. Do NOT use interchangeably without a physician's supervision.

**Store and dispense:** In the original container below 30°C (86°F). Do not refrigerate. Protect from freezing. Once opened, the contents must be used within 2 months.



50 mL

**SANDIMMUNE®**  
**Oral Solution**  
**(cyclosporine oral solution, USP)**

**100mg/mL**



**WARNING:** Sandimmune® (cyclosporine oral solution, USP) is NOT BIOEQUIVALENT to Neoral® (cyclosporine oral solution, USP) MODIFIED. Do NOT use interchangeably without a physician's supervision.

**Each mL contains:**  
cyclosporine, USP ..... 100 mg  
dehydrated alcohol, Ph. Helv. .... 12.5%

**Rx only**

Manufactured by:  
DELPHARM Huningue S.A.S.  
Huningue, France

Product of India

Distributed by:  
Novartis Pharmaceuticals Corporation  
East Hanover, New Jersey 07936

**NOVARTIS**

50 mL

**SANDIMMUNE®**  
**Oral Solution**  
**(cyclosporine oral solution, USP)**

**100mg/mL**

This bottle is not child-resistant (CR) until white CR cap is put on.

**Dosage:** See package insert for dosage information.

**WARNING:** Sandimmune® (cyclosporine oral solution, USP) is NOT BIOEQUIVALENT to Neoral® (cyclosporine oral solution, USP) MODIFIED. Do NOT use interchangeably without a physician's supervision.

**Store and dispense:** In the original container below 30°C (86°F). Do not refrigerate. Protect from freezing. Once opened, the contents must be used within 2 months.

©Novartis

GTIN: 00300780110228

EXP  
LOT



Read Direction (123)



Vilayat  
Sayeed

Digitally signed by Vilayat Sayeed

Date: 10/08/2025 07:33:15AM

GUID: 508da70600028a7aad41a1cc455d8261