

**CENTER FOR DRUG EVALUATION AND RESEARCH**

**APPROVAL PACKAGE FOR:**

**APPLICATION NUMBER(S)**

**NDA 20-262/S-036/S-037/S-038**

**Trade Name:** Taxol for Injection

**Generic Name(s):** (paclitaxel)

**Sponsor:** Bristol-Myers Squibb Company

**Agent:**

**Approval Date:** March 4, 2002

**Indication:** Provides for changes in the ADVERSE REACTIONS section of the package insert by incorporating ototoxicity into the list of adverse events reported during continuing surveillance of safety

**CENTER FOR DRUG EVALUATION AND RESEARCH**

**APPROVAL PACKAGE FOR:**

**APPLICATION NUMBER**

**NDA 20-262/S-036/S-037/S-038**

**Approval Letter(s)**



NDA 20-262/S-036/S-037/S-038

Bristol-Myers Squibb Company  
5 Research Parkway  
Wallingford, CT 06492

Attention: Susan H. Behling, Director  
Oncology Global Strategic Unit, Worldwide Regulatory Affairs

Dear Ms. Behling:

Please refer to your supplemental new drug application dated August 8, 2000 (S-038), received August 9, 2000, submitted under section 505(b) of the Federal Food, Drug, and Cosmetic Act for Taxol (paclitaxel) Injection.

We also refer to your supplemental new drug application dated May 11 (S-037), received on May 12, 2000 and your final printed labeling (FPL) for S-036 dated July 6, received on July 7, 2000.

The "Changes Being Effectuated" supplemental new drug application 037 dated May 11, 2000, provides for changes to the **ADVERSE REACTIONS** section of the package insert by incorporating ototoxicity into the list of adverse events reported during continuing surveillance of Taxol safety. The submission dated July 6, 2000, provides final printed labeling in follow-up to the Approval letter dated June 20, 2000, for supplement 036. Both of these supplements have been superseded by supplement 038 and will be Acknowledged and Retained.

The "Changes Being Effectuated" supplemental new drug application 038 provides for changes to the **ADVERSE REACTIONS** section of the package insert by incorporating Steven's Johnson Syndrome and toxic epidermal necrolysis to the list of adverse events reported during the continuing surveillance of Taxol safety.

We have completed the review of supplemental application 038 and have concluded that adequate information has been presented to demonstrate that the drug product is safe and effective for use as recommended in the submitted final printed labeling (package insert and patient package insert submitted on August 8, 2000). Accordingly, the supplemental application is approved effective on the date of this letter.

If a letter communicating important information about this drug product (i.e., a "Dear Health Care Professional" letter) is issued to physicians and others responsible for patient care, we request that you submit a copy of the letter to this NDA and a copy to the following address:

MEDWATCH, HF-2  
FDA  
5600 Fishers Lane  
Rockville, MD 20857

We remind you that you must comply with the requirements for an approved NDA set forth under 21 CFR 314.80 and 314.81.

If you have any questions, call Christy Wilson, Consumer Safety Officer, at (301) 594-5761.

Sincerely,

*{See appended electronic signature page}*

Richard Pazdur, M.D.  
Director  
Division of Oncology Drug Products  
Office of Drug Evaluation I  
Center for Drug Evaluation and Research

-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/

-----  
Richard Pazdur  
3/4/02 02:56:15 PM

**CENTER FOR DRUG EVALUATION AND RESEARCH**

**APPROVAL PACKAGE FOR:**

**APPLICATION NUMBER**

**NDA 20-262/S-036/S-037/S-038**

**Approved Labeling**

# TAXOL® (paclitaxel) Injection

(Patient Information Included)

## WARNING

TAXOL® (paclitaxel) Injection should be administered under the supervision of a physician experienced in the use of cancer chemotherapeutic agents. Appropriate management of complications is possible only when adequate diagnostic and treatment facilities are readily available.

Anaphylaxis and severe hypersensitivity reactions characterized by dyspnea and hypotension requiring treatment, angioedema, and generalized urticaria have occurred in 2%-4% of patients receiving TAXOL in clinical trials. Fatal reactions have occurred in patients despite premedication. All patients should be pretreated with corticosteroids, diphenhydramine, and H<sub>2</sub> antagonists. (See **DOSE AND ADMINISTRATION**.) Patients who experience severe hypersensitivity reactions to TAXOL should not be rechallenged with the drug.


TAXOL therapy should not be given to patients with solid tumors who have baseline neutrophil counts of less than 1,500 cells/mm<sup>3</sup> and should not be given to patients with AIDS-related Kaposi's sarcoma if the baseline neutrophil count is less than 1000 cells/mm<sup>3</sup>. In order to monitor the occurrence of bone marrow suppression, primarily neutropenia, which may be severe and result in infection, it is recommended that frequent peripheral blood cell counts be performed on all patients receiving TAXOL.

## DESCRIPTION

TAXOL® (paclitaxel) Injection is a clear colorless to slightly yellow viscous solution. It is supplied as a nonaqueous solution intended for dilution with a suitable parenteral fluid prior to intravenous infusion. TAXOL is available in 30 mg (5 mL), 100 mg (16.7 mL), and 300 mg (50 mL) multidose vials. Each mL of sterile nonpyrogenic solution contains 6 mg paclitaxel, 527 mg of purified Cremophor® EL<sup>1</sup> (polyoxyethylated castor oil) and 49.7% (w/v) dehydrated alcohol, USP.

Paclitaxel is a natural product with antitumor activity. TAXOL is obtained via a semi-synthetic process from *Taxus baccata*. The chemical name for paclitaxel is 5β,20-Epoxy-1,2α,4,7β,10β,13α-hexahydroxytax-11-en-9-one 4,10-diacetate 2-benzoate 13-ester with (2R,3S)-N-benzoyl-3-phenylisoserine.

Paclitaxel has the following structural formula:



Paclitaxel is a white to off-white crystalline powder with the empirical formula C<sub>47</sub>H<sub>51</sub>NO<sub>14</sub> and a molecular weight of 853.9. It is highly lipophilic, insoluble in water, and melts at around 216-217°C.

## CLINICAL PHARMACOLOGY

Paclitaxel is a novel antimicrotubule agent that promotes the assembly of microtubules from tubulin dimers and stabilizes microtubules by preventing depolymerization. This stability results in the inhibition of the normal dynamic reorganization of the microtubule network that is essential for vital interphase and mitotic cellular functions. In addition, paclitaxel induces abnormal arrays or "bundles" of microtubules throughout the cell cycle and multiple arrests of microtubules during mitosis.

Following intravenous administration of TAXOL, paclitaxel plasma concentrations declined in a biphasic manner. The initial rapid decline represents distribution to the peripheral compartment and elimination of the drug. The later phase is due, in part, to a relatively slow efflux of paclitaxel from the peripheral compartment.

Pharmacokinetic parameters of paclitaxel following 3- and 24-hour infusions of TAXOL at dose levels of 135 and 175 mg/m<sup>2</sup> were determined in a Phase 3 randomized study in ovarian cancer patients and are summarized in the following table:

Table 1: Summary of Pharmacokinetic Parameters - Mean Values

Dose (mg/m <sup>2</sup> )	Infusion Duration (h)	N (patients)	C <sub>max</sub> (ng/mL)	AUC (0-∞) (ng·h/mL)	T <sub>1/2</sub> (h)	CL <sub>T</sub> (L/h/m <sup>2</sup> )
135	24	2	195	6300	52.7	21.7
175	24	4	365	7993	15.7	23.8
135	3	7	2170	7952	13.1	17.7
175	3	5	3650	15007	20.2	12.2

C<sub>max</sub> = Maximum plasma concentration  
AUC (0-∞) = Area under the plasma concentration-time curve from time 0 to infinity  
CL<sub>T</sub> = Total body clearance

It appeared that with the 24-hour infusion of TAXOL, a 30% increase in dose (135 mg/m<sup>2</sup> versus 175 mg/m<sup>2</sup>) increased the C<sub>max</sub> by 87%, whereas the AUC (0-∞) remained proportional. However, with a 3-hour infusion, for a 30% increase in dose, the C<sub>max</sub> and AUC (0-∞) were increased by 68% and 89%, respectively. The mean apparent volume of distribution at steady state, with the 24-hour infusion of TAXOL, ranged from 227 to 688 L/m<sup>2</sup>, indicating extensive extravascular distribution and/or tissue binding of paclitaxel.

The pharmacokinetics of paclitaxel were also evaluated in adult cancer patients who received single doses of 15-135 mg/m<sup>2</sup> given by 1-hour infusions (n=15), 30-275 mg/m<sup>2</sup> given by 6-hour infusions (n=36), and 200-275 mg/m<sup>2</sup> given by 24-hour infusions (n=54) in Phase 1 & 2 studies. Values for CL<sub>T</sub> and volume of distribution were consistent with the findings in the Phase 3 study. The pharmacokinetics of TAXOL in patients with AIDS-related Kaposi's sarcoma have not been studied.

<sup>1</sup> Cremophor® EL is the registered trademark of BASF Aktiengesellschaft. Cremophor® EL is further purified by a Bristol-Myers Squibb Company proprietary process before use.

*In vitro* studies of binding to human serum proteins, using paclitaxel concentrations ranging from 0.1 to 50 µg/mL, indicate that between 89%-98% of drug is bound; the presence of cimetidine, ranitidine, dexamethasone, or diphenhydramine did not affect protein binding of paclitaxel.

After intravenous administration of 15-275 mg/m<sup>2</sup> doses of TAXOL (paclitaxel) injection as 1-, 6-, or 24-hour infusions, mean values for cumulative urinary recovery of unchanged drug ranged from 1.3% to 12.6% of the dose, indicating extensive non-renal clearance. In five patients administered a 225 or 250 mg/m<sup>2</sup> dose of radiolabeled TAXOL as a 3-hour infusion, a mean of 71% of the radioactivity was excreted in the feces in 120 hours, and 14% was recovered in the urine. Total recovery of radioactivity ranged from 56% to 101% of the dose. Paclitaxel represented a mean of 5% of the administered radioactivity recovered in the feces, while metabolites, primarily 6α-hydroxypaclitaxel, accounted for the balance. *In vitro* studies with human liver microsomes and tissue slices showed that paclitaxel was metabolized primarily to 6α-hydroxypaclitaxel by the cytochrome P450 isozyme CYP2C8; and to two minor metabolites, 3'-p-hydroxypaclitaxel and 6α, 3'-p-dihydroxypaclitaxel, by CYP3A4. *In vitro*, the metabolism of paclitaxel to 6α-hydroxypaclitaxel was inhibited by a number of agents (ketoconazole, verapamil, diazepam, quinidine, dexamethasone, cyclosporin, teniposide, etoposide, and vincristine), but the concentrations used exceeded those found *in vivo* following normal therapeutic doses. Testosterone, 17α-ethinyl estradiol, retinoic acid, and quercetin, a specific inhibitor of CYP2C8, also inhibited the formation of 6α-hydroxypaclitaxel *in vitro*. The pharmacokinetics of paclitaxel may also be altered *in vivo* as a result of interactions with compounds that are substrates, inducers, or inhibitors of CYP2C8 and/or CYP3A4. (See **PRECAUTIONS: Drug Interactions**.) The effect of renal or hepatic dysfunction on the disposition of paclitaxel has not been investigated.

Possible interactions of paclitaxel with concomitantly administered medications have not been formally investigated.

## CLINICAL STUDIES

### Ovarian Carcinoma:

**First-Line Data:** The safety and efficacy of TAXOL followed by cisplatin in patients with advanced ovarian cancer and no prior chemotherapy were evaluated in two Phase 3 multicenter, randomized, controlled trials. In an Intergroup study led by the European Organization for Research and Treatment of Cancer involving the Scandinavian Group NOCOVA, the National Cancer Institute of Canada, and the Scottish Group, 680 patients with Stage II<sub>b</sub>, III, or IV disease (optimally or non-optimally debulked) received either TAXOL 175 mg/m<sup>2</sup> infused over 3 hours followed by cisplatin 75 mg/m<sup>2</sup> (Tc) or cyclophosphamide 750 mg/m<sup>2</sup> followed by cisplatin 75 mg/m<sup>2</sup> (Cc) for a median of six courses. Although the protocol allowed further therapy, only 15% received both drugs for nine or more courses. In a study conducted by the Gynecological Oncology Group (GOG), 410 patients with Stage III or IV disease (>1 cm residual disease after staging laparotomy or distant metastases) received either TAXOL 135 mg/m<sup>2</sup> infused over 24 hours followed by cisplatin 75 mg/m<sup>2</sup> or cyclophosphamide 750 mg/m<sup>2</sup> followed by cisplatin 75 mg/m<sup>2</sup> for six courses.

In both studies, patients treated with TAXOL in combination with cisplatin had significantly higher response rate, longer time to progression, and longer survival time compared with standard therapy. These differences were also significant for the subset of patients in the Intergroup study with non-optimally debulked disease, although the study was not fully powered for subset analyses (Tables 2A and 2B). Kaplan-Meier survival curves for each study are shown in Figures 1 and 2.

Table 2A: Efficacy in the Phase 3 First-Line Ovarian Carcinoma Studies

	Intergroup (non-optimally debulked subset)		GOG-111	
	T175/3 <sup>a</sup> C75 <sup>b</sup> (n=218)	C750 <sup>b</sup> C75 (n=227)	T135/24 <sup>a</sup> C75 <sup>b</sup> (n=196)	C750 <sup>b</sup> C75 (n=214)
• <b>Clinical Response<sup>c</sup></b>	(n=153)	(n=153)	(n=113)	(n=127)
- rate (percent)	58	43	62	48
- p-value <sup>e</sup>	0.016		0.04	
• <b>Time to Progression</b>				
- median (months)	13.2	9.9	16.6	13.0
- p-value <sup>e</sup>	0.0060		0.0008	
- hazard ratio (HR) <sup>f</sup>	0.76		0.70	
- 95% CI <sup>f</sup>	0.62-0.92		0.56-0.86	
• <b>Survival</b>				
- median (months)	29.5	21.9	35.5	24.2
- p-value <sup>e</sup>	0.0057		0.0002	
- hazard ratio <sup>f</sup>	0.73		0.64	
- 95% CI <sup>f</sup>	0.58-0.91		0.50-0.81	

<sup>a</sup> TAXOL dose in mg/m<sup>2</sup>/infusion duration in hours; cyclophosphamide and cisplatin doses in mg/m<sup>2</sup>.  
<sup>b</sup> Among patients with measurable disease only.  
<sup>c</sup> Unstratified for the Intergroup Study. Stratified for Study GOG-111.

Table 2B: Efficacy in the Phase 3 First-Line Ovarian Carcinoma Intergroup Study

	T175/3 <sup>a</sup> C75 (n=342)	C750 <sup>b</sup> C75 (n=338)
• <b>Clinical Response<sup>c</sup></b>	(n=162)	(n=161)
- rate (percent)	59	45
- p-value <sup>e</sup>	0.014	
• <b>Time to Progression</b>		
- median (months)	15.3	11.5
- p-value <sup>e</sup>	0.0005	
- hazard ratio <sup>f</sup>	0.74	
- 95% CI <sup>f</sup>	0.63-0.88	
• <b>Survival</b>		
- median (months)	35.6	25.9
- p-value <sup>e</sup>	0.0016	
- hazard ratio <sup>f</sup>	0.73	
- 95% CI <sup>f</sup>	0.60-0.89	

<sup>a</sup> TAXOL dose in mg/m<sup>2</sup>/infusion duration in hours; cyclophosphamide and cisplatin doses in mg/m<sup>2</sup>.  
<sup>b</sup> Among patients with measurable disease only.  
<sup>c</sup> Unstratified.

Figure 1: Survival: Cc Versus Tc (Intergroup)





Figure 2: Survival: Cc Versus Tc (GOG-111)



The adverse event profile for patients receiving TAXOL (paclitaxel) Injection in combination with cisplatin in these studies was qualitatively consistent with that seen for the pooled analysis of data from 812 patients treated with single-agent TAXOL in 10 clinical studies. These adverse events and adverse events from the Phase 3 first-line ovarian carcinoma studies are described in the **ADVERSE REACTIONS** section in tabular (Tables 9 and 10) and narrative form.

**Second-Line Data:** Data from five Phase 1 & 2 clinical studies (189 patients), a multicenter randomized Phase 3 study (407 patients), as well as an interim referral center program were used in support of the use of TAXOL in patients who have failed initial or subsequent chemotherapy for metastatic carcinoma of the ovary. Two of the Phase 2 studies (92 patients) utilized an initial dose of 135 to 170 mg/m<sup>2</sup> in most patients (>90%) administered over 24 hours by continuous infusion. Response rates in these two studies were 22% (95% CI: 11% to 37%) and 30% (95% CI: 18% to 46%) with a total of 6 complete and 18 partial responses in 92 patients. The median duration of overall response in these two studies measured from the first day of treatment was 7.2 months (range: 3.5-15.8 months) and 7.5 months (range: 5.3-17.4 months), respectively. The median survival was 8.1 months (range: 0.2-36.7 months) and 15.9 months (range: 1.8-34.5 months).

The Phase 3 study had a bifactorial design and compared the efficacy and safety of TAXOL administered at two different doses (135 or 175 mg/m<sup>2</sup>) and schedules (3- or 24-hour infusion). The overall response rate for the 407 patients was 16.2% (95% CI: 12.8% to 20.2%), with 6 complete and 60 partial responses. Duration of response, measured from the first day of treatment was 8.3 months (range: 3.2-21.6 months). Median time to progression was 3.7 months (range: 0.1+ - 25.1+ months). Median survival was 11.5 months (range: 0.2-26.3+ months).

Response rates, median survival, and median time to progression for the 4 arms are given in the following table.

Table 3: Efficacy in the Phase 3 Second-Line Ovarian Carcinoma Study

	175/3 (n=96)	175/24 (n=106)	135/3 (n=99)	135/24 (n=106)
• <b>Response</b>				
- rate (percent)	14.6	21.7	15.2	13.2
- 95% Confidence Interval	(8.5-23.6)	(14.5-31.0)	(9.0-24.1)	(7.7-21.5)
• <b>Time to Progression</b>				
- median (months)	4.4	4.2	3.4	2.8
- 95% Confidence Interval	(3.0-5.6)	(3.5-5.1)	(2.8-4.2)	(1.9-4.0)
• <b>Survival</b>				
- median (months)	11.5	11.8	13.1	10.7
- 95% Confidence Interval	(8.4-14.4)	(8.9-14.6)	(9.1-14.6)	(8.1-13.6)

Analyses were performed as planned by the bifactorial study design described in the protocol, by comparing the two doses (135 or 175 mg/m<sup>2</sup>) irrespective of the schedule (3 or 24 hours) and the two schedules irrespective of dose. Patients receiving the 175 mg/m<sup>2</sup> dose had a response rate similar to that for those receiving the 135 mg/m<sup>2</sup> dose: 18% vs. 14% (p=0.28). No difference in response rate was detected when comparing the 3-hour with the 24-hour infusion: 15% vs. 17% (p=0.50). Patients receiving the 175 mg/m<sup>2</sup> dose of TAXOL had a longer time to progression than those receiving the 135 mg/m<sup>2</sup> dose: median 4.2 vs. 3.1 months (p=0.03). The median time to progression for patients receiving the 3-hour vs. the 24-hour infusion was 4.0 months vs. 3.7 months, respectively. Median survival was 11.6 months in patients receiving the 175 mg/m<sup>2</sup> dose of TAXOL and 11.0 months in patients receiving the 135 mg/m<sup>2</sup> dose (p=0.92). Median survival was 11.7 months for patients receiving the 3-hour infusion of TAXOL and 11.2 months for patients receiving the 24-hour infusion (p=0.91). These statistical analyses should be viewed with caution because of the multiple comparisons made.

TAXOL remained active in patients who had developed resistance to platinum-containing therapy (defined as tumor progression while on, or tumor relapse within 6 months from completion of, a platinum-containing regimen) with response rates of 14% in the Phase 3 study and 31% in the Phase 1 & 2 clinical studies.

The adverse event profile in this Phase 3 study was consistent with that seen for the pooled analysis of data from 812 patients treated in 10 clinical studies. These adverse events and adverse events from the Phase 3 second-line ovarian carcinoma study are described in the **ADVERSE REACTIONS** section in tabular (Tables 9 and 11) and narrative form.

The results of this randomized study support the use of TAXOL at doses of 135 to 175 mg/m<sup>2</sup>, administered by a 3-hour intravenous infusion. The same doses administered by 24-hour infusion were more toxic. However, the study had insufficient power to determine whether a particular dose and schedule produced superior efficacy.

**Breast Carcinoma:**

**Adjuvant Therapy:** A Phase 3 intergroup study (Cancer and Leukemia Group B [CALGB], Eastern Cooperative Oncology Group [ECOG], North Central Cancer Treatment Group [NCCTG], and Southwest Oncology Group [SWOG]) randomized 3170 patients with node-positive breast carcinoma to adjuvant therapy with TAXOL (paclitaxel) injection or to no further chemotherapy following four courses of doxorubicin and cyclophosphamide (AC). This multicenter trial was conducted in women with histologically positive lymph nodes following either a mastectomy or segmental mastectomy and nodal dissections. The 3 x 2 factorial study was designed to assess the efficacy and safety of three different dose levels of doxorubicin (A) and to evaluate the effect of the addition of TAXOL administered following the completion of AC therapy. After stratification for the number of positive lymph nodes (1-3, 4-9, or 10+), patients were randomized to receive cyclophosphamide at a dose of 600 mg/m<sup>2</sup> and doxorubicin at doses of either 60 mg/m<sup>2</sup> (on day 1), 75 mg/m<sup>2</sup> (in two divided doses on days 1 and 2), or 90 mg/m<sup>2</sup> (in two divided doses on days 1 and 2 with prophylactic G-CSF support and ciprofloxacin) every 3 weeks for four courses and either TAXOL 175 mg/m<sup>2</sup> as a 3-hour infusion every 3 weeks for four additional courses or no additional chemotherapy. Patients whose tumors were positive were to receive subsequent tamoxifen treatment (20 mg daily for 5 years); patients who received segmental mastectomies prior to study were to receive breast irradiation after recovery from treatment-related toxicities.

At the time of the current analysis, median follow-up was 30.1 months. Of the 2066 patients who were hormone receptor positive, 93% received tamoxifen. The primary analyses of disease-free survival and overall survival used multivariate Cox models, which included TAXOL administration, doxorubicin dose, number of positive lymph nodes, tumor size, menopausal status, and estrogen receptor status as factors. Based on the model for disease-free survival, patients receiving AC followed by TAXOL had a 22% reduction in the risk of disease recurrence compared to patients randomized to AC alone (Hazard Ratio [HR] = 0.78, 95% CI 0.67-0.91, p=0.0022). They also had a 26% reduction in the risk of death (HR = 0.74, 95% CI 0.60-0.92, p=0.0065). For disease-free survival and overall survival, p values were not adjusted for interim analyses. Kaplan-Meier curves are shown in Figures 3 and 4. Increasing the dose of doxorubicin higher than 60 mg/m<sup>2</sup> had no effect on either disease-free survival or overall survival.

Figure 3. Disease-Free Survival: AC Versus AC+T





Figure 4. Survival: AC Versus AC+T



**Subset analyses:** Subsets defined by variables of known prognostic importance in adjuvant breast carcinoma were examined, including number of positive lymph nodes, tumor size, hormone receptor status, and menopausal status. Such analyses must be interpreted with care, as the most secure finding is the overall study result. In general, a reduction in hazard similar to the overall reduction was seen with TAXOL for both disease-free and overall survival in all of the larger subsets with one exception; patients with receptor-positive tumors had a smaller reduction in hazard (HR = 0.92) for disease-free survival with TAXOL than other groups. Results of subset analyses are shown in Table 4.

Patient Subset	Disease-Free Survival		Overall Survival	
	No. of Patients	No. of Recurrences (95% CI)	No. of Deaths (95% CI)	Hazard Ratio (95% CI)
<b>No. of Positive Nodes</b>				
1-3	1449	221 (0.72) (0.55-0.94)	187 (0.76) (0.52-1.12)	
4-9	1310	274 (0.78) (0.61-0.99)	148 (0.66) (0.47-0.91)	
10+	360	129 (0.93) (0.66-1.31)	87 (0.59) (0.39-0.86)	
<b>Tumor Size (cm)</b>				
≤ 2	1096	153 (0.79) (0.57-1.08)	67 (0.73) (0.45-1.18)	
> 2 and ≤ 5	1611	358 (0.79) (0.64-0.97)	201 (0.74) (0.56-0.98)	
> 5	397	111 (0.75) (0.51-1.08)	72 (0.73) (0.46-1.16)	
<b>Menopausal Status</b>				
Pre	1929	374 (0.83) (0.67-1.01)	187 (0.72) (0.54-0.97)	
Post	1183	250 (0.73) (0.57-0.93)	155 (0.77) (0.56-1.06)	
<b>Receptor Status</b>				
Positive	2066	293 (0.92) (0.73-1.16)	126 (0.83) (0.59-1.18)	
Negative/Unknown <sup>a</sup>	1055	331 (0.68) (0.55-0.85)	216 (0.71) (0.54-0.93)	

<sup>a</sup>Positive for either estrogen or progesterone receptors.  
<sup>b</sup>Negative or missing for both estrogen and progesterone receptors (both missing: n=15).

These retrospective subgroup analyses suggest that the beneficial effect of TAXOL (paclitaxel) injection is clearly established in the receptor-negative subgroup, but the benefit in receptor-positive patients is not yet clear. With respect to menopausal status, the benefit of TAXOL is consistent (see Table 4 and Figures 5-8).

Figure 5. Disease-Free Survival - Receptor Status Negative/Unknown AC Versus AC+T




Figure 6. Disease-Free Survival - Receptor Status Positive AC Versus AC+T




Figure 7. Disease-Free Survival - Premenopausal AC Versus AC+T





Figure 8. Disease-Free Survival - Postmenopausal AC Versus AC+T



The adverse event profile for the patients who received TAXOL subsequent to AC was consistent with that seen in the pooled analysis of data from 812 patients (Table 9) treated with single-agent TAXOL in 10 clinical studies. These adverse events are described in the ADVERSE REACTIONS section in tabular (Tables 9 and 12) and narrative form.

**After Failure of Initial Chemotherapy:** Data from 83 patients accrued in three Phase 2 open label studies and from 471 patients enrolled in a Phase 3 randomized study were available to support the use of TAXOL in patients with metastatic breast carcinoma.

**Phase 2 open label studies:** Two studies were conducted in 53 patients previously treated with a maximum of one prior chemotherapeutic regimen. TAXOL was administered in these two trials as a 24-hour infusion at initial doses of 250 mg/m<sup>2</sup> (with G-CSF support) or 200 mg/m<sup>2</sup>. The response rates were 57% (95% CI: 37% to 75%) and 52% (95% CI: 32% to 72%), respectively. The third Phase 2 study was conducted in extensively pretreated patients who had failed anthracycline therapy and who had received a minimum of two chemotherapy regimens for the treatment of metastatic disease. The dose of TAXOL was 200 mg/m<sup>2</sup> as a 24-hour infusion with G-CSF support. Nine of 30 patients achieved a partial response, for a response rate of 30% (95% CI: 15% to 50%).

**Phase 3 randomized study:** This multicenter trial was conducted in patients previously treated with one or two regimens of chemotherapy. Patients were randomized to receive TAXOL at a dose of either 175 mg/m<sup>2</sup> or 135 mg/m<sup>2</sup> given as a 3-hour infusion. In the 471 patients enrolled, 60% had symptomatic disease with impaired performance status at study entry, and 73% had visceral metastases. These patients had failed prior chemotherapy either in the adjuvant setting (30%), the metastatic setting (39%), or both (31%). Sixty-seven percent of the patients had been previously exposed to anthracyclines and 23% of them had disease considered resistant to this class of agents.

The overall response rate for the 454 evaluable patients was 26% (95% CI: 22% to 30%), with 17 complete and 99 partial responses. The median duration of response, measured from the first day of treatment, was 8.1 months (range: 3.4-18.1 months). Overall for the 471 patients, the median time to progression was 3.5 months (range: 0.03-17.1 months). Median survival was 11.7 months (range: 0-18.9 months). Response rates, median survival and median time to progression for the 2 arms are given in the following table.

	175/3 (n=235)	135/3 (n=236)
<b>Response</b>		
- rate (percent)	28	22
- p-value		0.135
<b>Time to Progression</b>		
- median (months)	4.2	3.0
- p-value		0.027
<b>Survival</b>		
- median (months)	11.7	10.5
- p-value		0.321

The adverse event profile of the patients who received single-agent TAXOL (paclitaxel) injection in the Phase 3 study was consistent with that seen for the pooled analysis of data from 812 patients treated in 10 clinical studies. These adverse events and adverse events from the Phase 3 breast carcinoma study are described in the ADVERSE REACTIONS section in tabular (Tables 9 and 13) and narrative form.

**Non-Small Cell Lung Carcinoma (NSCLC):** In a Phase 3 open label randomized study conducted by the ECOG, 599 patients were randomized to either TAXOL (T) 135 mg/m<sup>2</sup> as a 24-hour infusion in combination with cisplatin (c) 75 mg/m<sup>2</sup>, TAXOL (T) 250 mg/m<sup>2</sup> as a 24-hour infusion in combination with cisplatin (c) 75 mg/m<sup>2</sup> with G-CSF support, or cisplatin (c) 75 mg/m<sup>2</sup> on day 1, followed by etoposide (VP) 100 mg/m<sup>2</sup> on days 1, 2, and 3 (control).

Response rates, median time to progression, median survival, and one-year survival rates are given in the following table. The reported p-values have not been adjusted for multiple comparisons. There were statistically significant differences favoring each of the TAXOL plus cisplatin arms for response rate and time to tumor progression. There was no statistically significant difference in survival between either TAXOL plus cisplatin arm and the cisplatin plus etoposide arm.

	T135/24 c75 (n=198)	T250/24 c75 (n=201)	VP100* c75 (n=200)
<b>Response</b>			
- rate (percent)	25	23	12
- p-value <sup>b</sup>	0.001	<0.001	
<b>Time to Progression</b>			
- median (months)	4.3	4.9	2.7
- p-value <sup>b</sup>	0.05	0.004	
<b>Survival</b>			
- median (months)	9.3	10.0	7.4
- p-value <sup>b</sup>	0.12	0.08	
<b>One-Year Survival</b>			
- percent of patients	36	40	32

\* Etoposide (VP) 100 mg/m<sup>2</sup> was administered I.V. on days 1, 2 and 3.  
<sup>b</sup> Compared to cisplatin/etoposide.

In the ECOG study, the Functional Assessment of Cancer Therapy-Lung (FACT-L) questionnaire had seven subscales that measured subjective assessment of treatment. Of the seven, the Lung Cancer Specific Symptoms subscale favored the TAXOL 135 mg/m<sup>2</sup>/24 hour plus cisplatin arm compared to the cisplatin/etoposide arm. For all other factors, there was no difference in the treatment groups.

The adverse event profile for patients who received TAXOL in combination with cisplatin in this study was generally consistent with that seen for the pooled analysis of data from 812 patients treated with single-agent TAXOL in 10 clinical studies. These adverse events and adverse events from the Phase 3 first-line NSCLC study are described in the ADVERSE REACTIONS section in tabular (Tables 9 and 14) and narrative form.

**AIDS-Related Kaposi's Sarcoma:** Data from two Phase 2 open label studies support the use of TAXOL as second-line therapy in patients with AIDS-related Kaposi's sarcoma. Fifty-nine of the 85 patients enrolled in these studies had previously received systemic therapy, including interferon alpha (32%), DaunoXome<sup>®</sup> (31%), DOXIL<sup>®</sup> (2%), and doxorubicin containing chemotherapy (42%), with 64% having received prior anthracyclines. Eighty-five percent of the pretreated patients had progressed on, or could not tolerate, prior systemic therapy.

In Study CA139-174 patients received TAXOL at 135 mg/m<sup>2</sup> as a 3-hour infusion every 3 weeks (intended dose intensity 45 mg/m<sup>2</sup>/week). If no dose-limiting toxicity was observed, patients were to receive 155 mg/m<sup>2</sup> and 175 mg/m<sup>2</sup> in subsequent courses. Hematopoietic growth factors were not to be used initially. In Study CA139-281 patients received TAXOL at 100 mg/m<sup>2</sup> as a 3-hour infusion every 2 weeks (intended dose intensity 50 mg/m<sup>2</sup>/week). In this study patients could be receiving hematopoietic growth factors before the start of TAXOL therapy, or this support was to be initiated as indicated; the dose of TAXOL was not increased. The dose intensity of TAXOL used in this patient population was lower than the dose intensity recommended for other solid tumors.

All patients had widespread and poor-risk disease. Applying the ACTG staging criteria to patients with prior systemic therapy, 93% were poor risk for extent of disease (T<sub>1</sub>), 88% had a CD4 count <200 cells/mm<sup>3</sup> (I<sub>1</sub>), and 97% had poor risk considering their systemic illness (S<sub>1</sub>).

All patients in Study CA139-174 had a Karnofsky performance status of 80 or 90 at baseline; in Study CA139-281, there were 26 (46%) patients with a Karnofsky performance status of 70 or worse at baseline.

	Percent of Patients	
	Prior Systemic Therapy (n=59)	
Visceral ± edema ± oral ± cutaneous	42	
Edema or lymph nodes ± oral ± cutaneous	41	
Oral ± cutaneous	10	
Cutaneous only	7	

Although the planned dose intensity in the two studies was slightly different (45 mg/m<sup>2</sup>/week in Study CA139-174 and 50 mg/m<sup>2</sup>/week in Study CA139-281), delivered dose intensity was 38-39 mg/m<sup>2</sup>/week in both studies, with a similar range (20-24 to 51-61).

DaunoXome<sup>®</sup> is a registered trademark of NeXstar Pharmaceuticals, Inc. DOXIL<sup>®</sup> is a registered trademark of Sequus Pharmaceuticals, Inc.

**Efficacy** - The efficacy of TAXOL (paclitaxel) Injection was evaluated by assessing cutaneous tumor response according to the amended ACTG criteria and by seeking evidence of clinical benefit in patients in six domains of symptoms and/or conditions that are commonly related to AIDS-related Kaposi's sarcoma.

**Cutaneous Tumor Response (Amended ACTG Criteria)** - The objective response rate was 59% (95% CI: 46% to 72%)(35 of 59 patients) in patients with prior systemic therapy. Cutaneous responses were primarily defined as flattening of more than 50% of previously raised lesions.

Table 8: Overall Best Response (Amended ACTG Criteria)	
	Percent of Patients
	Prior Systemic Therapy
	(n=59)
Complete response	3
Partial response	56
Stable disease	29
Progression	8
Early death/toxicity	3

The median time to response was 8.1 weeks and the median duration of response measured from the first day of treatment was 10.4 months (95% CI: 7.0 to 11.0 months) for the patients who had previously received systemic therapy. The median time to progression was 6.2 months (95% CI: 4.6 to 8.7 months).

**Additional Clinical Benefit** - Most data on patient benefit were assessed retrospectively (plans for such analyses were not included in the study protocols). Nonetheless, clinical descriptions and photographs indicated clear benefit in some patients, including instances of improved pulmonary function in patients with pulmonary involvement, improved ambulation, resolution of ulcers, and decreased analgesic requirements in patients with KS involving the feet and resolution of facial lesions and edema in patients with KS involving the face, extremities, and genitalia.

**Safety** - The adverse event profile of TAXOL administered to patients with advanced HIV disease and poor-risk AIDS-related Kaposi's sarcoma was generally similar to that seen in a pooled analysis of data from 812 patients with solid tumors. These adverse events and adverse events from the Phase 2 second-line Kaposi's sarcoma studies are described in the **ADVERSE REACTIONS** section in tabular (Tables 9 and 15) and narrative form. In this immunosuppressed patient population, however, a lower dose intensity of TAXOL and supportive therapy including hematopoietic growth factors in patients with severe neutropenia are recommended. Patients with AIDS-related Kaposi's sarcoma may have more severe hematologic toxicities than patients with solid tumors.

#### INDICATIONS

TAXOL is indicated as first-line and subsequent therapy for the treatment of advanced carcinoma of the ovary. As first-line therapy, TAXOL is indicated in combination with cisplatin.

TAXOL is indicated for the adjuvant treatment of node-positive breast cancer administered sequentially to standard doxorubicin-containing combination chemotherapy. In the clinical trial, there was an overall favorable effect on disease-free and overall survival in the total population of patients with receptor-positive and receptor-negative tumors, but the benefit has been specifically demonstrated by available data (median follow-up 30 months) only in the patients with estrogen and progesterone receptor-negative tumors. (See **CLINICAL STUDIES: Breast Carcinoma**.)

TAXOL is indicated for the treatment of breast cancer after failure of combination chemotherapy for metastatic disease or relapse within 6 months of adjuvant chemotherapy. Prior therapy should have included an anthracycline unless clinically contraindicated.

TAXOL, in combination with cisplatin, is indicated for the first-line treatment of non-small cell lung cancer in patients who are not candidates for potentially curative surgery and/or radiation therapy.

TAXOL is indicated for the second-line treatment of AIDS-related Kaposi's sarcoma.

#### CONTRAINDICATIONS

TAXOL is contraindicated in patients who have a history of hypersensitivity reactions to TAXOL or other drugs formulated in Cremophor® EL (polyoxyethylated castor oil).

TAXOL should not be used in patients with solid tumors who have baseline neutrophil counts of <1500 cells/mm<sup>3</sup> or in patients with AIDS-related Kaposi's sarcoma with baseline neutrophil counts of <1000 cells/mm<sup>3</sup>.

#### WARNINGS

Anaphylaxis and severe hypersensitivity reactions characterized by dyspnea and hypotension requiring treatment, angioedema, and generalized urticaria have occurred in 2%-4% of patients receiving TAXOL in clinical trials. Fatal reactions have occurred in patients despite premedication. All patients should be pretreated with corticosteroids, diphenhydramine, and H<sub>2</sub> antagonists. (See **DOSE AND ADMINISTRATION**.) Patients who experience severe hypersensitivity reactions to TAXOL should not be rechallenged with the drug.

Bone marrow suppression (primarily neutropenia) is dose-dependent and is the dose-limiting toxicity. Neutrophil nadirs occurred at a median of 11 days. TAXOL should not be administered to patients with baseline neutrophil counts of less than 1500 cells/mm<sup>3</sup> (<1000 cells/mm<sup>3</sup> for patients with KS). Frequent monitoring of blood counts should be instituted during TAXOL treatment. Patients should not be re-treated with subsequent cycles of TAXOL until neutrophils recover to a level >1500 cells/mm<sup>3</sup> (>1000 cells/mm<sup>3</sup> for patients with KS) and platelets recover to a level >100,000 cells/mm<sup>3</sup>.

Severe conduction abnormalities have been documented in <1% of patients during TAXOL therapy and in some cases requiring pacemaker placement. If patients develop significant conduction abnormalities during TAXOL infusion, appropriate therapy should be administered and continuous cardiac monitoring should be performed during subsequent therapy with TAXOL.

**Pregnancy:** TAXOL can cause fetal harm when administered to a pregnant woman. Administration of paclitaxel during the period of organogenesis to rabbits at doses of 3.0 mg/kg/day (about 0.2 the daily maximum recommended human dose on a mg/m<sup>2</sup> basis) caused embryo- and fetotoxicity, as indicated by intrauterine mortality, increased resorptions, and increased fetal deaths. Maternal toxicity was also observed at this dose. No teratogenic effects were observed at 1.0 mg/kg/day (about 1/15 the daily maximum recommended human dose on a mg/m<sup>2</sup> basis); teratogenic potential could not be assessed at higher doses due to extensive fetal mortality.

There are no adequate and well-controlled studies in pregnant women. If TAXOL is used during pregnancy, or if the patient becomes pregnant while receiving this drug, the patient should be apprised of the potential hazard to the fetus. Women of childbearing potential should be advised to avoid becoming pregnant.

#### PRECAUTIONS

Contact of the undiluted concentrate with plasticized polyvinyl chloride (PVC) equipment or devices used to prepare solutions for infusion is not recommended. In order to minimize patient exposure to the plasticizer DEHP [di-(2-ethylhexyl)phthalate], which may be leached from PVC infusion bags or sets, diluted TAXOL (paclitaxel) Injection solutions should preferably be stored in bottles (glass, polypropylene) or plastic bags (polypropylene, polyolefin) and administered through polyethylene-lined administration sets.

TAXOL should be administered through an in-line filter with a micro-porous membrane not greater than 0.22 microns. Use of filter devices such as IVEX-2® filters which incorporate short inlet and outlet PVC-coated tubing has not resulted in significant leaching of DEHP.

**Drug Interactions:** In a Phase I trial using escalating doses of TAXOL (110-200 mg/m<sup>2</sup>) and cisplatin (50 or 75 mg/m<sup>2</sup>) given as sequential infusions, myelosuppression was more profound when TAXOL was given after cisplatin than with the alternate sequence (i.e., TAXOL before cisplatin). Pharmacokinetic data from these patients demonstrated a decrease in paclitaxel clearance of approximately 33% when TAXOL was administered following cisplatin.

The metabolism of TAXOL is catalyzed by cytochrome P450 isoenzymes CYP2C8 and CYP3A4. In the absence of formal clinical drug interaction studies, caution should be exercised when administering TAXOL concomitantly with known substrates or inhibitors of the cytochrome P450 isoenzymes CYP2C8 and CYP3A4. (See **CLINICAL PHARMACOLOGY**.)

Potential interactions between TAXOL, a substrate of CYP3A4, and protease inhibitors (ritonavir, saquinavir, indinavir, and nelfinavir), which are substrates and/or inhibitors of CYP3A4, have not been evaluated in clinical trials.

Reports in the literature suggest that plasma levels of doxorubicin (and its active metabolite doxorubicinol) may be increased when paclitaxel and doxorubicin are used in combination.

**Hematology:** TAXOL therapy should not be administered to patients with baseline neutrophil counts of less than 1,500 cells/mm<sup>3</sup>. In order to monitor the occurrence of myelotoxicity, it is recommended that frequent peripheral blood cell counts be performed on all patients receiving TAXOL. Patients should not be re-treated with subsequent cycles of TAXOL until neutrophils recover to a level >1500 cells/mm<sup>3</sup> and platelets recover to a level >100,000 cells/mm<sup>3</sup>. In the case of severe neutropenia (<500 cells/mm<sup>3</sup> for seven days or more) during a course of TAXOL therapy, a 20% reduction in dose for subsequent courses of therapy is recommended.

For patients with advanced HIV disease and poor-risk AIDS-related Kaposi's sarcoma, TAXOL, at the recommended dose for this disease, can be initiated and repeated if the neutrophil count is at least 1000 cells/mm<sup>3</sup>.

**Hypersensitivity Reactions:** Patients with a history of severe hypersensitivity reactions to products containing Cremophor® EL (e.g., cyclosporin for injection concentrate and teniposide for injection concentrate) should not be treated with TAXOL. In order to avoid the occurrence of severe hypersensitivity reactions, all patients treated with TAXOL should be premedicated with corticosteroids (such as dexamethasone), diphenhydramine and H<sub>2</sub> antagonists (such as cimetidine or ranitidine). Minor symptoms such as flushing, skin reactions, dyspnea, hypotension, or tachycardia do not require interruption of therapy. However, severe reactions, such as hypotension requiring treatment, dyspnea requiring bronchodilators, angioedema, or generalized urticaria require immediate discontinuation of TAXOL and aggressive symptomatic therapy. Patients who have developed severe hypersensitivity reactions should not be rechallenged with TAXOL.

**Cardiovascular:** Hypotension, bradycardia, and hypertension have been observed during administration of TAXOL, but generally do not require treatment. Occasionally TAXOL infusions must be interrupted or discontinued because of initial or recurrent hypertension. Frequent vital sign monitoring, particularly during the first hour of TAXOL infusion, is recommended. Continuous cardiac monitoring is not required except for patients with serious conduction abnormalities. (See **WARNINGS**.)

**Nervous System:** Although the occurrence of peripheral neuropathy is frequent, the development of severe symptomatology is unusual and requires a dose reduction of 20% for all subsequent courses of TAXOL.

TAXOL contains dehydrated alcohol USP, 396 mg/mL; consideration should be given to possible CNS and other effects of alcohol. (See **PRECAUTIONS: Pediatric Use**.)

**Hepatic:** There is evidence that the toxicity of TAXOL is enhanced in patients with elevated liver enzymes. Caution should be exercised when administering TAXOL to patients with moderate to severe hepatic impairment and dose adjustments should be considered.

**Injection Site Reaction:** Injection site reactions, including reactions secondary to extravasation, were usually mild and consisted of erythema, tenderness, skin discoloration, or swelling at the injection site. These reactions have been observed more frequently with the 24-hour infusion than with the 3-hour infusion. Recurrence of skin reactions at a site of previous extravasation following administration of TAXOL at a different site, i.e., "recall", has been reported rarely.

Rare reports of more severe events such as phlebitis, cellulitis, induration, skin exfoliation, necrosis, and fibrosis have been received as part of the continuing surveillance of TAXOL safety. In some cases the onset of the injection site reaction either occurred during a prolonged infusion or was delayed by a week to ten days.

A specific treatment for extravasation reactions is unknown at this time. Given the possibility of extravasation, it is advisable to closely monitor the infusion site for possible infiltration during drug administration.

**Carcinogenesis, Mutagenesis, Impairment of Fertility:** The carcinogenic potential of TAXOL has not been studied.

Paclitaxel has been shown to be clastogenic *in vitro* (chromosome aberrations in human lymphocytes) and *in vivo* (micronucleus test in mice). Paclitaxel was not mutagenic in the Ames test or the CHO/HGPRT gene mutation assay.

Administration of paclitaxel prior to and during mating produced impairment of fertility in male and female rats at doses equal to or greater than 1 mg/kg/day (about 0.04 the daily maximum recommended human dose on a mg/m<sup>2</sup> basis). At this dose, paclitaxel caused reduced fertility and reproductive indices, and increased embryo- and fetotoxicity. (See **WARNINGS**.)

**Pregnancy:** Pregnancy "Category D". (See **WARNINGS**.)

**Nursing Mothers:** It is not known whether the drug is excreted in human milk. Following intravenous administration of carbon-14 labeled TAXOL to rats on days 9 to 10 postpartum, concentrations of radioactivity in milk were higher than in plasma and declined in parallel with the plasma concentrations. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants, it is recommended that nursing be discontinued when receiving TAXOL therapy.

**Pediatric Use:** The safety and effectiveness of TAXOL in pediatric patients have not been established.

There have been reports of central nervous system (CNS) toxicity (rarely associated with death) in a clinical trial in pediatric patients in which TAXOL was infused intravenously over 3 hours at doses ranging from 350 mg/m<sup>2</sup> to 420 mg/m<sup>2</sup>. The toxicity is most likely attributable to the

high dose of the ethanol component of the TAXOL (paclitaxel) Injection vehicle given over a short infusion time. The use of concomitant antihistamines may intensify this effect. Although a direct effect of the paclitaxel itself cannot be discounted, the high doses used in this study (over twice the recommended adult dosage) must be considered in assessing the safety of TAXOL for use in this population.

**Information for Patients:** (See Patient Information Leaflet.)

#### ADVERSE REACTIONS

**Pooled Analysis of Adverse Event Experiences from Single-Agent Studies:** Data in the following table are based on the experience of 812 patients (493 with ovarian carcinoma and 319 with breast carcinoma) enrolled in 10 studies who received single-agent TAXOL. Two hundred and seventy-five patients were treated in eight Phase 2 studies with TAXOL doses ranging from 135 to 300 mg/m<sup>2</sup> administered over 24 hours (in four of these studies, G-CSF was administered as hematopoietic support). Three hundred and one patients were treated in the randomized Phase 3 ovarian carcinoma study which compared two doses (135 or 175 mg/m<sup>2</sup>) and two schedules (3 or 24 hours) of TAXOL. Two hundred and thirty-six patients with breast carcinoma received TAXOL (135 or 175 mg/m<sup>2</sup>) administered over 3 hours in a controlled study.

Table 9: Summary* of Adverse Events in Patients With Solid Tumors Receiving Single-Agent TAXOL	
	Percent of Patients (n=812)
• <b>Bone Marrow</b>	
- Neutropenia	< 2,000/mm <sup>3</sup> 90
	< 500/mm <sup>3</sup> 52
	< 4,000/mm <sup>3</sup> 90
	< 1,000/mm <sup>3</sup> 17
- Thrombocytopenia	< 100,000/mm <sup>3</sup> 20
	< 50,000/mm <sup>3</sup> 7
- Anemia	< 11 g/dL 78
	< 8 g/dL 16
- Infections	30
- Bleeding	14
- Red Cell Transfusions	25
- Platelet Transfusions	2
• <b>Hypersensitivity Reaction*</b>	
- All	41
- Severe†	2
• <b>Cardiovascular</b>	
- Vital Sign Changes*	
- Bradycardia (n=537)	3
- Hypotension (n=532)	12
- Significant Cardiovascular Events	1
• <b>Abnormal ECG</b>	
- All Pts	23
- Pts with normal baseline (n=559)	14
• <b>Peripheral Neuropathy</b>	
- Any symptoms	60
- Severe symptoms†	3
• <b>Myalgia/Arthralgia</b>	
- Any symptoms	60
- Severe symptoms†	8
• <b>Gastrointestinal</b>	
- Nausea and vomiting	52
- Diarrhea	38
- Mucositis	31
• <b>Atopeia</b>	87
• <b>Hepatic (Pts with normal baseline and on study data)</b>	
- Bilirubin elevations (N=765)	7
- Alkaline phosphatase elevations (N=575)	22
- AST (SGOT) elevations (N=591)	19
• <b>Injection Site Reaction</b>	13

\* Based on worst course analysis.

† All patients received premedication.

‡ During the first 3 hours of infusion.

§ Severe events are defined as at least Grade III toxicity.

None of the observed toxicities were clearly influenced by age. **Disease-Specific Adverse Event Experiences First-Line Ovary in Combination:** For the 1084 patients who were evaluable for safety in the Phase 3 first-line ovary combination therapy studies, Table 10 shows the incidence of important adverse events. For both studies, the analysis of safety was based on all courses of therapy (six courses for the GOG-111 study and up to nine courses for the Intergroup study).

Table 10: Frequency* of Important Adverse Events in the Phase 3 First-Line Ovarian Carcinoma Studies				
	Percent of Patients			
	Intergroup	GOG-111		
	T175/3 <sup>b</sup> c75 <sup>c</sup> (n=339)	C750 <sup>c</sup> c75 <sup>c</sup> (n=336)	T135/24 <sup>b</sup> c75 <sup>c</sup> (n=196)	C750 <sup>c</sup> c75 <sup>c</sup> (n=213)
• <b>Bone Marrow</b>				
- Neutropenia				
< 2000/mm <sup>3</sup>	91 <sup>d</sup>	95 <sup>d</sup>	96	92
< 500/mm <sup>3</sup>	33 <sup>d</sup>	43 <sup>d</sup>	81 <sup>d</sup>	58 <sup>d</sup>
- Thrombocytopenia				
< 100,000/mm <sup>3</sup>	21 <sup>d</sup>	33 <sup>d</sup>	26	30
< 50,000/mm <sup>3</sup>	3 <sup>d</sup>	7 <sup>d</sup>	10	9
- Anemia				
< 11 g/dL <sup>e</sup>	96	97	88	86
< 8 g/dL	3 <sup>d</sup>	8 <sup>d</sup>	13	9
- Infections	25	27	21	15
- Febrile Neutropenia	4	7	15 <sup>d</sup>	4 <sup>d</sup>
• <b>Hypersensitivity Reaction</b>				
- All	41 <sup>d</sup>	6 <sup>d</sup>	8 <sup>4a</sup>	1 <sup>4a</sup>
- Severe†	1	1	3 <sup>4a</sup>	— <sup>4a</sup>

\* Based on worst course analysis.

<sup>b</sup> TAXOL (T) dose in mg/m<sup>2</sup>/infusion duration in hours.

<sup>c</sup> Cyclophosphamide (C) or cisplatin (c) dose in mg/m<sup>2</sup>.

<sup>d</sup> p<0.05 by Fisher exact test.

<sup>e</sup> <130,000/mm<sup>3</sup> in the Intergroup study.

<sup>f</sup> <12 g/dL in the Intergroup study.

<sup>g</sup> All patients received premedication.

<sup>h</sup> Severe events are defined as at least Grade III toxicity.

**Table 10: Frequency\* of Important Adverse Events in the Phase 3 First-Line Ovarian Carcinoma Studies (cont.)**

	Percent of Patients			
	Intergroup		GOG-111	
	T175/3 <sup>a</sup> c75 <sup>a</sup> (n=339)	C750 <sup>a</sup> c75 <sup>a</sup> (n=336)	T135/24 <sup>a</sup> c75 <sup>a</sup> (n=196)	C750 <sup>a</sup> c75 <sup>a</sup> (n=213)
<b>• Neurotoxicity<sup>b</sup></b>				
- Any symptoms	87 <sup>d</sup>	52 <sup>d</sup>	25	20
- Severe symptoms <sup>c</sup>	21 <sup>e</sup>	2 <sup>e</sup>	3 <sup>e</sup>	— <sup>d</sup>
<b>• Nausea and Vomiting</b>				
- Any symptoms	88	93	65	69
- Severe symptoms <sup>c</sup>	18	24	10	11
<b>• Myalgia/Arthralgia</b>				
- Any symptoms	60 <sup>d</sup>	27 <sup>d</sup>	9 <sup>d</sup>	2 <sup>d</sup>
- Severe symptoms <sup>c</sup>	6 <sup>d</sup>	1 <sup>d</sup>	1	—
<b>• Diarrhea</b>				
- Any symptoms	37 <sup>d</sup>	29 <sup>d</sup>	16 <sup>d</sup>	8 <sup>d</sup>
- Severe symptoms <sup>c</sup>	2	3	4	1
<b>• Asthenia</b>				
- Any symptoms	NC	NC	17 <sup>d</sup>	10 <sup>d</sup>
- Severe symptoms <sup>c</sup>	NC	NC	1	1
<b>• Alopecia</b>				
- Any symptoms	96 <sup>d</sup>	89 <sup>d</sup>	55 <sup>d</sup>	37 <sup>d</sup>
- Severe symptoms <sup>c</sup>	51 <sup>d</sup>	21 <sup>d</sup>	6	8

<sup>a</sup> Based on worst course analysis.  
<sup>b</sup> TAXOL (T) dose in mg/m<sup>2</sup>/infusion duration in hours.  
<sup>c</sup> Cyclophosphamide (C) or cisplatin (c) dose in mg/m<sup>2</sup>.  
<sup>d</sup> p<0.05 by Fisher exact test.  
<sup>e</sup> <130,000/mm<sup>3</sup> in the intergroup study.  
<sup>f</sup> <12 g/dL in the intergroup study.  
<sup>g</sup> All patients received premedication.  
<sup>h</sup> In the GOG-111 study, neurotoxicity was collected as peripheral neuropathy and in the intergroup study, neurotoxicity was collected as either neuromotor or neurosensory symptoms.  
<sup>i</sup> Severe events are defined as at least Grade III toxicity.  
 NC Not Collected.

**Second-Line Ovary:** For the 403 patients who received single-agent TAXOL (paclitaxel) injection in the Phase 3 second-line ovarian carcinoma study, the following table shows the incidence of important adverse events.

**Table 11: Frequency\* of Important Adverse Events in the Phase 3 Second-Line Ovarian Carcinoma Study**

	Percent of Patients			
	175/3 <sup>a</sup> 175/24 <sup>a</sup> 135/3 <sup>a</sup> 135/24 <sup>a</sup>		135/24 <sup>a</sup> 135/24 <sup>a</sup>	
	(n=95)	(n=105)	(n=98)	(n=105)
<b>• Bone Marrow</b>				
- Neutropenia	< 2,000/mm <sup>3</sup>	78	98	78
	< 500/mm <sup>3</sup>	27	75	14
	< 100,000/mm <sup>3</sup>	4	18	8
	< 50,000/mm <sup>3</sup>	1	7	2
	< 11 g/dL	84	90	68
	< 8 g/dL	11	12	6
- Infections		26	29	20
<b>• Hypersensitivity Reaction<sup>b</sup></b>				
- All		41	45	38
- Severe <sup>c</sup>		2	0	2
<b>• Peripheral Neuropathy</b>				
- Any symptoms		63	60	55
- Severe symptoms <sup>c</sup>		1	2	0
<b>• Mucositis</b>				
- Any symptoms		17	35	21
- Severe symptoms <sup>c</sup>		0	3	0

<sup>a</sup> Based on worst course analysis.  
<sup>b</sup> TAXOL dose in mg/m<sup>2</sup>/infusion duration in hours.  
<sup>c</sup> All patients received premedication.  
<sup>d</sup> Severe events are defined as at least Grade III toxicity.

Myelosuppression was dose and schedule related, with the schedule effect being more prominent. The development of severe hypersensitivity reactions (HSRs) was rare; 1% of the patients and 0.2% of the courses overall. There was no apparent dose or schedule effect seen for the HSRs. Peripheral neuropathy was clearly dose-related, but schedule did not appear to affect the incidence.

**Adjuvant Breast:** For the Phase 3 adjuvant breast carcinoma study, the following table shows the incidence of important severe adverse events for the 3121 patients (total population) who were evaluable for safety as well as for a group of 325 patients (early population) who, per the study protocol, were monitored more intensively than other patients.

**Table 12: Frequency\* of Important Severe<sup>b</sup> Adverse Events in the Phase 3 Adjuvant Breast Carcinoma Study**

	Percent of Patients			
	Early Population		Total Population	
	AC <sup>c</sup> (n=166)	AC <sup>c</sup> followed by T <sup>d</sup> (n=159)	AC <sup>c</sup> (n=1551)	AC <sup>c</sup> followed by T <sup>d</sup> (n=1570)
<b>• Bone Marrow<sup>a</sup></b>				
- Neutropenia				
	< 500/mm <sup>3</sup>	79	76	48
	< 100,000/mm <sup>3</sup>	27	25	11
- Anemia				
	< 8 g/dL	17	21	8
	< 11 g/dL	6	14	5
	< 8 g/dL	—	3	<1
- Fever without Infection		4	1	2
<b>• Hypersensitivity Reaction<sup>b</sup></b>				
- All		2	1	2
- Severe <sup>c</sup>		1	2	1
<b>• Neuromotor Toxicity</b>				
- Any symptoms		1	<1	1
- Severe symptoms <sup>c</sup>		—	—	3
<b>• Neurosensory Toxicity</b>				
- Any symptoms		1	<1	1
- Severe symptoms <sup>c</sup>		—	—	2
<b>• Myalgia/Arthralgia</b>				
- Any symptoms		13	18	8
- Severe symptoms <sup>c</sup>		4	6	5

<sup>a</sup> Based on worst course analysis.  
<sup>b</sup> Severe events are defined as at least Grade III toxicity.  
<sup>c</sup> Patients received 600 mg/m<sup>2</sup> cyclophosphamide and doxorubicin (AC) at doses of either 60 mg/m<sup>2</sup>, 75 mg/m<sup>2</sup>, or 90 mg/m<sup>2</sup> (with prophylactic G-CSF support and ciprofloxacin), every 3 weeks for four courses.  
<sup>d</sup> TAXOL (T) following four courses of AC at a dose of 175 mg/m<sup>2</sup>/3 hours every 3 weeks for four courses.  
<sup>e</sup> The incidence of febrile neutropenia was not reported in this study.  
<sup>f</sup> All patients were to receive premedication.

The incidence of an adverse event for the total population likely represents an underestimation of the actual incidence given that safety data were collected differentially based on enrollment cohort. However, since safety data were collected consistently across regimens, the safety of the sequential addition of TAXOL (paclitaxel) injection following AC therapy may be compared with AC therapy alone. Compared to patients who received AC alone, patients who received AC followed by TAXOL experienced more Grade III/IV neurosensory toxicity, more Grade III/IV myalgia/arthralgia, more Grade III/IV neurologic pain (5% vs 1%), more Grade III/IV flu-like symptoms (5% vs 3%), and more Grade III/IV hyperglycemia (3% vs 1%). During the additional four courses of treatment with TAXOL, two deaths (0.1%) were attributed to treatment. During TAXOL treatment, Grade IV neutropenia was reported for 15% of patients, Grade III/IV neurosensory toxicity for 15%, Grade III/IV myalgias for 23%, and alopecia for 46%.

The incidences of severe hematologic toxicities, infections, mucositis, and cardiovascular events increased with higher doses of doxorubicin. **Breast Cancer After Failure of Initial Chemotherapy:** For the 458 patients who received single-agent TAXOL in the Phase 3 breast carcinoma study, the following table shows the incidence of important adverse events by treatment arm (each arm was administered by a 3-hour infusion).

**Table 13: Frequency\* of Important Adverse Events in the Phase 3 Study of Breast Cancer After Failure of Initial Chemotherapy or Within 6 Months of Adjuvant Chemotherapy**

	Percent of Patients	
	175/3 <sup>a</sup> (n=229)	135/3 <sup>a</sup> (n=229)
	<b>• Bone Marrow</b>	
- Neutropenia	< 2,000/mm <sup>3</sup>	90
	< 500/mm <sup>3</sup>	28
- Thrombocytopenia	< 100,000/mm <sup>3</sup>	11
	< 50,000/mm <sup>3</sup>	3
- Anemia	< 11 g/dL	55
	< 8 g/dL	4
- Infections		23
- Febrile Neutropenia		2
<b>• Hypersensitivity Reaction<sup>b</sup></b>		
- All		36
- Severe <sup>c</sup>		0
<b>• Peripheral Neuropathy</b>		
- Any symptoms		70
- Severe symptoms <sup>c</sup>		7
<b>• Mucositis</b>		
- Any symptoms		23
- Severe symptoms <sup>c</sup>		3

<sup>a</sup> Based on worst course analysis.  
<sup>b</sup> TAXOL dose in mg/m<sup>2</sup>/infusion duration in hours.  
<sup>c</sup> All patients received premedication.  
<sup>d</sup> Severe events are defined as at least Grade III toxicity.

Myelosuppression and peripheral neuropathy were dose related. There was one severe hypersensitivity reaction (HSR) observed at the dose of 135 mg/m<sup>2</sup>. **First-Line NSCLC in Combination:** In the study conducted by the Eastern Cooperative Oncology Group (ECOG), patients were randomized to either TAXOL (T) 135 mg/m<sup>2</sup> as a 24-hour infusion in combination with cisplatin (c) 75 mg/m<sup>2</sup>, TAXOL (T) 250 mg/m<sup>2</sup> as a 24-hour infusion in combination with cisplatin (c) 75 mg/m<sup>2</sup> with G-CSF support, or cisplatin (c) 75 mg/m<sup>2</sup> on day 1, followed by etoposide (VP) 100 mg/m<sup>2</sup> on days 1, 2 and 3 (control). The following table shows the incidence of important adverse events.

**Table 14: Frequency\* of Important Adverse Events in the Phase 3 Study for First-Line NSCLC**

	Percent of Patients		
	T135/24 <sup>a</sup> c75 (n=195)	T250/24 <sup>a</sup> c75 (n=197)	VP100 <sup>a</sup> c75 (n=196)
	<b>• Bone Marrow</b>		
- Neutropenia	< 2,000/mm <sup>3</sup>	89	86
	< 500/mm <sup>3</sup>	74 <sup>e</sup>	65
- Thrombocytopenia	< normal	48	68
	< 50,000/mm <sup>3</sup>	6	12
- Anemia	< normal	94	96
	< 8 g/dL	22	19
- Infections		38	31
<b>• Hypersensitivity Reaction<sup>b</sup></b>			
- All		16	27
- Severe <sup>c</sup>		1	4 <sup>e</sup>
<b>• Arthralgia/Myalgia</b>			
- Any symptoms		21 <sup>e</sup>	42 <sup>e</sup>
- Severe symptoms <sup>c</sup>		3	11
<b>• Nausea/Vomiting</b>			
- Any symptoms		85	87
- Severe symptoms <sup>c</sup>		27	29
<b>• Mucositis</b>			
- Any symptoms		18	28
- Severe symptoms <sup>c</sup>		1	4
<b>• Neuromotor Toxicity</b>			
- Any symptoms		37	47
- Severe symptoms <sup>c</sup>		6	12
<b>• Neurosensory Toxicity</b>			
- Any symptoms		48	61
- Severe symptoms <sup>c</sup>		13	28 <sup>e</sup>
<b>• Cardiovascular Events</b>			
- Any symptoms		33	39
- Severe symptoms <sup>c</sup>		13	12

<sup>a</sup> Based on worst course analysis.  
<sup>b</sup> TAXOL (T) dose in mg/m<sup>2</sup>/infusion duration in hours; cisplatin (c) dose in mg/m<sup>2</sup>.  
<sup>c</sup> TAXOL dose in mg/m<sup>2</sup>/infusion duration in hours with G-CSF support; cisplatin dose in mg/m<sup>2</sup>.  
<sup>d</sup> Etoposide (VP) dose in mg/m<sup>2</sup> was administered I.V. on days 1, 2 and 3; cisplatin dose in mg/m<sup>2</sup>.  
<sup>e</sup> p<0.05.  
<sup>f</sup> All patients received premedication.  
<sup>g</sup> Severe events are defined as at least Grade III toxicity.

Toxicity was generally more severe in the high-dose TAXOL (paclitaxel) injection treatment arm (T250/c75) than in the low-dose TAXOL arm (T135/c75). Compared to the cisplatin/toposide arm, patients in the low-dose TAXOL arm experienced more arthralgia/myalgia of any grade and more severe neutropenia. The incidence of febrile neutropenia was not reported in this study.

**Kaposi's Sarcoma:** The following table shows the frequency of important adverse events in the 85 patients with KS treated with two different single-agent TAXOL regimens.

**Table 15: Frequency\* of Important Adverse Events in the AIDS-Related Kaposi's Sarcoma Studies**

	Percent of Patients	
	Study CA139-174 TAXOL 135/3 <sup>a</sup> q 3 wk (n=29)	Study CA139-281 TAXOL 100/3 <sup>a</sup> q 2 wk (n=56)
	<b>• Bone Marrow</b>	
- Neutropenia	< 2,000/mm <sup>3</sup>	100
	< 500/mm <sup>3</sup>	76
- Thrombocytopenia	< 100,000/mm <sup>3</sup>	52
	< 50,000/mm <sup>3</sup>	17
- Anemia	< 11 g/dL	86
	< 8 g/dL	34
- Febrile Neutropenia		55
<b>• Opportunistic Infection</b>		
- Any		76
- Cytomegalovirus		45
- Herpes Simplex		38
- <i>Pneumocystis carinii</i>		14
- <i>M. avium-intracellulare</i>		24
- Candidiasis, esophageal		7
- Cryptosporidiosis		7
- Cryptococcal meningitis		3
- Leukoencephalopathy		—
<b>• Hypersensitivity Reaction<sup>b</sup></b>		
- All		14
<b>• Cardiovascular</b>		
- Hypotension		17
- Bradycardia		3
<b>• Peripheral Neuropathy</b>		
- Any		79
- Severe <sup>c</sup>		10
<b>• Myalgia/Arthralgia</b>		
- Any		93
- Severe <sup>c</sup>		14
<b>• Gastrointestinal</b>		
- Nausea and Vomiting		69
- Diarrhea		90
- Mucositis		45
<b>• Renal (creatinine elevation)</b>		
- Any		34
- Severe <sup>c</sup>		7
<b>• Discontinuation for drug toxicity</b>		
- Any		34
- Severe <sup>c</sup>		7

<sup>a</sup> Based on worst course analysis.  
<sup>b</sup> TAXOL dose in mg/m<sup>2</sup>/infusion duration in hours.  
<sup>c</sup> All patients received premedication.  
<sup>d</sup> Severe events are defined as at least Grade III toxicity.

As demonstrated in this table, toxicity was more pronounced in the study utilizing TAXOL at a dose of 135 mg/m<sup>2</sup> every 3 weeks than in the study utilizing TAXOL at a dose of 100 mg/m<sup>2</sup> every 2 weeks. Notably, severe neutropenia (76% versus 35%), febrile neutropenia (55% versus 9%), and opportunistic infections (76% versus 54%) were more common with the former dose and schedule. The differences between the two studies with respect to dose escalation and use of hematopoietic growth factors, as described above, should be taken into account. (See CLINICAL STUDIES: AIDS-Related Kaposi's Sarcoma.) Note also that only 26% of the 85 patients in these studies received concomitant treatment with protease inhibitors, whose effect on paclitaxel metabolism has not yet been studied.

**Adverse Event Experiences by Body System:** Unless otherwise noted, the following discussion refers to the overall safety database of 812 patients with solid tumors treated with single-agent TAXOL in clinical studies. Toxicities that occurred with greater severity or frequency in previously untreated patients with ovarian carcinoma or NSCLC who received TAXOL in combination with cisplatin or in patients with breast cancer who received TAXOL after doxorubicin/cyclophosphamide in the adjuvant setting and that occurred with a difference that was clinically significant in these populations are also described. The frequency and severity of important adverse events for the Phase 3 ovarian carcinoma, breast carcinoma, NSCLC, and the Phase 2 Kaposi's sarcoma studies are presented above in tabular form by treatment arm. In addition, rare events have been reported from postmarketing experience or from other clinical studies. The frequency and severity of adverse events have been generally similar for patients receiving TAXOL for the treatment of ovarian, breast, or lung carcinoma or Kaposi's sarcoma, but patients with AIDS-related Kaposi's sarcoma may have more frequent and severe hematologic toxicity, infections, and febrile neutropenia. These patients require a lower dose intensity and supportive care. (See CLINICAL STUDIES: AIDS-Related Kaposi's Sarcoma.) Toxicities that were observed only in or were noted to have occurred with greater severity in the population with Kaposi's sarcoma and that occurred with a difference that was clinically significant in this population are described.

**Hematologic:** Bone marrow suppression was the major dose-limiting toxicity of TAXOL. Neutropenia, the most important hematologic toxicity, was dose and schedule dependent and was generally rapidly reversible. Among patients treated in the Phase 3 second-line ovarian study with a 3-hour infusion, neutrophil counts declined below 500 cells/mm<sup>3</sup> in 14% of the patients treated with a dose of 135 mg/m<sup>2</sup> compared to 27% at a dose of 175 mg/m<sup>2</sup> (p=0.05). In the same study, severe neutropenia (<500 cells/mm<sup>3</sup>) was more frequent with the 24-hour than with the 3-hour infusion; infusion duration had a greater impact on myelosuppression than dose. Neutropenia did not appear to increase with cumulative exposure and did not appear to be more frequent nor more severe for patients previously treated with radiation therapy.

In the study where TAXOL was administered to patients with ovarian carcinoma at a dose of 135 mg/m<sup>2</sup>/24 hours in combination with cisplatin versus the control arm of cyclophosphamide plus cisplatin, the incidences of grade IV neutropenia and of febrile neutropenia were significantly greater in the TAXOL plus cisplatin arm than in the control arm. Grade

IV neutropenia occurred in 81% on the TAXOL (paclitaxel) Injection plus cisplatin arm versus 58% on the cyclophosphamide plus cisplatin arm, and febrile neutropenia occurred in 15% and 4% respectively. On the TAXOL/cisplatin arm, there were 35/1074 (3%) courses with fever in which Grade IV neutropenia was reported at some time during the course. When TAXOL followed by cisplatin was administered to patients with advanced NSCLC in the ECOG study, the incidences of Grade IV neutropenia were 74% (TAXOL 135 mg/m<sup>2</sup>/24 hours followed by cisplatin) and 65% (TAXOL 250 mg/m<sup>2</sup>/24 hours followed by cisplatin and G-CSF) compared with 55% in patients who received cisplatin/etoposide.

Fever was frequent (12% of all treatment courses). Infectious episodes occurred in 30% of all patients and 9% of all courses; these episodes were fatal in 1% of all patients, and included sepsis, pneumonia and peritonitis. In the Phase 3 second-line ovarian study, infectious episodes were reported in 20% and 26% of the patients treated with a dose of 135 mg/m<sup>2</sup> or 175 mg/m<sup>2</sup> given as 3-hour infusions, respectively. Urinary tract infections and upper respiratory tract infections were the most frequently reported infectious complications. In the immunosuppressed patient population with advanced HIV disease and poor-risk AIDS-related Kaposi's sarcoma, 61% of the patients reported at least one opportunistic infection. (See **CLINICAL STUDIES: AIDS-Related Kaposi's Sarcoma**.) The use of supportive therapy, including G-CSF, is recommended for patients who have experienced severe neutropenia. (See **DOSE AND ADMINISTRATION**.)

Thrombocytopenia was uncommon, and almost never severe (<50,000 cells/mm<sup>3</sup>). Twenty percent of the patients experienced a drop in their platelet count below 100,000 cells/mm<sup>3</sup> at least once while on treatment; 7% had a platelet count <50,000 cells/mm<sup>3</sup> at the time of their worst nadir. Bleeding episodes were reported in 4% of all courses and by 14% of all patients but most of the hemorrhagic episodes were localized and the frequency of these events was unrelated to the TAXOL dose and schedule. In the Phase 3 second-line ovarian study, bleeding episodes were reported in 10% of the patients; no patients treated with the 3-hour infusion received platelet transfusions. In the adjuvant breast carcinoma trial, the incidence of severe thrombocytopenia and platelet transfusions increased with higher doses of doxorubicin.

Anemia (Hb <11 g/dL) was observed in 78% of all patients and was severe (Hb <8 g/dL) in 16% of the cases. No consistent relationship between dose or schedule and the frequency of anemia was observed. Among all patients with normal baseline hemoglobin, 69% became anemic on study but only 7% had severe anemia. Red cell transfusions were required in 25% of all patients and in 12% of those with normal baseline hemoglobin levels.

**Hypersensitivity Reactions (HSRs):** All patients received premedication prior to TAXOL (see **WARNINGS** and **PRECAUTIONS: Hypersensitivity Reactions**). The frequency and severity of HSRs were not affected by the dose or schedule of TAXOL administration. In the Phase 3 second-line ovarian study, the 3-hour infusion was not associated with a greater increase in HSRs when compared to the 24-hour infusion. Hypersensitivity reactions were observed in 20% of all courses and in 41% of all patients. These reactions were severe in less than 2% of the patients and 1% of the courses. No severe reactions were observed after course 3 and severe symptoms occurred generally within the first hour of TAXOL infusion. The most frequent symptoms observed during these severe reactions were dyspnea, flushing, chest pain, and tachycardia.

The minor hypersensitivity reactions consisted mostly of flushing (28%), rash (12%), hypotension (4%), dyspnea (2%), tachycardia (2%), and hypertension (1%). The frequency of hypersensitivity reactions remained relatively stable during the entire treatment period.

Rare reports of chills and reports of back pain in association with hypersensitivity reactions have been received as part of the continuing surveillance of TAXOL safety.

**Cardiovascular:** Hypotension, during the first 3 hours of infusion, occurred in 12% of all patients and 3% of all courses administered. Bradycardia, during the first 3 hours of infusion, occurred in 3% of all patients and 1% of all courses. In the Phase 3 second-line ovarian study, neither dose nor schedule had an effect on the frequency of hypotension and bradycardia. These vital sign changes most often caused no symptoms and required neither specific therapy nor treatment discontinuation. The frequency of hypotension and bradycardia were not influenced by prior anthracycline therapy.

Significant cardiovascular events possibly related to single-agent TAXOL occurred in approximately 1% of all patients. These events included syncope, rhythm abnormalities, hypertension and venous thrombosis. One of the patients with syncope treated with TAXOL at 175 mg/m<sup>2</sup> over 24 hours had progressive hypotension and died. The arrhythmias included asymptomatic ventricular tachycardia, bigeminy and complete AV block requiring pacemaker placement. Among patients with NSCLC treated with TAXOL in combination with cisplatin in the Phase 3 study, significant cardiovascular events occurred in 12%-13%. This apparent increase in cardiovascular events is possibly due to an increase in cardiovascular risk factors in patients with lung cancer.

Electrocardiogram (ECG) abnormalities were common among patients at baseline. ECG abnormalities on study did not usually result in symptoms, were not dose-limiting, and required no intervention. ECG abnormalities were noted in 23% of all patients. Among patients with a normal ECG prior to study entry, 14% of all patients developed an abnormal tracing while on study. The most frequently reported ECG modifications were non-specific repolarization abnormalities, sinus bradycardia, sinus tachycardia, and premature beats. Among patients with normal ECGs at baseline, prior therapy with anthracyclines did not influence the frequency of ECG abnormalities.

Cases of myocardial infarction have been reported rarely. Congestive heart failure has been reported typically in patients who have received other chemotherapy, notably anthracyclines. (See **PRECAUTIONS: Drug Interactions**.)

Rare reports of atrial fibrillation and supraventricular tachycardia have been received as part of the continuing surveillance of TAXOL safety.

**Respiratory:** Rare reports of interstitial pneumonia, lung fibrosis, and pulmonary embolism have been received as part of the continuing surveillance of TAXOL safety. Rare reports of radiation pneumonitis have been received in patients receiving concurrent radiotherapy.

**Neurologic:** The assessment of neurologic toxicity was conducted differently among the studies as evident from the data reported in each individual study (see Tables 9-15). Moreover, the frequency and severity of neurologic manifestations were influenced by prior and/or concomitant therapy with neurotoxic agents.

In general, the frequency and severity of neurologic manifestations were dose-dependent in patients receiving single-agent TAXOL (paclitaxel) Injection. Peripheral neuropathy was observed in 60% of all patients (3% severe) and in 52% (2% severe) of the patients without pre-existing neuropathy. The frequency of peripheral neuropathy increased with cumulative dose. Neurologic symptoms were observed in 27% of the patients after the first course of treatment and in 34%-51% from course 2 to 10. Peripheral neuropathy was the cause of TAXOL discontinuation in 1% of all patients. Sensory symptoms have usually improved or resolved within several months of TAXOL discontinuation. Pre-existing neuropathies resulting from prior therapies are not a contraindication for TAXOL therapy.

In the Intergroup first-line ovarian carcinoma study (see Table 10), neurotoxicity included reports of neuromotor and neurosensory events. The regimen with TAXOL 175 mg/m<sup>2</sup> given by 3-hour infusion plus cisplatin 75 mg/m<sup>2</sup> resulted in a greater incidence and severity of neurotoxicity than the regimen containing cyclophosphamide and cisplatin, 87% (21% severe) versus 52% (2% severe), respectively. The duration of grade III or IV neurotoxicity cannot be determined with precision for the Intergroup study since the resolution dates of adverse events were not collected in the case report forms for this trial and complete follow-up documentation was available only in a minority of these patients. In the GOG first-line ovarian carcinoma study, neurotoxicity was reported as peripheral neuropathy. The regimen with TAXOL 135 mg/m<sup>2</sup> given by 24-hour infusion plus cisplatin 75 mg/m<sup>2</sup> resulted in an incidence of neurotoxicity that was similar to the regimen containing cyclophosphamide plus cisplatin, 25% (3% severe) versus 20% (0% severe), respectively. Cross-study comparison of neurotoxicity in the Intergroup and GOG trials suggests that when TAXOL is given in combination with cisplatin 75 mg/m<sup>2</sup>, the incidence of severe neurotoxicity is more common at a TAXOL dose of 175 mg/m<sup>2</sup> given by 3-hour infusion (21%) than at a dose of 135 mg/m<sup>2</sup> given by 24-hour infusion (3%).

In patients with NSCLC, administration of TAXOL followed by cisplatin resulted in a greater incidence of severe neurotoxicity compared to the incidence in patients with ovarian or breast cancer treated with single-agent TAXOL. Severe neurosensory symptoms were noted in 13% of NSCLC patients receiving TAXOL 135 mg/m<sup>2</sup> by 24-hour infusion followed by cisplatin 75 mg/m<sup>2</sup> and 8% of NSCLC patients receiving cisplatin/etoposide (see Table 14).

Other than peripheral neuropathy, serious neurologic events following TAXOL administration have been rare (<1%) and have included grand mal seizures, syncope, ataxia, and neuroencephalopathy.

Rare reports of autonomic neuropathy resulting in paralytic ileus have been received as part of the continuing surveillance of TAXOL safety. Optic nerve and/or visual disturbances (scintillating scotomata) have also been reported, particularly in patients who have received higher doses than those recommended. These effects generally have been reversible. However, rare reports in the literature of abnormal visual evoked potentials in patients have suggested persistent optic nerve damage. Postmarketing reports of ototoxicity (hearing loss and tinnitus) have also been received.

**Arthralgia/Myalgia:** There was no consistent relationship between dose or schedule of TAXOL and the frequency or severity of arthralgia/myalgia. Sixty percent of all patients treated experienced arthralgia/myalgia; 8% experienced severe symptoms. The symptoms were usually transient, occurred two or three days after TAXOL administration, and resolved within a few days. The frequency and severity of musculoskeletal symptoms remained unchanged throughout the treatment period.

**Hepatic:** No relationship was observed between liver function abnormalities and either dose or schedule of TAXOL administration. Among patients with normal baseline liver function 7%, 22%, and 19% had elevations in bilirubin, alkaline phosphatase, and AST (SGOT), respectively. Prolonged exposure to TAXOL was not associated with cumulative hepatic toxicity.

Rare reports of hepatic necrosis and hepatic encephalopathy leading to death have been received as part of the continuing surveillance of TAXOL safety.

**Renal:** Among the patients treated for Kaposi's sarcoma with TAXOL, five patients had renal toxicity of grade III or IV severity. One patient with suspected HIV nephropathy of grade IV severity had to discontinue therapy. The other four patients had renal insufficiency with reversible elevations of serum creatinine.

**Gastrointestinal (GI):** Nausea/vomiting, diarrhea, and mucositis were reported by 52%, 38%, and 31% of all patients, respectively. These manifestations were usually mild to moderate. Mucositis was schedule dependent and occurred more frequently with the 24-hour than with the 3-hour infusion.

In patients with poor-risk AIDS-related Kaposi's sarcoma, nausea/vomiting, diarrhea, and mucositis were reported by 69%, 79%, and 28% of patients, respectively. One third of patients with Kaposi's sarcoma complained of diarrhea prior to study start. (See **CLINICAL STUDIES: AIDS-Related Kaposi's Sarcoma**.)

In the first-line Phase 3 ovarian carcinoma studies, the incidence of nausea and vomiting when TAXOL was administered in combination with cisplatin appeared to be greater compared with the database for single-agent TAXOL in ovarian and breast carcinoma. In addition, diarrhea of any grade was reported more frequently compared to the control arm, but there was no difference for severe diarrhea in these studies.

Rare reports of intestinal obstruction, intestinal perforation, pancreatitis, ischemic colitis, and dehydration have been received as part of the continuing surveillance of TAXOL safety. Rare reports of neutropenic enterocolitis (typhlitis), despite the coadministration of G-CSF, were observed in patients treated with TAXOL alone and in combination with other chemotherapeutic agents.

**Injection Site Reaction:** Injection site reactions, including reactions secondary to extravasation, were usually mild and consisted of erythema, tenderness, skin discoloration, or swelling at the injection site. These reactions have been observed more frequently with the 24-hour infusion than with the 3-hour infusion. Recurrence of skin reactions at a site of previous extravasation following administration of TAXOL at a different site, i.e., "recall", has been reported rarely.

Rare reports of more severe events such as phlebitis, cellulitis, induration, skin exfoliation, necrosis, and fibrosis have been received as part of the continuing surveillance of TAXOL safety. In some cases the onset of the injection site reaction either occurred during a prolonged infusion or was delayed by a week to ten days.

A specific treatment for extravasation reactions is unknown at this time. Given the possibility of extravasation, it is advisable to closely monitor the infusion site for possible infiltration during drug administration.

**Other Clinical Events:** Alopecia was observed in almost all (87%) of the patients. Transient skin changes due to TAXOL (paclitaxel) Injection-related hypersensitivity reactions have been observed, but no other skin toxicities were significantly associated with TAXOL administration. Nail changes (changes in pigmentation or discoloration of nail bed) were uncommon (2%). Edema was reported in 21% of all patients (17% of those without baseline edema); only 1% had severe edema and none of these patients required treatment discontinuation. Edema was most commonly focal and disease-related. Edema was observed in 5% of all courses for patients with normal baseline and did not increase with time on study.

Rare reports of skin abnormalities related to radiation recall as well as reports of maculopapular rash, pruritus, Stevens-Johnson syndrome, and toxic epidermal necrolysis have been received as part of the continuing surveillance of TAXOL safety.

Reports of asthenia and malaise have been received as part of the continuing surveillance of TAXOL safety. In the Phase 3 trial of TAXOL 135 mg/m<sup>2</sup> over 24 hours in combination with cisplatin as first-line therapy of ovarian cancer, asthenia was reported in 17% of the patients, significantly greater than the 10% incidence observed in the control arm of cyclophosphamide/cisplatin.

**Accidental Exposure:** Upon inhalation, dyspnea, chest pain, burning eyes, sore throat, and nausea have been reported. Following topical exposure, events have included tingling, burning, and redness.

#### OVERDOSAGE

There is no known antidote for TAXOL overdose. The primary anticipated complications of overdose would consist of bone marrow suppression, peripheral neurotoxicity, and mucositis. Overdoses in pediatric patients may be associated with acute ethanol toxicity (see **PRECAUTIONS: Pediatric Use**).

#### DOSE AND ADMINISTRATION

**Note:** Contact of the undiluted concentrate with plasticized PVC equipment or devices used to prepare solutions for infusion is not recommended. In order to minimize patient exposure to the plasticizer DEHP [di-(2-ethylhexyl)phthalate], which may be leached from PVC infusion bags or sets, diluted TAXOL solutions should be stored in bottles (glass, polypropylene) or plastic bags (polypropylene, polyolefin) and administered through polyethylene-lined administration sets.

All patients should be premedicated prior to TAXOL administration in order to prevent severe hypersensitivity reactions. Such premedication may consist of dexamethasone 20 mg PO administered approximately 12 and 6 hours before TAXOL, diphenhydramine (or its equivalent) 50 mg I.V. 30 to 60 minutes prior to TAXOL, and cetirizine (300 mg) or ranitidine (50 mg) I.V. 30 to 60 minutes before TAXOL.

For patients with carcinoma of the ovary, the following regimens are recommended (see **CLINICAL STUDIES: Ovarian Carcinoma**):

- 1) For previously untreated patients with carcinoma of the ovary, one of the following recommended regimens may be given every 3 weeks. In selecting the appropriate regimen, differences in toxicities should be considered (see Table 10 in **ADVERSE REACTIONS: Disease-Specific Adverse Event Experiences**).

- a. TAXOL administered intravenously over 3 hours at a dose of 175 mg/m<sup>2</sup> followed by cisplatin at a dose of 75 mg/m<sup>2</sup>; or
- b. TAXOL administered intravenously over 24 hours at a dose of 135 mg/m<sup>2</sup> followed by cisplatin at a dose of 75 mg/m<sup>2</sup>.

- 2) In patients previously treated with chemotherapy for carcinoma of the ovary, TAXOL has been used at several doses and schedules; however, the optimal regimen is not yet clear. The recommended regimen is TAXOL 135 mg/m<sup>2</sup> or 175 mg/m<sup>2</sup> administered intravenously over 3 hours every 3 weeks.

For patients with carcinoma of the breast, the following regimens are recommended (see **CLINICAL STUDIES: Breast Carcinoma**):

- 1) For the adjuvant treatment of node-positive breast cancer, the recommended regimen is TAXOL, at a dose of 175 mg/m<sup>2</sup> intravenously over 3 hours every 3 weeks for four courses administered sequentially to doxorubicin-containing combination chemotherapy. The clinical trial used four courses of doxorubicin and cyclophosphamide (see **CLINICAL STUDIES: Breast Carcinoma**).

- 2) After failure of initial chemotherapy for metastatic disease or relapse within 6 months of adjuvant chemotherapy, TAXOL at a dose of 175 mg/m<sup>2</sup> administered intravenously over 3 hours every 3 weeks has been shown to be effective.

For patients with non-small cell lung carcinoma, the recommended regimen, given every 3 weeks, is TAXOL administered intravenously over 24 hours at a dose of 135 mg/m<sup>2</sup> followed by cisplatin, 75 mg/m<sup>2</sup>.

For patients with AIDS-related Kaposi's sarcoma, TAXOL administered at a dose of 135 mg/m<sup>2</sup> given intravenously over 3 hours every 3 weeks or at a dose of 100 mg/m<sup>2</sup> given intravenously over 3 hours every 2 weeks is recommended (dose intensity 45-50 mg/m<sup>2</sup>/week). In the two clinical trials evaluating these schedules (see **CLINICAL STUDIES: AIDS-Related Kaposi's Sarcoma**), the former schedule (135 mg/m<sup>2</sup> every 3 weeks) was more toxic than the latter. In addition, all patients with low performance status were treated with the latter schedule (100 mg/m<sup>2</sup> every 2 weeks).

Based upon the immunosuppression in patients with advanced HIV disease, the following modifications are recommended in these patients:

- 1) Reduce the dose of dexamethasone as one of the three premedication drugs to 10 mg PO (instead of 20 mg PO);
- 2) Initiate or repeat treatment with TAXOL only if the neutrophil count is at least 1000 cells/mm<sup>3</sup>;
- 3) Reduce the dose of subsequent courses of TAXOL by 20% for patients who experience severe neutropenia (neutrophil <500 cells/mm<sup>3</sup> for a week or longer); and
- 4) Initiate concomitant hematopoietic growth factor (G-CSF) as clinically indicated.

For the therapy of patients with solid tumors (ovary, breast, and NSCLC), courses of TAXOL should not be repeated until the neutrophil count is at least 1,500 cells/mm<sup>3</sup> and the platelet count is at least 100,000 cells/mm<sup>3</sup>. TAXOL should not be given to patients with AIDS-related Kaposi's sarcoma if the baseline or subsequent neutrophil count is less than 1000 cells/mm<sup>3</sup>. Patients who experience severe neutropenia

(neutrophil <500 cells/mm<sup>3</sup> for a week or longer) or severe peripheral neuropathy during TAXOL (paclitaxel) Injection therapy should have dosage reduced by 20% for subsequent courses of TAXOL. The incidence of neurotoxicity and the severity of neutropenia increase with dose.

**Preparation and Administration Precautions:** TAXOL is a cytotoxic anticancer drug and, as with other potentially toxic compounds, caution should be exercised in handling TAXOL. The use of gloves is recommended. If TAXOL solution contacts the skin, wash the skin immediately and thoroughly with soap and water. Following topical exposure, events have included tingling, burning, and redness. If TAXOL contacts mucous membranes, the membranes should be flushed thoroughly with water. Upon inhalation, dyspnea, chest pain, burning eyes, sore throat, and nausea have been reported.

Given the possibility of extravasation, it is advisable to closely monitor the infusion site for possible infiltration during drug administration. (See **PRECAUTIONS: Injection Site Reaction.**)

**Preparation for Intravenous Administration:** TAXOL must be diluted prior to infusion. TAXOL should be diluted in 0.9% Sodium Chloride Injection, USP; 5% Dextrose Injection, USP; 5% Dextrose and 0.9% Sodium Chloride Injection, USP, or 5% Dextrose in Ringer's Injection to a final concentration of 0.3 to 1.2 mg/mL. The solutions are physically and chemically stable for up to 27 hours at ambient temperature (approximately 25°C) and room lighting conditions. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

Upon preparation, solutions may show haziness, which is attributed to the formulation vehicle. No significant losses in potency have been noted following simulated delivery of the solution through I.V. tubing containing an in-line (0.22 micron) filter.

Data collected for the presence of the extractable plasticizer DEHP [di-(2-ethylhexyl)phthalate] show that levels increase with time and concentration when dilutions are prepared in PVC containers. Consequently, the use of plasticized PVC containers and administration sets is not recommended. TAXOL solutions should be prepared and stored in glass, polypropylene, or polyolefin containers. Non-PVC containing administration sets, such as those which are polyethylene-lined, should be used.

TAXOL should be administered through an in-line filter with a microporous membrane not greater than 0.22 microns. Use of filter devices such as IVEX-2<sup>®</sup> filters which incorporate short inlet and outlet PVC-coated tubing has not resulted in significant leaching of DEHP.

The Chemo Dispensing Pin™ device or similar devices with spikes should not be used with vials of TAXOL since they can cause the stopper to collapse resulting in loss of sterile integrity of the TAXOL solution.

**Stability:** Unopened vials of TAXOL Injection are stable until the date indicated on the package when stored between 20°-25°C (68°-77°F), in the original package. Neither freezing nor refrigeration adversely affects the stability of the product. Upon refrigeration components in the TAXOL vial may precipitate, but will redissolve upon reaching room temperature with little or no agitation. There is no impact on product quality under these circumstances. If the solution remains cloudy or if an insoluble precipitate is noted, the vial should be discarded. Solutions for infusion prepared as recommended are stable at ambient temperature (approximately 25°C) and lighting conditions for up to 27 hours.

#### HOW SUPPLIED

NDC 0015-3475-30	30 mg/5 mL multidose vial individually packaged in a carton.
NDC 0015-3476-30	100 mg/16.7 mL multidose vial individually packaged in a carton.
NDC 0015-3479-11	300 mg/50 mL multidose vial individually packaged in a carton.

**Storage:** Store the vials in original cartons between 20°-25°C (68°-77°F). Retain in the original package to protect from light.

**Handling and Disposal:** Procedures for proper handling and disposal of anticancer drugs should be considered. Several guidelines on this subject have been published.<sup>1-7</sup> There is no general agreement that all of the procedures recommended in the guidelines are necessary or appropriate.

#### REFERENCES

1. Recommendations for the Safe Handling of Parenteral Antineoplastic Drugs. NIH Publication No. 83-2621. For sale by the Superintendent of Documents, US Government Printing Office, Washington, DC 20402.
2. AMA Council Report. Guidelines for Handling Parenteral Antineoplastics. *JAMA* 1985; 253(11):1590-1592.
3. National Study Commission on Cytotoxic Exposure—Recommendations for Handling Cytotoxic Agents. Available from Louis P. Jeffrey, Chairman, National Study Commission on Cytotoxic Exposure, Massachusetts College of Pharmacy and Allied Health Sciences, 179 Longwood Avenue, Boston, Massachusetts, 02115.
4. Clinical Oncological Society of Australia. Guidelines and Recommendations for Safe Handling of Antineoplastic Agents. *Med J Australia* 1983; 1:426-428.
5. Jones RB, et al: Safe Handling of Chemotherapeutic Agents: A Report from the Mount Sinai Medical Center. *CA-A Cancer Journal for Clinicians* 1983; (Sept/Oct) 258-263.
6. American Society of Hospital Pharmacists Technical Assistance Bulletin on Handling Cytotoxic and Hazardous Drugs. *Am J Hosp Pharm* 1990; 47:1033-1049.
7. Controlling Occupational Exposure to Hazardous Drugs. (OSHA WORK-PRACTICE GUIDELINES.) *Am J Health-Syst Pharm* 1996; 53:1669-1685.

## PATIENT INFORMATION

# TAXOL<sup>®</sup> Injection

(generic name = paclitaxel)

#### WHAT IS TAXOL?

TAXOL is a prescription cancer medicine. It is injected into a vein and it is used to treat different types of tumors. The tumors include advanced ovary and breast cancer. The tumors also include certain lung cancers (non-small cell) in people who cannot have surgery or radiation therapy. TAXOL may also be used to treat AIDS-related Kaposi's sarcoma.

#### WHAT IS CANCER?

Under normal conditions, the cells in your body divide and grow in an orderly, controlled way. Cell division and growth are necessary for the human body to perform its functions and to repair itself, when necessary. Cancer cells are different from normal cells because they are not able to control their own growth. The reasons for this abnormal growth are not yet fully understood.

A tumor is a mass of unhealthy cells that are dividing and growing fast and in an uncontrolled way. When a tumor invades surrounding healthy body tissue it is known as a malignant tumor. A malignant tumor can spread (metastasize) from its original site to other parts of the body if not found and treated early.

#### HOW DOES TAXOL WORK?

TAXOL is a type of medical treatment called chemotherapy. The purpose of chemotherapy is to kill cancer cells or prevent their growth.

All cells, whether they are healthy cells or cancer cells, go through several stages of growth. During one of the stages, the cell starts to divide. TAXOL may stop the cells from dividing and growing, so they eventually die. In addition, normal cells may also be affected by TAXOL causing some of the side effects. (See **WHAT ARE THE POSSIBLE SIDE EFFECTS OF TAXOL?** below.)

#### WHO SHOULD NOT TAKE TAXOL?

Patients who have a history of hypersensitivity (allergic reactions) to TAXOL or other drugs containing Cremophor<sup>®</sup> EL\* (polyoxyethylated castor oil), like cyclosporine or teniposide, should not be given TAXOL. In addition, TAXOL should not be given to patients with dangerously low white blood cell counts.

#### HOW IS TAXOL GIVEN?

TAXOL is injected into a vein [intravenous (I.V.) infusion]. Before you are given TAXOL, you will have to take certain medicines (premedications) to prevent or reduce the chance you will have a serious allergic reaction. Such reactions have occurred in a small number of patients while receiving TAXOL and have been rarely fatal. (See **WHAT ARE THE POSSIBLE SIDE EFFECTS OF TAXOL?** below.)

#### WHAT ARE THE POSSIBLE SIDE EFFECTS OF TAXOL?

Most patients taking TAXOL will experience side effects, although it is not always possible to tell whether such effects are caused by TAXOL, another medicine they may be taking, or the cancer itself. Important side effects are described below; however, some patients may experience other side effects that are less common. *Report any unusual symptoms to your doctor.*

Important side effects observed in studies of patients taking TAXOL were as follows:

- **Allergic reactions.** Allergic reactions can vary in degrees of severity. They may cause death in rare cases. When a severe allergic reaction develops, it usually occurs at the time the medicine is entering the body (during TAXOL infusion). Allergic reactions may cause trouble breathing, very low blood pressure, sudden swelling, and/or hives or rash. The likelihood of a serious allergic reaction is lowered by the use of several kinds of medicines that are given to you before the TAXOL infusion.
- **Heart and blood vessel (cardiovascular) effects.** TAXOL may cause a drop in heart rate (bradycardia) and low blood pressure (hypotension). The patient usually does not notice these changes. These changes usually do not require treatment. Your heart function, including blood pressure and pulse, will be monitored while you are receiving the medicine. You should notify your doctor if you have a history of heart disease.
- **Infections due to low white blood cell count.** Among the body's defenses against bacterial infections are white blood cells. Between your TAXOL treatment cycles, you will often have blood tests to check your white blood cell counts. TAXOL usually causes a brief drop in white blood cells. *If you have a fever (temperature above 100.4°F) or other sign of infection, tell your doctor right away. Sometimes serious infections develop that require treatment in the hospital with antibiotics. Serious illness or death could result if such infections are not treated when white blood cell counts are low.*
- **Hair loss.** Complete hair loss, or alopecia, almost always occurs with TAXOL. This usually involves the loss of eyebrows, eyelashes, and pubic hair, as well as scalp hair. It can occur suddenly after treatment has begun, but usually happens 14 to 21 days after treatment. *Hair generally grows back after you've finished your TAXOL treatment.*

- **Joint and muscle pain.** You may get joint and muscle pain a few days after your TAXOL (paclitaxel) Injection treatment. These symptoms usually disappear in a few days. Although pain medicine may not be necessary, tell your doctor if you are uncomfortable.
- **Irritation at the injection site.** TAXOL sometimes causes irritation at the site where it enters the vein. Reactions may include discomfort, redness, swelling, inflammation (of the surrounding skin or of the vein itself), and ulceration (open sores). These reactions are usually caused by the I.V. (intravenous) fluid leaking into the surrounding area. *If you notice anything unusual at the site of the injection (needle), either during or after treatment, tell your doctor right away.*
- **Low red blood cell count.** Red blood cells deliver oxygen to tissues throughout all parts of the body and take carbon dioxide from the tissues by using a protein called hemoglobin. A lowering of the volume of red blood cells may occur following TAXOL treatment causing anemia. Some patients may need a blood transfusion to treat the anemia. Patients can feel tired, tire easily, appear pale, and become short of breath. Contact your doctor if you experience any of these symptoms following TAXOL treatment.
- **Mouth or lip sores (mucositis).** Some patients develop redness and/or sores in the mouth or on the lips. These symptoms might occur a few days after the TAXOL treatment and usually decrease or disappear within one week. Talk with your doctor about proper mouth care and other ways to prevent or reduce your chances of developing mucositis.
- **Numbness, tingling, or burning in the hands and/or feet (neuropathy).** These symptoms occur often with TAXOL and usually get better or go away without medication within several months of completing treatment. However, if you are uncomfortable, tell your doctor so that he/she can decide the best approach for relief of your symptoms.
- **Stomach upset and diarrhea.** Some patients experience nausea, vomiting, and/or diarrhea following TAXOL use. If you experience nausea or stomach upset, tell your doctor. Diarrhea will usually disappear without treatment; however, *if you experience severe abdominal or stomach area pain and/or severe diarrhea, tell your doctor right away.*

Talk with your doctor or other healthcare professional to discuss ways to prevent or reduce some of these side effects. Because this leaflet does not include all possible side effects that can occur with TAXOL, it is important to talk with your doctor about other possible side effects.

#### CAN I TAKE TAXOL IF I AM PREGNANT OR NURSING A BABY?

TAXOL could harm the fetus when given to a pregnant woman. Women should avoid becoming pregnant while they are undergoing treatment with TAXOL. *Tell your doctor if you become pregnant or plan to become pregnant while taking TAXOL.*

Because studies have shown TAXOL to be present in the breast milk of animals receiving the drug, it may be present in human breast milk as well. Therefore, nursing a baby while taking TAXOL is NOT recommended.

This medicine was prescribed for your particular condition. This summary does not include everything there is to know about TAXOL. Medicines are sometimes prescribed for purposes other than those listed in a Patient Information Leaflet. If you have questions or concerns, or want more information about TAXOL, your doctor and pharmacist have the complete prescribing information upon which this guide is based. You may want to read it and discuss it with your doctor. Remember, no written summary can replace careful discussion with your doctor.

\* Cremophor<sup>®</sup> EL is the registered trademark of BASF Aktiengesellschaft. Cremophor<sup>®</sup> EL is further purified by a Bristol-Myers Squibb Company proprietary process before use.

This Patient Information Leaflet has been approved by the U.S. Food and Drug Administration.



K4-B001-9-00  
347630D1M-13

Based on: 1109663A7  
Revised July 2000

**CENTER FOR DRUG EVALUATION AND RESEARCH**

**APPROVAL PACKAGE FOR:**

**APPLICATION NUMBER**

**NDA 20-262/S-036/S-037/S-038**

**Medical Review(s)**

## Medical Officer Labeling Review

**Application:** sNDA 20-262  
**Sponsor:** Bristol-Myers Squibb Pharmaceutical Research Institute  
**Drug:** Paclitaxel

**Submissions:** 5/11/00 SLR-037  
7/6/00 SE2-036  
8/8/00 SLR-038  
2/15/02 SNC to SE2-036

**Date to Reviewer:** 2/6/02  
**Review date:** 2/6/02  
2/28/02

The sponsor submitted 3 labeling supplements. The contents of the supplements are reviewed below.

### SLR-037

This supplement contained formatting and punctuation changes. In addition, the risk of ototoxicity was added to the Adverse Events section of the label.

#### Reviewer Comment:

All changes are acceptable as proposed and are supported by a review of post-marketing events reported to AERS, confirmed by the reviewer.

### SLR-038

This supplement added the risks of Stevens-Johnson syndrome and toxic epidermal necrolysis to the label, which are supported by a review of post-marketing events reported to AERS, confirmed by the reviewer.

#### Reviewer Comment:

All changes are acceptable as proposed.

### SE2-036

This supplement corrected the number of deaths observed for ER(+) women in the adjuvant breast cancer study in Table 4, referred patients to a Patient Information Leaflet, and updated adverse events.

#### Reviewer Comments:

1. The following comment refers to the updated number of death for the breast cancer study:

The hazard ratio of 0.83 (95% CI 0.59-1.18) reported in Table 4 for the receptor positive subset of the adjuvant breast cancer trial was calculated based on 67 deaths. You have updated the number of deaths to 126 but have not provided an updated hazard ratio.

This change is unacceptable unless the HR and its 95% CI are changed accordingly, or unless you can provide documentation that the HR as written was calculated using the updated death rate.

2. Table 10 now contains a footnote that indicates that the difference in any severity of diarrheal symptoms in the Intergroup study (first-line ovarian cancer) is statistically significantly worse on the taxol arm. This footnote was previously approved for the difference in diarrheal symptoms in GOG-111. This change is acceptable.

3. The proposed re-wording of the neurotoxicity warning in the Adverse Events section is acceptable.

### **Summary**

1. The following comment should be sent to the sponsor for SE2-036:

The hazard ratio of 0.83 (95% CI 0.59-1.18) reported in Table 4 for the receptor positive subset of the adjuvant breast cancer trial was calculated based on 67 deaths. You have updated the number of deaths to 126 but have not provided an updated hazard ratio. This change is unacceptable unless the HR and its 95% CI are changed accordingly, or unless you can provide documentation that the HR as written was calculated using the updated death rate.

2. All other proposed changes for the 3 supplements are acceptable.

### ***Addendum:***

Comment 1 was sent to the sponsor. The sponsor replied 2/15/02 SNC to SE2-036:

"There were 67 deaths on the doxorubicin/cyclophosphamide (AC) arm and 59 deaths on the AC plus Taxol (AC+T) arm, as can be seen from Figure S.7.7F on page 287 of the final study report for protocol CA139-223 (Document Control Number 910071561). This study report was submitted to the Agency on April 26, 1999 (S-033). The hazard ratio and CI estimate are based on those 126 events. The change in the number of deaths from 67 to 126 in the FPL was simply to correct the error."

This reply is acceptable. The changes are acceptable, and the 3 supplements are approved.

**/S/**

---

Susan Flamm Honig, M.D.  
Medical Reviewer

**/S/**

---

Grant Williams, M.D  
Team Leader

-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/

-----  
Susan Honig  
2/28/02 03:42:07 PM  
MEDICAL OFFICER

Grant, you have already signed this review in hard copy.

Grant Williams  
3/1/02 06:48:07 AM  
MEDICAL OFFICER

**CENTER FOR DRUG EVALUATION AND RESEARCH**

**APPROVAL PACKAGE FOR:**

**APPLICATION NUMBER**

**NDA 20-262/S-036/S-037/S-038**

**Administrative/Correspondence Reviews**

**DIVISION OF ONCOLOGY DRUG PRODUCTS  
CSO LABELING REVIEW**

**NDA:** NDA 20-262/S-036 FA  
NDA 20-262/S-037  
NDA 20-262/S-038

**DRUG:** TAXOL (paclitaxel) Injection

**SPONSOR:** Bristol –Myers Squibb Company

**DATES OF SUBMISSIONS:**

<u>Supplement Number</u>	<u>Submission Date</u>	<u>Receipt Date</u>
S-036 FA	July 6, 2000	July 7, 2000
S-037	May 11, 2000	May 12, 2000
S-038	August 8, 2000	August 9, 2000

**BACKGROUND:**

Supplement 036 FA provides final printed labeling in follow-up to the Approval letter dated June 20, 2000 for SE1-036. This FPL also contains the Patient Package Insert (PPI) and incorporates the CBE changes requested in SLR-037.

Special Supplement Changes Being Effected 037 provides for changes to the **ADVERSE REACTIONS** section by incorporating ototoxicity into the list of adverse events reported during the continuing surveillance of Taxol safety. Supplement 037 was submitted with final printed labeling (including a PPI) and is superseded by supplements 036 FA and 038.

Special Supplement Changes Being Effected 038 provides for changes to the **ADVERSE REACTIONS** section by incorporating Steven's Johnson Syndrome and toxic epidermal necrolysis to the list of adverse events reported during the continuing surveillance of Taxol safety. Supplement 038 was submitted with final printed labeling (including a PPI).

All of the proposed package inserts were compared to the most recently approved labeling for SE1-036 dated June 20, 2000. The PPIs submitted were compared to the most recently approved PPI dated December 10, 1999, submitted as supplement 032. All changes and/or discrepancies are outlined below.

**DISCUSSION:**

1. In the **CLINICAL STUDIES** section, **Breast Carcinoma** subsection, *Subset analyses* subsection, **Table 4**, the number of deaths reported in the Overall Survival column, **Receptor Status Positive** row, has been changed from 67 to 126.

*Comment: This change was reviewed by the Medical Officer on 2-6-02. The following comment was sent to the sponsor for clarification by fax on 2-12-02:*

*“The hazard ratio of 0.83 (95% CI 0.59 – 1.18) reported in Table 4 for the receptor positive subset of the adjuvant breast cancer trial was calculated based on 67 deaths. You have updated the number of deaths to 126 but have not provided an updated hazard ratio. This change is unacceptable unless the HR and its 95% CI are changed accordingly, or unless you can provide documentation that the HR as written was calculated using the updated death rate.”*

*In a fax to the Division dated February 15, 2002, the sponsor provided the following response:*

*“There were 67 deaths on the doxorubicin/cyclophosphamide (AC) arm and 59 deaths on the AC plus Taxol (AC+T) arm, as can be seen from Figure S.7.7F on page 287 of the final study report for protocol CA139-223 (Document Control Number 910071561). This study report was submitted to the Agency on April 26, 1999 (S-033). The hazard ratio and CI estimate are based on those 126 events. The change in the number of deaths from 67 to 126 in the FPL was simply to correct the error.”*

*The Medical Officer reviewed the sponsor’s response on 2-26-02 and found it adequate. The change is acceptable.*

2. The following statement has been added at the end of the **PRECAUTIONS** section:

**“Information for Patients: (See Patient Information Leaflet).”**

*Comment: This change was requested by the FDA in the approval letter for Supplement 032 dated December 10, 1999. It is acceptable.*

3. In the **ADVERSE REACTIONS** section, **Disease-Specific Adverse Event Experiences** subsection, **First-Line Ovary in Combination** subsection, **Table 10, Diarrhea (Any symptoms)** row, **Intergroup** columns, the notation <sup>“d”</sup> has been added to the numbers 37 and 29. The notation means  $p < 0.05$  by Fisher exact test.

*Comment: This change was reviewed by the Medical Officer on 2-6-02 and found to be acceptable.*

4. In the **ADVERSE REACTIONS** section, **Adverse Event Experiences by Body System** subsection, **Neurologic** subsection, 3<sup>rd</sup> paragraph, 2<sup>nd</sup> sentence, has been changed as follows (*in italics*):

from

“The regimen with TAXOL 175 mg/m<sup>2</sup> given by 3-hour infusion plus cisplatin 75 mg/m<sup>2</sup> resulted in *a greater incidence of neurotoxicity and severe neurotoxicity* than the regimen containing cyclophosphamide and cisplatin, 87% (21% severe) versus 52% (2% severe) respectively.”

to

“The regimen with TAXOL 175 mg/m<sup>2</sup> given by 3-hour infusion plus cisplatin 75 mg/m<sup>2</sup> resulted in *greater incidence and severity* of neurotoxicity than the regimen containing cyclophosphamide and cisplatin, 87% (21% severe) versus 52% (2% severe) respectively.”

*Comment: This change was reviewed by the Medical Officer on 2-6-02 and found to be acceptable.*

5. In the **ADVERSE REACTIONS** section, **Adverse Event Experiences by Body System** subsection, **Neurologic** subsection, the following has been added as the last sentence of the last paragraph:

“Postmarketing reports of ototoxicity (hearing loss and tinnitus) have also been received.”

*Comment: This change was submitted in Supplement 037 based on reports during the continuing surveillance of Taxol safety. This change was reviewed by the Medical Officer on 2-6-02 and found to be acceptable.*

6. In the **ADVERSE REACTIONS** section, **Adverse Event Experiences by Body System** subsection, **Other Clinical Events** subsection, 2<sup>nd</sup> paragraph, “Stevens-Johnson syndrome, and toxic epidermal necrolysis” have both been added.

*Comment: This change was submitted in Supplement 038 based on reports during the continuing surveillance of Taxol safety. Stevens-Johnson syndrome was previously reported by the NCI in a Dear Doctor letter during the year 2000. This change was reviewed by the Medical Officer on 2-6-02 and found to be acceptable.*

7. Minor editorial changes have been made throughout the package insert.

*Comment: The editorial changes made are acceptable.*



-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/

-----  
Christy Wilson  
2/28/02 02:31:03 PM  
CSO

Final review. Draft review signed by DPease on 1-29-02;  
by SHonig on 2-6-02 (and again on 2-26-02);  
and by GrWilliams on 2-8-02.

Draft review signed by DPease on 1-29-02; SHonig on  
2-6-02 and 2-26-02; and GrWilliams on 2-8-02

Dotti Pease  
2/28/02 02:43:47 PM  
CSO

Susan Honig  
2/28/02 03:31:01 PM  
MEDICAL OFFICER

Grant Williams  
3/1/02 06:44:36 AM  
MEDICAL OFFICER