

**CENTER FOR DRUG EVALUATION AND RESEARCH**

**Approval Package for:**

***APPLICATION NUMBER:***

**50-587 / S-052**

**Trade Name: Primaxin I. V.**

**Generic Name: Imipenem and cilastatin for injection**

**Sponsor: Merck & Company, Inc.**

**Approval Date: August 28, 2000**

# CENTER FOR DRUG EVALUATION AND RESEARCH

*APPLICATION NUMBER:*

**50-587 / S-052**

## CONTENTS

### **Reviews / Information Included in this NDA Review.**

<b>Approval Letter</b>	<b>X</b>
<b>Approvable Letter</b>	
<b>Labeling</b>	<b>X</b>
<b>Medical Review(s)</b>	<b>X</b>
<b>Chemistry Review(s)</b>	
<b>Pharmacology Review(s)</b>	
<b>Statistical Review(s)</b>	
<b>Microbiology Review(s)</b>	
<b>Clinical Pharmacology/ Biopharmaceutics Review(s)</b>	
<b>Administrative/Correspondence Document(s)</b>	

**CENTER FOR DRUG EVALUATION AND RESEARCH**

*APPLICATION NUMBER:*

**50-587 / S-052**

**APPROVAL LETTER**

NDA 50-587/S-052

AUG 28 2000

Merck Research Laboratories  
Attention: Dennis M. Erb, Ph.D.  
Senior Director, Regulatory Affairs  
Sumneytown Pike  
P.O. Box 4, BLA-20  
West Point, PA 19486

Dear Dr. Erb:

Please refer to your supplemental new drug application dated August 18, 1999, received August 23, 1999, submitted under section 505(b) of the Federal Food, Drug, and Cosmetic Act for Primaxin I.V. (imipenem and cilastatin for injection). We note that this application is subject to the exemption provisions contained in section 1 25(d)(2) of Title I of the FDA Modernization Act of 1997.

This supplemental new drug application provides for revisions in the DESCRIPTION, PREPARATION OF SOLUTION, COMPATIBILITY AND STABILITY, and HOW SUPPLIED sections of the label to incorporate the approved MONO VIAL™ Vial text, as well as, editorial changes to the ADVERSE REACTIONS and PREPARATION OF SOLUTION sections of the label.

We have completed the review of this supplemental application and have concluded that adequate information has been presented to demonstrate that the drug product is safe and effective for use as recommended in the submitted final printed labeling (package insert submitted August 18, 1999). Accordingly, the supplemental application is approved effective on the date of this letter.

If a letter communicating important information about this drug product (i.e., a 'Dear Health Care Practitioner' letter) is issued to physicians and others responsible for patient care, we request that you submit a copy of the letter to this NDA and a copy to the following address:

MED WATCH, HF-2  
FDA  
5600 Fishers Lane  
Rockville, MD 20857

We remind you that you must comply with the requirements for an approved NDA set forth under 21 CFR 314.80 and 314.81.

**CENTER FOR DRUG EVALUATION AND RESEARCH**

*APPLICATION NUMBER:*

**50-587 / S-052**

**LABELING**


**PRIMAXIN® I.V.**  
**(IMIPENEM AND CILASTATIN FOR INJECTION)**

**For Intravenous Injection Only**


**DESCRIPTION**

PRIMAXIN<sup>†</sup> I.V. (Imipenem and Cilastatin for Injection) is a sterile formulation of imipenem (a thienamycin antibiotic) and cilastatin sodium (the inhibitor of the renal dipeptidase, dehydropeptidase I), with sodium bicarbonate added as a buffer. PRIMAXIN I.V. is a potent broad spectrum antibacterial agent for intravenous administration.

Imipenem (N-formimidoylthienamycin monohydrate) is a crystalline derivative of thienamycin, which is produced by *Streptomyces cattleya*. Its chemical name is (5*R*,6*S*)-3-[[2-(formimidoylamino)ethyl]thio]-6-[[*R*]-1-hydroxyethyl]-7-oxo-1-azabicyclo[3.2.0]hept-2-ene-2-carboxylic acid monohydrate. It is an off-white, nonhygroscopic crystalline compound with a molecular weight of 317.37. It is sparingly soluble in water and slightly soluble in methanol. Its empirical formula is C<sub>12</sub>H<sub>17</sub>N<sub>3</sub>O<sub>4</sub>S•H<sub>2</sub>O, and its structural formula is:



Cilastatin sodium is the sodium salt of a derivatized heptenoic acid. Its chemical name is sodium (Z)-7-[[*R*]-2-amino-2-carboxyethyl]thio]-2-[[*S*]-2,2-dimethylcyclopropanecarboxamido]-2-heptenoate. It is an off-white to yellowish-white, hygroscopic, amorphous compound with a molecular weight of 380.43. It is very soluble in water and in methanol. Its empirical formula is C<sub>16</sub>H<sub>25</sub>N<sub>2</sub>O<sub>5</sub>S Na, and its structural formula is:



PRIMAXIN I.V. is buffered to provide solutions in the pH range of 6.5 to 8.5. There is no significant change in pH when solutions are prepared and used as directed. (See **COMPATIBILITY AND STABILITY**.) PRIMAXIN I.V. 250 contains 18.8 mg of sodium (0.8 mEq) and PRIMAXIN I.V. 500 contains 37.5 mg of sodium (1.6 mEq). Solutions of PRIMAXIN I.V. range from colorless to yellow. Variations of color within this range do not affect the potency of the product.

**CLINICAL PHARMACOLOGY**

*Adults*

*Intravenous Administration*

Intravenous infusion of PRIMAXIN I.V. over 20 minutes results in peak plasma levels of imipenem antimicrobial activity that range from 14 to 24 µg/mL for the 250 mg dose, from 21 to 58 µg/mL for the 500 mg dose, and from 41 to 83 µg/mL for the 1000 mg dose. At these doses, plasma levels of imipenem antimicrobial activity decline to below 1 µg/mL or less in 4 to 6 hours. Peak plasma levels of cilastatin

<sup>†</sup> Registered trademark of MERCK & CO., Inc.  
COPYRIGHT © MERCK & CO., Inc., 1987, 1994, 1998  
All rights reserved

following a 20-minute intravenous infusion of PRIMAXIN I.V., range from 15 to 25 µg/mL for the 250 mg dose, from 31 to 49 µg/mL for the 500 mg dose, and from 56 to 88 µg/mL for the 1000 mg dose.

The plasma half-life of each component is approximately 1 hour. The binding of imipenem to human serum proteins is approximately 20% and that of cilastatin is approximately 40%. Approximately, 70% of the administered imipenem is recovered in the urine within 10 hours after which no further urinary excretion is detectable. Urine concentrations of imipenem in excess of 10 µg/mL can be maintained for up to 8 hours with PRIMAXIN I.V. at the 500-mg dose. Approximately, 70% of the cilastatin sodium dose is recovered in the urine within 10 hours of administration of PRIMAXIN I.V.

No accumulation of imipenem/cilastatin in plasma or urine is observed with regimens administered as frequently as every 6 hours in patients with normal renal function.

Imipenem, when administered alone, is metabolized in the kidneys by dehydropeptidase I resulting in relatively low levels in urine. Cilastatin sodium, an inhibitor of this enzyme, effectively prevents renal metabolism of imipenem so that when imipenem and cilastatin sodium are given concomitantly, fully adequate antibacterial levels of imipenem are achieved in the urine.

After a 1 gram dose of PRIMAXIN I.V., the following average levels of imipenem were measured (usually at 1 hour post dose except where indicated) in the tissues and fluids listed:

Tissue or Fluid	n	Imipenem Level µg/mL or µg/g	Range
Vitreous Humor	3	3.4 (3.5 hours post dose)	2.88-3.6
Aqueous Humor	5	2.99 (2 hours post dose)	2.4-3.9
Lung Tissue	8	5.6 (median)	3.5-15.5
Sputum	1	2.1	—
Pleural	1	22.0	—
Peritoneal	12	23.9 S.D.±5.3(2 hours post dose)	—
Bile	2	5.3 (2.25 hours post dose)	4.6-6.0
CSF (uninflamed)	5	1.0 (4 hours post dose)	0.26-2.0
CSF (inflamed)	7	2.6 (2 hours post dose)	0.5-5.5
Fallopian Tubes	1	13.6	—
Endometrium	1	11.1	—
Myometrium	1	5.0	—
Bone	10	2.6	0.4-5.4
Interstitial Fluid	12	16.4	10.0-22.6
Skin	12	4.4	NA
Fascia	12	4.4	NA

Imipenem-cilastatin sodium is hemodialyzable. However, usefulness of this procedure in the overdosage setting is questionable. (See **OVERDOSAGE**.)

#### Microbiology

The bactericidal activity of imipenem results from the inhibition of cell wall synthesis. Its greatest affinity is for penicillin binding proteins (PBPs) 1A, 1B, 2, 4, 5 and 6 of *Escherichia coli*, and 1A, 1B, 2, 4 and 5 of *Pseudomonas aeruginosa*. The lethal effect is related to binding to PBP 2 and PBP 1B.

Imipenem has a high degree of stability in the presence of beta-lactamases, both penicillinases and cephalosporinases produced by gram-negative and gram-positive bacteria. It is a potent inhibitor of beta-lactamases from certain gram-negative bacteria which are inherently resistant to most beta-lactam antibiotics, e.g., *Pseudomonas aeruginosa*, *Serratia* spp., and *Enterobacter* spp.

Imipenem has *in vitro* activity against a wide range of gram-positive and gram-negative organisms. Imipenem has been shown to be active against most strains of the following microorganisms, both *in vitro* and in clinical infections treated with the intravenous formulation of imipenem-cilastatin sodium as described in the **INDICATIONS AND USAGE** section.

#### Gram-positive aerobes:

*Enterococcus faecalis* (formerly *S. faecalis*)

(NOTE: Imipenem is inactive *in vitro* against *Enterococcus faecium* [formerly *S. faecium*].)

*Staphylococcus aureus* including penicillinase-producing strains

*Staphylococcus epidermidis* including penicillinase-producing strains

(NOTE: Methicillin-resistant staphylococci should be reported as resistant to imipenem.)

*Streptococcus agalactiae* (Group B streptococci)

*Streptococcus pneumoniae*

*Streptococcus pyogenes*

#### Gram-negative aerobes:

*Acinetobacter* spp.

*Citrobacter* spp.

*Enterobacter* spp.

*Escherichia coli*

*Gardnerella vaginalis*  
*Haemophilus influenzae*  
*Haemophilus parainfluenzae*  
*Klebsiella* spp.  
*Morganella morganii*  
*Proteus vulgaris*  
*Providencia rettgeri*  
*Pseudomonas aeruginosa*

(NOTE: Imipenem is inactive *in vitro* against *Xanthomonas (Pseudomonas) maltophilia* and some strains of *P. cepacia*.)

*Serratia* spp., including *S. marcescens*

Gram-positive anaerobes:

*Bifidobacterium* spp.  
*Clostridium* spp.  
*Eubacterium* spp.  
*Peptococcus* spp.  
*Peptostreptococcus* spp.  
*Propionibacterium* spp.

Gram-negative anaerobes:

*Bacteroides* spp., including *B. fragilis*  
*Fusobacterium* spp.

The following *in vitro* data are available, **but their clinical significance is unknown.**

Imipenem exhibits *in vitro* minimum inhibitory concentrations (MICs) of 4 µg/mL or less against most (≥90%) strains of the following microorganisms; however, the safety and effectiveness of imipenem in treating clinical infections due to these microorganisms have not been established in adequate and well-controlled clinical trials.

Gram-positive aerobes:

*Bacillus* spp.  
*Listeria monocytogenes*  
*Nocardia* spp.  
*Staphylococcus saprophyticus*  
 Group C streptococci  
 Group G streptococci  
 Viridans group streptococci

Gram-negative aerobes:

*Aeromonas hydrophila*  
*Alcaligenes* spp.  
*Capnocytophaga* spp.  
*Haemophilus ducreyi*  
*Neisseria gonorrhoeae* including penicillinase-producing strains  
*Pasteurella* spp.  
*Providencia stuartii*

Gram-negative anaerobes:

*Prevotella bivia*  
*Prevotella disiens*  
*Prevotella melaninogenica*  
*Veillonella* spp.

*In vitro* tests show imipenem to act synergistically with aminoglycoside antibiotics against some isolates of *Pseudomonas aeruginosa*.

**Susceptibility Tests:**

Measurement of MIC or minimum bactericidal concentration (MBC) and achieved antimicrobial compound concentrations may be appropriate to guide therapy in some infections. (See **CLINICAL PHARMACOLOGY** section for further information on drug concentrations achieved in infected body sites and other pharmacokinetic properties of this antimicrobial drug product.)

**Dilution Techniques:**

Quantitative methods that are used to determine MICs provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. One such procedure uses a standardized dilution method<sup>1</sup> (broth, agar, or microdilution) or equivalent with imipenem powder.

The MIC values obtained should be interpreted according to the following criteria:

MIC (µg/mL)	Interpretation
≤4	Susceptible (S)
8	Intermediate (I)
≥16	Resistant (R)

A report of "Susceptible" indicates that the pathogen is likely to be inhibited by usually achievable concentrations of the antimicrobial compound in blood. A report of "Intermediate" indicates that the result should be considered equivocal, and, if the microorganism is not fully susceptible to alternative, clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where high dosage of drug can be used. This category also provides a buffer zone that prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of "Resistant" indicates that usually achievable concentrations of the antimicrobial compound in the blood are unlikely to be inhibitory and that other therapy should be selected.

Standardized susceptibility test procedures require the use of laboratory control microorganisms. Standard imipenem powder should provide the following MIC values:

Microorganism	MIC (µg/mL)
<i>E. coli</i> ATCC 25922	0.06–0.25
<i>S. aureus</i> ATCC 29213	0.015–0.06
<i>E. faecalis</i> ATCC 29212	0.5–2.0
<i>P. aeruginosa</i> ATCC 27853	1.0–4.0

#### Diffusion Techniques:

Quantitative methods that require measurement of zone diameters provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. One such standardized procedure<sup>2</sup> that has been recommended for use with disks to test the susceptibility of microorganisms to imipenem uses the 10-µg imipenem disk. Interpretation involves correlation of the diameter obtained in the disk test with the MIC for imipenem.

Reports from the laboratory providing results of the standard single-disk susceptibility test with a 10-µg imipenem disk should be interpreted according to the following criteria:

Zone Diameter (mm)	Interpretation
≥16	Susceptible (S)
14–15	Intermediate (I)
≤13	Resistant (R)

Interpretation should be as stated above for results using dilution techniques.

Standardized susceptibility test procedures require the use of laboratory control microorganisms. The 10-µg imipenem disk should provide the following diameters in these laboratory test quality control strains:

Microorganism	Zone Diameter (mm)
<i>E. coli</i> ATCC 25922	26–32
<i>P. aeruginosa</i> ATCC 27853	20–28

#### Anaerobic techniques:

For anaerobic bacteria, the susceptibility to imipenem can be determined by the reference agar dilution method or by alternate standardized test methods.<sup>3</sup>

The MIC values obtained should be interpreted according to the following criteria:

MIC (µg/mL)	Interpretation
≤4	Susceptible (S)
8	Intermediate (I)
≥16	Resistant (R)

As with other susceptibility techniques, the use of laboratory control microorganisms is required. Standard imipenem powder should provide the following MIC values:

#### Reference Agar Dilution Testing:

Microorganism	MIC (µg/mL)
<i>B. fragilis</i> ATCC 25285	0.03–0.12
<i>B. thetaiotaomicron</i> ATCC 29741	0.06–0.25
<i>E. lentum</i> ATCC 43055	0.25–1.0

#### Broth Microdilution Testing:

Microorganism	MIC (µg/mL)
<i>B. thetaiotaomicron</i> ATCC 29741	0.06–0.25
<i>E. lentum</i> ATCC 43055	0.12–0.5

## INDICATIONS AND USAGE

PRIMAXIN I.V. is indicated for the treatment of serious infections caused by susceptible strains of the designated microorganisms in the conditions listed below:

(1) **Lower respiratory tract infections.** *Staphylococcus aureus* (penicillinase-producing strains), *Acinetobacter* species, *Enterobacter* species, *Escherichia coli*, *Haemophilus influenzae*, *Haemophilus parainfluenzae*\*, *Klebsiella* species, *Serratia marcescens*

(2) **Urinary tract infections** (complicated and uncomplicated). *Enterococcus faecalis*, *Staphylococcus aureus* (penicillinase-producing strains)\*, *Enterobacter* species, *Escherichia coli*, *Klebsiella* species, *Morganella morganii*\*, *Proteus vulgaris*\*, *Providencia rettgeri*\*, *Pseudomonas aeruginosa*

(3) **Intra-abdominal infections.** *Enterococcus faecalis*, *Staphylococcus aureus* (penicillinase-producing strains)\*, *Staphylococcus epidermidis*, *Citrobacter* species, *Enterobacter* species, *Escherichia coli*, *Klebsiella* species, *Morganella morganii*\*, *Proteus* species, *Pseudomonas aeruginosa*, *Bifidobacterium* species, *Clostridium* species, *Eubacterium* species, *Peptococcus* species, *Peptostreptococcus* species, *Propionibacterium* species\*, *Bacteroides* species including *B. fragilis*, *Fusobacterium* species

(4) **Gynecologic infections.** *Enterococcus faecalis*, *Staphylococcus aureus* (penicillinase-producing strains)\*, *Staphylococcus epidermidis*, *Streptococcus agalactiae* (Group B streptococci), *Enterobacter* species\*, *Escherichia coli*, *Gardnerella vaginalis*, *Klebsiella* species\*, *Proteus* species, *Bifidobacterium* species\*, *Peptococcus* species\*, *Peptostreptococcus* species, *Propionibacterium* species\*, *Bacteroides* species including *B. fragilis*\*

(5) **Bacterial septicemia.** *Enterococcus faecalis*, *Staphylococcus aureus* (penicillinase-producing strains), *Enterobacter* species, *Escherichia coli*, *Klebsiella* species, *Pseudomonas aeruginosa*, *Serratia* species\*, *Bacteroides* species including *B. fragilis*\*

(6) **Bone and joint infections.** *Enterococcus faecalis*, *Staphylococcus aureus* (penicillinase-producing strains), *Staphylococcus epidermidis*, *Enterobacter* species, *Pseudomonas aeruginosa*

(7) **Skin and skin structure infections.** *Enterococcus faecalis*, *Staphylococcus aureus* (penicillinase-producing strains), *Staphylococcus epidermidis*, *Acinetobacter* species, *Citrobacter* species, *Enterobacter* species, *Escherichia coli*, *Klebsiella* species, *Morganella morganii*, *Proteus vulgaris*, *Providencia rettgeri*\*, *Pseudomonas aeruginosa*, *Serratia* species, *Peptococcus* species, *Peptostreptococcus* species, *Bacteroides* species including *B. fragilis*, *Fusobacterium* species\*

(8) **Endocarditis.** *Staphylococcus aureus* (penicillinase-producing strains)

(9) **Polymicrobial infections.** PRIMAXIN I.V. is indicated for polymicrobial infections including those in which *S. pneumoniae* (pneumonia, septicemia), *S. pyogenes* (skin and skin structure), or nonpenicillinase-producing *S. aureus* is one of the causative organisms. However, monobacterial infections due to these organisms are usually treated with narrower spectrum antibiotics, such as penicillin G.

PRIMAXIN I.V. is not indicated in patients with meningitis because safety and efficacy have not been established.

For Pediatric Use information, See **PRECAUTIONS**, *Pediatric Use*, and **DOSAGE AND ADMINISTRATION** sections.

Because of its broad spectrum of bactericidal activity against gram-positive and gram-negative aerobic and anaerobic bacteria, PRIMAXIN I.V. is useful for the treatment of mixed infections and as presumptive therapy prior to the identification of the causative organisms.

Although clinical improvement has been observed in patients with cystic fibrosis, chronic pulmonary disease, and lower respiratory tract infections caused by *Pseudomonas aeruginosa*, bacterial eradication may not necessarily be achieved.

As with other beta-lactam antibiotics, some strains of *Pseudomonas aeruginosa* may develop resistance fairly rapidly during treatment with PRIMAXIN I.V. During therapy of *Pseudomonas aeruginosa* infections, periodic susceptibility testing should be done when clinically appropriate.

Infections resistant to other antibiotics, for example, cephalosporins, penicillin, and aminoglycosides, have been shown to respond to treatment with PRIMAXIN I.V.

## CONTRAINDICATIONS

PRIMAXIN I.V. is contraindicated in patients who have shown hypersensitivity to any component of this product.

\* Efficacy for this organism in this organ system was studied in fewer than 10 infections.

## WARNINGS

SERIOUS AND OCCASIONALLY FATAL HYPERSENSITIVITY (ANAPHYLACTIC) REACTIONS HAVE BEEN REPORTED IN PATIENTS RECEIVING THERAPY WITH BETA-LACTAMS. THESE REACTIONS ARE MORE APT TO OCCUR IN PERSONS WITH A HISTORY OF SENSITIVITY TO MULTIPLE ALLERGENS.

THERE HAVE BEEN REPORTS OF PATIENTS WITH A HISTORY OF PENICILLIN HYPERSENSITIVITY WHO HAVE EXPERIENCED SEVERE HYPERSENSITIVITY REACTIONS WHEN TREATED WITH ANOTHER BETA-LACTAM. BEFORE INITIATING THERAPY WITH PRIMAXIN I.V., CAREFUL INQUIRY SHOULD BE MADE CONCERNING PREVIOUS HYPERSENSITIVITY REACTIONS TO PENICILLINS, CEPHALOSPORINS, OTHER BETA-LACTAMS, AND OTHER ALLERGENS. IF AN ALLERGIC REACTION OCCURS, PRIMAXIN SHOULD BE DISCONTINUED.

**SERIOUS ANAPHYLACTIC REACTIONS REQUIRE IMMEDIATE EMERGENCY TREATMENT WITH EPINEPHRINE, OXYGEN, INTRAVENOUS STEROIDS, AND AIRWAY MANAGEMENT, INCLUDING INTUBATION, MAY ALSO BE ADMINISTERED AS INDICATED.**

Seizures and other CNS adverse experiences, such as confusional states and myoclonic activity, have been reported during treatment with PRIMAXIN I.V. (See **PRECAUTIONS**.)

**Pseudomembranous colitis has been reported with nearly all antibacterial agents, including imipenem-cilastatin sodium, and may range in severity from mild to life threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhea subsequent to the administration of antibacterial agents.**

Treatment with antibacterial agents alters the normal flora of the colon and may permit overgrowth of clostridia. Studies indicate that a toxin produced by *Clostridium difficile* is one primary cause of "antibiotic-associated colitis".

After the diagnosis of pseudomembranous colitis has been established, therapeutic measures should be initiated. Mild cases of pseudomembranous colitis usually respond to drug discontinuation alone. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation and treatment with an antibacterial drug clinically effective against *C. difficile* colitis.

## PRECAUTIONS

### General

CNS adverse experiences such as confusional states, myoclonic activity, and seizures have been reported during treatment with PRIMAXIN I.V., especially when recommended dosages were exceeded. These experiences have occurred most commonly in patients with CNS disorders (e.g., brain lesions or history of seizures) and/or compromised renal function. However, there have been reports of CNS adverse experiences in patients who had no recognized or documented underlying CNS disorder or compromised renal function.

When recommended doses were exceeded, adult patients with creatinine clearances of  $\leq 20$  mL/min/1.73 m<sup>2</sup>, whether or not undergoing hemodialysis, had a higher risk of seizure activity than those without impairment of renal function. Therefore, close adherence to the dosing guidelines for these patients is recommended. (See **DOSAGE AND ADMINISTRATION**.)

Patients with creatinine clearances of  $\leq 5$  mL/min/1.73 m<sup>2</sup> should not receive PRIMAXIN I.V. unless hemodialysis is instituted within 48 hours.

For patients on hemodialysis, PRIMAXIN I.V. is recommended only when the benefit outweighs the potential risk of seizures.

Close adherence to the recommended dosage and dosage schedules is urged, especially in patients with known factors that predispose to convulsive activity. Anticonvulsant therapy should be continued in patients with known seizure disorders. If focal tremors, myoclonus, or seizures occur, patients should be evaluated neurologically, placed on anticonvulsant therapy if not already instituted, and the dosage of PRIMAXIN I.V. re-examined to determine whether it should be decreased or the antibiotic discontinued.

As with other antibiotics, prolonged use of PRIMAXIN I.V. may result in overgrowth of nonsusceptible organisms. Repeated evaluation of the patient's condition is essential. If superinfection occurs during therapy, appropriate measures should be taken.

### Laboratory Tests

While PRIMAXIN I.V. possesses the characteristic low toxicity of the beta-lactam group of antibiotics, periodic assessment of organ system functions, including renal, hepatic, and hematopoietic, is advisable during prolonged therapy.

### *Drug Interactions*

Generalized seizures have been reported in patients who received ganciclovir and PRIMAXIN. These drugs should not be used concomitantly unless the potential benefits outweigh the risks.

Since concomitant administration of PRIMAXIN and probenecid results in only minimal increases in plasma levels of imipenem and plasma half-life, it is not recommended that probenecid be given with PRIMAXIN.

PRIMAXIN should not be mixed with or physically added to other antibiotics. However, PRIMAXIN may be administered concomitantly with other antibiotics, such as aminoglycosides.

### *Carcinogenesis, Mutagenesis, Impairment of Fertility*

Long term studies in animals have not been performed to evaluate carcinogenic potential of imipenem-cilastatin. Genetic toxicity studies were performed in a variety of bacterial and mammalian tests *in vivo* and *in vitro*. The tests used were: V79 mammalian cell mutagenesis assay (imipenem-cilastatin sodium alone and imipenem alone), Ames test (cilastatin sodium alone and imipenem alone), unscheduled DNA synthesis assay (imipenem-cilastatin sodium) and *in vivo* mouse cytogenetics test (imipenem-cilastatin sodium). None of these tests showed any evidence of genetic alterations.

Reproductive tests in male and female rats were performed with imipenem-cilastatin sodium at dosage levels up to 11 times<sup>††</sup> the usual human dose of the intravenous formulation (on a mg/kg basis). Slight decreases in live fetal body weight were restricted to the highest dosage level. No other adverse effects were observed on fertility, reproductive performance, fetal viability, growth or postnatal development of pups. Similarly, no adverse effects on the fetus or on lactation were observed when imipenem-cilastatin sodium was administered to rats late in gestation.

### *Pregnancy: Teratogenic Effects*

*Pregnancy Category C:* Teratology studies with cilastatin sodium in rabbits and rats at 6 to 20 times<sup>††</sup> the maximum recommended human dose of the intravenous formulation of imipenem-cilastatin sodium (50 mg/kg/day<sup>††</sup>), respectively, showed no evidence of adverse effect on the fetus. No evidence of teratogenicity was observed in rabbits and rats given imipenem at doses up to 1 and 18 times<sup>††</sup> the maximum recommended daily human dose of the intravenous formulation of imipenem-cilastatin sodium, respectively.

Teratology studies with imipenem-cilastatin sodium at doses up to 11 times<sup>††</sup> the usual recommended human dose of the intravenous formulation (30 mg/kg/day<sup>††</sup>) in pregnant mice and rats during the period of major organogenesis revealed no evidence of teratogenicity.

Imipenem-cilastatin sodium, when administered to pregnant rabbits at dosages equivalent to the usual human dose of the intravenous formulation and higher, caused body weight loss, diarrhea, and maternal deaths. When comparable doses of imipenem-cilastatin sodium were given to non-pregnant rabbits, body weight loss, diarrhea, and deaths were also observed. This intolerance is not unlike that seen with other beta-lactam antibiotics in this species and is probably due to alteration of gut flora.

A teratology study in pregnant cynomolgus monkeys given imipenem-cilastatin sodium at doses of 40 mg/kg/day (bolus intravenous injection) or 160 mg/kg/day (subcutaneous injection) resulted in maternal toxicity including emesis, inappetence, body weight loss, diarrhea, abortion, and death in some cases. In contrast, no significant toxicity was observed when non-pregnant cynomolgus monkeys were given doses of imipenem-cilastatin sodium up to 180 mg/kg/day (subcutaneous injection). When doses of imipenem-cilastatin sodium (approximately 100 mg/kg/day or approximately 2 times<sup>††</sup> the maximum recommended daily human dose of the intravenous formulation) were administered to pregnant cynomolgus monkeys at an intravenous infusion rate which mimics human clinical use, there was minimal maternal intolerance (occasional emesis), no maternal deaths, no evidence of teratogenicity, but an increase in embryonic loss relative to control groups.

There are, however, no adequate and well-controlled studies in pregnant women. PRIMAXIN I.V. should be used during pregnancy only if the potential benefit justifies the potential risk to the mother and fetus.

### *Nursing Mothers*

It is not known whether imipenem-cilastatin sodium is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when PRIMAXIN I.V. is administered to a nursing woman.

### *Pediatric Use*

Use of PRIMAXIN I.V. in pediatric patients, neonates to 16 years of age, is supported by evidence from adequate and well-controlled studies of PRIMAXIN I.V. in adults and by the following clinical studies and

<sup>††</sup> Based on patient weight of 70 kg.

published literature in pediatric patients: Based on published studies of 178\*\* pediatric patients  $\geq 3$  months of age (with non-CNS infections), the recommended dose of PRIMAXIN I.V. is 15-25 mg/kg/dose administered every six hours. Doses of 25 mg/kg/dose in patients 3 months to  $< 3$  years of age, and 15 mg/kg/dose in patients 3-12 years of age were associated with mean trough plasma concentrations of imipenem of  $1.1 \pm 0.4$   $\mu\text{g/mL}$  and  $0.6 \pm 0.2$   $\mu\text{g/mL}$  following multiple 60-minute infusions, respectively; trough urinary concentrations of imipenem were in excess of 10  $\mu\text{g/mL}$  for both doses. These doses have provided adequate plasma and urine concentrations for the treatment of non-CNS infections. Based on studies in adults, the maximum daily dose for treatment of infections with fully susceptible organisms is 2.0 g per day, and of infections with moderately susceptible organisms (primarily some strains of *P. aeruginosa*) is 4.0 g/day. (See Table 1, **DOSAGE AND ADMINISTRATION.**) Higher doses (up to 90 mg/kg/day in older children) have been used in patients with cystic fibrosis. (See **DOSAGE AND ADMINISTRATION.**)

Based on studies of 135\*\*\* pediatric patients  $\leq 3$  months of age (weighing  $\geq 1,500$  gms), the following dosage schedule is recommended for non-CNS infections:

<1 wk of age: 25 mg/kg every 12 hrs

1-4 wks of age: 25 mg/kg every 8 hrs

4 wks-3 mos. of age: 25 mg/kg every 6 hrs.

In a published dose-ranging study of smaller premature infants (670-1,890 gms) in the first week of life, a dose of 20 mg/kg q12h by 15-30 minutes infusion was associated with mean peak and trough plasma imipenem concentrations of 43  $\mu\text{g/mL}$  and 1.7  $\mu\text{g/mL}$  after multiple doses, respectively. However, moderate accumulation of cilastatin in neonates may occur following multiple doses of PRIMAXIN I.V. The safety of this accumulation is unknown.

PRIMAXIN I.V. is not recommended in pediatric patients with CNS infections because of the risk of seizures.

PRIMAXIN I.V. is not recommended in pediatric patients  $< 30$  kg with impaired renal function, as no data are available.

## ADVERSE REACTIONS

### Adults

PRIMAXIN I.V. is generally well tolerated. Many of the 1,723 patients treated in clinical trials were severely ill and had multiple background diseases and physiological impairments, making it difficult to determine causal relationship of adverse experiences to therapy with PRIMAXIN I.V.

### Local Adverse Reactions

Adverse local clinical reactions that were reported as possibly, probably, or definitely related to therapy with PRIMAXIN I.V. were:

Phlebitis/thrombophlebitis — 3.1%

Pain at the injection site — 0.7%

Erythema at the injection site — 0.4%

Vein induration — 0.2%

Infused vein infection — 0.1%

### Systemic Adverse Reactions

The most frequently reported systemic adverse clinical reactions that were reported as possibly, probably, or definitely related to PRIMAXIN I.V. were nausea (2.0%), diarrhea (1.8%), vomiting (1.5%), rash (0.9%), fever (0.5%), hypotension (0.4%), seizures (0.4%) (see **PRECAUTIONS**), dizziness (0.3%), pruritus (0.3%), urticaria (0.2%), somnolence (0.2%).

Additional adverse systemic clinical reactions reported as possibly, probably, or definitely drug related occurring in less than 0.2% of the patients or reported since the drug was marketed are listed within each body system in order of decreasing severity: *Gastrointestinal* — pseudomembranous colitis (the onset of pseudomembranous colitis symptoms may occur during or after antibacterial treatment, see **WARNINGS**), hemorrhagic colitis, hepatitis, jaundice, gastroenteritis, abdominal pain, glossitis, tongue papillar hypertrophy, staining of the teeth and/or tongue, heartburn, pharyngeal pain, increased salivation; *Hematologic* — pancytopenia, bone marrow depression, thrombocytopenia, neutropenia, leukopenia, hemolytic anemia; *CNS* — encephalopathy, tremor, confusion, myoclonus, paresthesia, vertigo, headache, psychic disturbances including hallucinations; *Special Senses* — hearing loss, tinnitus, taste perversion; *Respiratory* — chest discomfort, dyspnea, hyperventilation, thoracic spine pain;

\*\* Two patients were less than 3 months of age.

\*\*\* One patient was greater than 3 months of age.

*Cardiovascular* — palpitations, tachycardia; *Skin* — Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme, angioneurotic edema, flushing, cyanosis, hyperhidrosis, skin texture changes, candidiasis, pruritus vulvae; *Body as a whole* — polyarthralgia, asthenia/weakness, drug fever; *Renal* — acute renal failure, oliguria/anuria, polyuria, urine discoloration. The role of PRIMAXIN I.V. in changes in renal function is difficult to assess, since factors predisposing to pre-renal azotemia or to impaired renal function usually have been present.

*Adverse Laboratory Changes*

Adverse laboratory changes without regard to drug relationship that were reported during clinical trials or reported since the drug was marketed were:

*Hepatic:* Increased ALT (SGPT), AST (SGOT), alkaline phosphatase, bilirubin, and LDH

*Hemic:* Increased eosinophils, positive Coombs test, increased WBC, increased platelets, decreased hemoglobin and hematocrit, agranulocytosis, increased monocytes, abnormal prothrombin time, increased lymphocytes, increased basophils

*Electrolytes:* Decreased serum sodium, increased potassium, increased chloride

*Renal:* Increased BUN, creatinine

*Urinalysis:* Presence of urine protein, urine red blood cells, urine white blood cells, urine casts, urine bilirubin, and urine urobilinogen.

*Pediatric Patients*

In studies of 178 pediatric patients ≥3 months of age, the following adverse events were noted:

The Most Common Clinical Adverse Experiences Without Regard to Drug Relationship (Patient Incidence >1%)

Adverse Experience	No. of Patients (%)
<i>Digestive System</i>	
Diarrhea	7* (3.9)
Gastroenteritis	2 (1.1)
Vomiting	2* (1.1)
<i>Skin</i>	
Rash	4 (2.2)
Irritation, I.V. site	2 (1.1)
<i>Urogenital System</i>	
Urine discoloration	2 (1.1)
<i>Cardiovascular System</i>	
Phlebitis	4 (2.2)
*One patient had both vomiting and diarrhea and is counted in each category.	

In studies of 135 patients (newborn to 3 months of age), the following adverse events were noted:

The Most Common Clinical Adverse Experiences Without Regard to Drug Relationship (Patient Incidence >1%)

Adverse Experience	No. of Patients (%)
<i>Digestive System</i>	
Diarrhea	4 (3.0%)
Oral Candidiasis	2 (1.5%)
<i>Skin</i>	
Rash	2 (1.5%)
<i>Urogenital System</i>	
Oliguria/anuria	3 (2.2%)
<i>Cardiovascular System</i>	
Tachycardia	2 (1.5%)
<i>Nervous System</i>	
Convulsions	8 (5.9%)

Patients (≥3 Months of Age) With Normal Pretherapy but Abnormal During Therapy Laboratory Values

Laboratory Parameter	Abnormality	No. of Patients With Abnormalities/ No. of Patients With Lab Done (%)
Hemoglobin	Age <5 mos.: <10 gm % 6 mos.-12 yrs.: <11.5 gm%	19/129 (14.7)
Hematocrit	Age <5 mos.: <30 vol % 6 mos.-12 yrs.: <34.5 vol %	23/129 (17.8)
Neutrophils	≤1000/mm <sup>3</sup> (absolute)	4/123 (3.3)
Eosinophils	≥7%	15/117 (12.8)
Platelet Count	≥500 ths/mm <sup>3</sup>	16/119 (13.4)
Urine Protein	≥1	8/97 (8.2)
Serum Creatinine	>1.2 mg/dL	0/105 (0)
BUN	>22 mg/dL	0/108 (0)
AST (SGOT)	>36 IU/L	14/78 (17.9)
ALT (SGPT)	>30 IU/L	10/93 (10.8)

Patients (&lt;3 Months of Age) With Normal Pretherapy but Abnormal During Therapy Laboratory Values

Laboratory Parameter	No. of Patients With Abnormalities* (%)
Eosinophil Count↑	11 (9.0%)
Hematocrit↓	3 (2.0%)
Hematocrit↑	1 (1.0%)
Platelet Count↑	5 (4.0%)
Platelet Count↓	2 (2.0%)
Serum Creatinine↑	5 (5.0%)
Bilirubin↑	3 (3.0%)
Bilirubin↓	1 (1.0%)
AST (SGOT)↑	5 (6.0%)
ALT (SGPT)↑	3 (3.0%)
Serum Alkaline Phosphate↑	2 (3.0%)

\*The denominator used for percentages was the number of patients for whom the test was performed during or post-treatment and, therefore, varies by test.

Examination of published literature and spontaneous adverse event reports suggested a similar spectrum of adverse events in adult and pediatric patients.

## OVERDOSAGE

The acute intravenous toxicity of imipenem-cilastatin sodium in a ratio of 1:1 was studied in mice at doses of 751 to 1359 mg/kg. Following drug administration, ataxia was rapidly produced and clonic convulsions were noted in about 45 minutes. Deaths occurred within 4-56 minutes at all doses.

The acute intravenous toxicity of imipenem-cilastatin sodium was produced within 5-10 minutes in rats at doses of 771 to 1583 mg/kg. In all dosage groups, females had decreased activity, bradypnea, and ptosis with clonic convulsions preceding death; in males, ptosis was seen at all dose levels while tremors and clonic convulsions were seen at all but the lowest dose (771 mg/kg). In another rat study, female rats showed ataxia, bradypnea, and decreased activity in all but the lowest dose (550 mg/kg); deaths were preceded by clonic convulsions. Male rats showed tremors at all doses and clonic convulsions and ptosis were seen at the two highest doses (1130 and 1734 mg/kg). Deaths occurred between 6 and 88 minutes with doses of 771 to 1734 mg/kg.

In the case of overdosage, discontinue PRIMAXIN I.V., treat symptomatically, and institute supportive measures as required. Imipenem-cilastatin sodium is hemodialyzable. However, usefulness of this procedure in the overdosage setting is questionable.

**DOSAGE AND ADMINISTRATION****Adults**

The dosage recommendations for PRIMAXIN I.V. represent the quantity of imipenem to be administered. An equivalent amount of cilastatin is also present in the solution. Each 125 mg, 250 mg, or 500 mg dose should be given by intravenous administration over 20 to 30 minutes. Each 750 mg or 1000 mg dose should be infused over 40 to 60 minutes. In patients who develop nausea during the infusion, the rate of infusion may be slowed.

The total daily dosage for PRIMAXIN I.V. should be based on the type or severity of infection and given in equally divided doses based on consideration of degree of susceptibility of the pathogen(s), renal function, and body weight. Adult patients with impaired renal function, as judged by creatinine clearance  $\leq 70$  mL/min/1.73 m<sup>2</sup>, require adjustment of dosage as described in the succeeding section of these guidelines.

***Intravenous Dosage Schedule for Adults with Normal Renal Function and Body Weight  $\geq 70$  kg***

Doses cited in Table I are based on a patient with normal renal function and a body weight of 70 kg. These doses should be used for a patient with a creatinine clearance of  $\geq 71$  mL/min/1.73 m<sup>2</sup> and a body weight of  $\geq 70$  kg. A reduction in dose must be made for a patient with a creatinine clearance of  $\leq 70$  mL/min/1.73 m<sup>2</sup> and/or a body weight less than 70 kg. (See Tables II and III.)

Dosage regimens in column A of Table I are recommended for infections caused by fully susceptible organisms which represent the majority of pathogenic species. Dosage regimens in column B of Table I are recommended for infections caused by organisms with moderate susceptibility to imipenem, primarily some strains of *P. aeruginosa*.

TABLE I  
INTRAVENOUS DOSAGE SCHEDULE FOR ADULTS WITH  
NORMAL RENAL FUNCTION AND BODY WEIGHT  $\geq 70$  kg

Type or Severity of Infection	A Fully susceptible organisms including gram-positive and gram-negative aerobes and anaerobes	B Moderately susceptible organisms, primarily some strains of <i>P. aeruginosa</i>
Mild	250 mg q6h (TOTAL DAILY DOSE = 1.0g)	500 mg q6h (TOTAL DAILY DOSE = 2.0g)
Moderate	500 mg q8h (TOTAL DAILY DOSE = 1.5g) or 500 mg q6h (TOTAL DAILY DOSE = 2.0g)	500 mg q6h (TOTAL DAILY DOSE = 2.0g) or 1 g q8h (TOTAL DAILY DOSE = 3.0g)
Severe, life threatening only	500 mg q6h (TOTAL DAILY DOSE = 2.0g)	1 g q8h (TOTAL DAILY DOSE = 3.0g) or 1 g q6h (TOTAL DAILY DOSE = 4.0g)
Uncomplicated urinary tract infection	250 mg q6h (TOTAL DAILY DOSE = 1.0g)	250 mg q6h (TOTAL DAILY DOSE = 1.0g)
Complicated urinary tract infection	500 mg q6h (TOTAL DAILY DOSE = 2.0g)	500 mg q6h (TOTAL DAILY DOSE = 2.0g)

Due to the high antimicrobial activity of PRIMAXIN I.V., it is recommended that the maximum total daily dosage not exceed 50 mg/kg/day or 4.0 g/day, whichever is lower. There is no evidence that higher doses provide greater efficacy. However, patients over twelve years of age with cystic fibrosis and normal renal function have been treated with PRIMAXIN I.V. at doses up to 90 mg/kg/day in divided doses, not exceeding 4.0 g/day.

***Reduced Intravenous Schedule for Adults with Impaired Renal Function and/or Body Weight  $< 70$  kg***

Patients with creatinine clearance of  $\leq 70$  mL/min/1.73 m<sup>2</sup> and/or body weight less than 70 kg require dosage reduction of PRIMAXIN I.V. as indicated in the tables below. Creatinine clearance may be calculated from serum creatinine concentration by the following equation:

$$T_{cc} \text{ (Males)} = \frac{(\text{wt. in kg}) (140 - \text{age})}{(72) (\text{creatinine in mg/dL})}$$

$$T_{cc} \text{ (Females)} = 0.85 \times \text{above value}$$

To determine the dose for adults with impaired renal function and/or reduced body weight:

1. Choose a total daily dose from Table I based on infection characteristics.
2. a) If the total daily dose is 1.0 g, 1.5 g, or 2.0 g, use the appropriate subsection of Table II and continue with step 3.  
 b) If the total daily dose is 3.0 g or 4.0 g, use the appropriate subsection of Table III and continue with step 3.
3. From Table II or III:
  - a) Select the body weight on the far left which is closest to the patient's body weight (kg).
  - b) Select the patient's creatinine clearance category.
  - c) Where the row and column intersect is the reduced dosage regimen.

**TABLE II**  
 REDUCED INTRAVENOUS DOSAGE OF PRIMAXIN I.V. IN ADULT PATIENTS WITH  
 IMPAIRED RENAL FUNCTION AND/OR BODY WEIGHT <70 kg

and Body Weight (kg) is:	if TOTAL DAILY DOSE from TABLE I is:											
	1.0 g/day				1.5 g/day				2.0 g/day			
	and creatinine clearance (mL/min/1.73 m <sup>2</sup> ) is:				and creatinine clearance (mL/min/1.73 m <sup>2</sup> ) is:				and creatinine clearance (mL/min/1.73 m <sup>2</sup> ) is:			
	≥71	41-70	21-40	6-20	≥71	41-70	21-40	6-20	≥71	41-70	21-40	6-20
	then the reduced dosage regimen (mg) is:				then the reduced dosage regimen (mg) is:				then the reduced dosage regimen (mg) is:			
≥70	250 q6h	250 q8h	250 q12h	250 q12h	500 q8h	250 q6h	250 q8h	250 q12h	500 q6h	500 q8h	250 q6h	250 q12h
60	250 q8h	125 q6h	250 q12h	125 q12h	250 q6h	250 q8h	250 q8h	250 q12h	500 q8h	250 q6h	250 q8h	250 q12h
50	125 q6h	125 q6h	125 q8h	125 q12h	250 q6h	250 q8h	250 q12h	250 q12h	250 q6h	250 q6h	250 q8h	250 q12h
40	125 q6h	125 q8h	125 q12h	125 q12h	250 q8h	125 q6h	125 q8h	125 q12h	250 q6h	250 q8h	250 q12h	250 q12h
30	125 q8h	125 q8h	125 q12h	125 q12h	125 q6h	125 q8h	125 q8h	125 q12h	250 q8h	125 q6h	125 q8h	125 q12h

**TABLE III**  
 REDUCED INTRAVENOUS DOSAGE OF PRIMAXIN I.V. IN ADULT PATIENTS WITH  
 IMPAIRED RENAL FUNCTION AND/OR BODY WEIGHT <70 kg

and Body Weight (kg) is:	if TOTAL DAILY DOSE from TABLE I is:							
	3.0 g/day				4.0 g/day			
	and creatinine clearance (mL/min/1.73 m <sup>2</sup> ) is:				and creatinine clearance (mL/min/1.73 m <sup>2</sup> ) is:			
	≥71	41-70	21-40	6-20	≥71	41-70	21-40	6-20
	then the reduced dosage regimen (mg) is:				then the reduced dosage regimen (mg) is:			
≥70	1000 q8h	500 q6h	500 q8h	500 q12h	1000 q6h	750 q8h	500 q6h	500 q12h
60	750 q8h	500 q8h	500 q8h	500 q12h	1000 q8h	750 q8h	500 q8h	500 q12h
50	500 q6h	500 q8h	250 q6h	250 q12h	750 q8h	500 q6h	500 q8h	500 q12h
40	500 q8h	250 q6h	250 q8h	250 q12h	500 q6h	500 q8h	250 q6h	250 q12h
30	250 q6h	250 q8h	250 q8h	250 q12h	500 q8h	250 q6h	250 q8h	250 q12h

Patients with creatinine clearances of 6 to 20 mL/min/1.73 m<sup>2</sup> should be treated with PRIMAXIN I.V. 125 mg or 250 mg every 12 hours for most pathogens. There may be an increased risk of seizures when doses of 500 mg every 12 hours are administered to these patients.

Patients with creatinine clearance  $\leq 5$  mL/min/1.73 m<sup>2</sup> should not receive PRIMAXIN I.V. unless hemodialysis is instituted within 48 hours. There is inadequate information to recommend usage of PRIMAXIN I.V. for patients undergoing peritoneal dialysis.

#### *Hemodialysis*

When treating patients with creatinine clearances of  $\leq 5$  mL/min/1.73 m<sup>2</sup> who are undergoing hemodialysis, use the dosage recommendations for patients with creatinine clearances of 6-20 mL/min/1.73 m<sup>2</sup>. (See *Reduced Intravenous Dosage Schedule for Adults with Impaired Renal Function and/or Body Weight <70 kg.*) Both imipenem and cilastatin are cleared from the circulation during hemodialysis. The patient should receive PRIMAXIN I.V. after hemodialysis and at 12 hour intervals timed from the end of that hemodialysis session. Dialysis patients, especially those with background CNS disease, should be carefully monitored; for patients on hemodialysis, PRIMAXIN I.V. is recommended only when the benefit outweighs the potential risk of seizures. (See **PRECAUTIONS.**)

#### *Pediatric Patients*

See **PRECAUTIONS, Pediatric Patients.**

For pediatric patients  $\geq 3$  months of age, the recommended dose for non-CNS infections is 15-25 mg/kg/dose administered every six hours. Based on studies in adults, the maximum daily dose for treatment of infections with fully susceptible organisms is 2.0 g per day, and of infections with moderately susceptible organisms (primarily some strains of *P. aeruginosa*) is 4.0 g/day. Higher doses (up to 90 mg/kg/day in older children) have been used in patients with cystic fibrosis.

For pediatric patients  $\leq 3$  months of age (weighing  $\geq 1,500$  gms), the following dosage schedule is recommended for non-CNS infections:

<1 wk of age: 25 mg/kg every 12 hrs

1-4 wks of age: 25 mg/kg every 8 hrs

4 wks-3 mos. of age: 25 mg/kg every 6 hrs.

Doses less than or equal to 500 mg should be given by intravenous infusion over 15 to 30 minutes. Doses greater than 500 mg should be given by intravenous infusion over 40 to 60 minutes.

PRIMAXIN I.V. is not recommended in pediatric patients with CNS infections because of the risk of seizures.

PRIMAXIN I.V. is not recommended in pediatric patients <30 kg with impaired renal function, as no data are available.

## PREPARATION OF SOLUTION

### *Infusion Bottles*

Contents of the infusion bottles of PRIMAXIN I.V. Powder should be restored with 100 mL of diluent (see list of diluents under **COMPATIBILITY AND STABILITY**) and shaken until a clear solution is obtained.

### *Vials*

Contents of the vials must be suspended and transferred to 100 mL of an appropriate infusion solution.

A suggested procedure is to add approximately 10 mL from the appropriate infusion solution (see list of diluents under **COMPATIBILITY AND STABILITY**) to the vial. Shake well and transfer the resulting suspension to the infusion solution container.

**Benzyl alcohol as a preservative has been associated with toxicity in neonates. While toxicity has not been demonstrated in pediatric patients greater than three months of age, small pediatric patients in this age range may also be at risk for benzyl alcohol toxicity. Therefore, diluents containing benzyl alcohol should not be used when PRIMAXIN I.V. is constituted for administration to pediatric patients in this age range.**

**CAUTION: THE SUSPENSION IS NOT FOR DIRECT INFUSION.**

Repeat with an additional 10 mL of infusion solution to ensure complete transfer of vial contents to the infusion solution. **The resulting mixture should be agitated until clear.**

### *ADD-Vantage<sup>®</sup>††† Vials*

See separate INSTRUCTIONS FOR USE OF 'PRIMAXIN I.V.' IN ADD-Vantage<sup>®</sup> VIALS. PRIMAXIN I.V. in ADD-Vantage<sup>®</sup> vials should be reconstituted with ADD-Vantage<sup>®</sup> diluent containers containing 100 mL of either 0.9% Sodium Chloride Injection or 100 mL 5% Dextrose Injection.

††† Registered trademark of Abbott Laboratories, Inc.

**MONOVIAL® Vials**

See separate INSTRUCTIONS FOR USE OF 'PRIMAXIN I.V.' in MONOVIAL® VIALS. PRIMAXIN I.V. in MONOVIAL® vials should be reconstituted using an appropriate diluent in an infusion bag, with a maximum port length of 14 mm.

The MONOVIAL vial is not compatible with the ADD-Vantage® diluent bags.

**COMPATIBILITY AND STABILITY***Before Reconstitution:*

The dry powder should be stored at a temperature below 25°C (77°F).

*Reconstituted Solutions:*

Solutions of PRIMAXIN I.V. range from colorless to yellow. Variations of color within this range do not affect the potency of the product.

PRIMAXIN I.V., as supplied in single use infusion bottles, vials and MONOVIAL® vials and reconstituted with the following diluents (See **PREPARATION OF SOLUTION**), maintains satisfactory potency for 4 hours at room temperature or for 24 hours under refrigeration (5°C). Solutions of PRIMAXIN I.V. should not be frozen.

- 0.9% Sodium Chloride Injection
- 5% or 10% Dextrose Injection
- 5% Dextrose and 0.9% Sodium Chloride Injection
- 5% Dextrose Injection with 0.225% or 0.45% saline solution
- 5% Dextrose Injection with 0.15% potassium chloride solution
- Mannitol 5% and 10%

PRIMAXIN I.V., as supplied in single dose ADD-Vantage® vials and reconstituted with the following diluents (See **PREPARATION OF SOLUTION**), maintains satisfactory potency for 4 hours at room temperature.

- 0.9% Sodium Chloride Injection
- 5% Dextrose Injection

PRIMAXIN I.V. should not be mixed with or physically added to other antibiotics. However, PRIMAXIN I.V. may be administered concomitantly with other antibiotics, such as aminoglycosides.

**HOW SUPPLIED**

PRIMAXIN I.V. is supplied as a sterile powder mixture in single dose containers including vials, infusion bottles, ADD-Vantage® vials, and MONOVIAL® vials containing imipenem (anhydrous equivalent) and cilastatin sodium as follows:

No. 3514 — 250 mg imipenem equivalent and 250 mg cilastatin equivalent and 10 mg sodium bicarbonate as a buffer

**NDC 0006-3514-58** in trays of 25 vials  
(6505-01-332-4793 250 mg, 25's).

No. 3516 — 500 mg imipenem equivalent and 500 mg cilastatin equivalent and 20 mg sodium bicarbonate as a buffer

**NDC 0006-3516-59** in trays of 25 vials  
(6505-01-332-4794 500 mg, 25's).

No. 3517 — 500 mg imipenem equivalent and 500 mg cilastatin equivalent and 20 mg sodium bicarbonate as a buffer

**NDC 0006-3517-75** in trays of 10 infusion bottles  
(6505-01-234-0240 infusion bottle, 10's).

No. 3551 — 250 mg imipenem equivalent and 250 mg cilastatin equivalent and 10 mg sodium bicarbonate as a buffer

**NDC 0006-3551-58** in trays of 25 ADD-Vantage® vials.

No. 3552 — 500 mg imipenem equivalent and 500 mg cilastatin equivalent and 20 mg sodium bicarbonate as a buffer

**NDC 0006-3552-59** in trays of 25 ADD-Vantage® vials  
(6505-01-279-9627 500 mg ADD-Vantage®, 25's).

No. 3666 — 500 mg imipenem equivalent and 500 mg cilastatin equivalent and 20 mg sodium bicarbonate as a buffer

‡ Registered trademark of Becton Dickinson and Company.

PRIMAXIN® I.V. (Imipenem and Cilastatin for Injection)

7882125

NDC 0006-3666-59 in trays of 25 MONOVIAL® vials.

#### REFERENCES

1. National Committee for Clinical Laboratory Standards, Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically-Fourth Edition. Approved Standard NCCLS Document M7-A4, Vol. 17, No. 2 NCCLS, Villanova, PA, 1997.
  2. National Committee for Clinical Laboratory Standards, Performance Standards for Antimicrobial Disk Susceptibility Tests-Sixth Edition. Approved Standard NCCLS Document M2-A6, Vol. 17, No. 1 NCCLS, Villanova, PA, 1997.
  3. National Committee for Clinical Laboratory Standards, Method for Antimicrobial Susceptibility Testing of Anaerobic Bacteria-Third Edition. Approved Standard NCCLS Document M11-A3, Vol. 13, No. 26 NCCLS, Villanova, PA, 1993.
- 

 Dist. by.  
**MERCK & CO., INC.**, West Point, PA 19486, USA

Issued February 1999  
Printed in USA

**CENTER FOR DRUG EVALUATION AND RESEARCH**

***APPLICATION NUMBER:***

**50-587 / S-052**

**MEDICAL REVIEW**

**Medical Officer Review  
Labeling Supplement  
NDA 50,587**

Date of Submission Letter: 8/18/99 (SLR-052)  
Date Received by CDER: 8/23/99  
Date Given to MO: 4/27/00  
Date of MO's Review: 6/1/00  
NDA: 50,587  
Applicant: Merck Research Laboratories  
Drug: Primaxin  
Type of Drug: Carbapenem  
Route of Administration: Intravenous  
Dosage Form: PRIMAXIN I.V. in MONOVIAL (500 mg  
imipenem equivalent and 500 mg cilastin equivalent  
and 20 mg sodium bicarbonate as buffer strength)  
Purpose of Submission: Proposal for labeling changes

**I. Purpose of Submission**

The Applicant has submitted this labeling supplement to request the following changes to the currently approved label for Primaxin:

- A. Revision of the DESCRIPTION, PREPARATION OF SOLUTION, COMPATIBILITY AND STABILITY, and HOW SUPPLIED sections of the label to incorporate the approved MONOVIAL™ Vials text. (The supplemental application for use of PRIMAXIN I.V. in MONOVIAL vials was approved in a letter to the Applicant from the Agency dated July 24, 1998.)
- B. Editorial changes to ADVERSE REACTIONS and PREPARATION OF SOLUTION sections of the label.

**II. Changes Made**

Revisions made to incorporate MONOVIAL text:

A. DESCRIPTION

Current label reads: PRIMAXIN I.V. is buffered to provide solutions in the pH range of

Revised label reads: PRIMAXIN I.V. is buffered to provide solutions in the pH range of 6.5 to 8.5.

*Medical Officer's Comment: This change was not reflected in the proposed label included in the supplemental application for MONOVIAL vial use. However, it was included in the Finished Product Specifications portion of the submission, which has been previously reviewed by the Chemistry team.*

## B. PREPARATION OF SOLUTION

The following additional text has been added:

### *MONOVIAL VIALS*

See separate INSTRUCTIONS FOR USE OF 'PRIMAXIN I.V.' in MONOVIAL VIALS. PRIMAXIN I.V. in MONOVIAL vials should be reconstituted using an appropriate diluent in an infusion bag, with a maximum port length of 14 mm.

The MONOVIAL vial is not compatible with the ADD-Vantage diluent bags.

*Medical Officer's Comment: This statement and the accompanying insert for use of the MONOVIAL vial system provide clear and concise instructions for use.*

## C. COMPATIBILITY AND STABILITY

Current label reads: PRIMAXIN I.V., as supplied in infusion bottles and vials and reconstituted as above with the following diluents, maintains satisfactory potency for four hours at room temperature or for 24 hours under refrigeration (5°C). Solutions of PRIMAXIN I.V. should not be frozen.

- 0.9% Sodium Chloride Injection
- 5% or 10% Dextrose Injection
- 5% Dextrose and 0.9% Sodium Chloride Injection
- 5% Dextrose Injection with 0.225% or 0.45% saline solution
- 5% Dextrose Injection with 0.15% potassium chloride solution
- Mannitol 5% and 10%

PRIMAXIN I.V. is supplied in single dose ADD-Vantage vials and should be prepared as directed in the accompanying INSTRUCTIONS FOR USE OF 'PRIMAXIN I.V.' in ADD-Vantage VIALS using ADD-Vantage diluent containers containing 100 mL of either 0.9% Sodium Chloride Injection or 5% Dextrose Injection. When prepared with either of these diluents, PRIMAXIN I.V. maintains satisfactory potency for 4 hours at room temperature.

Revised label reads: PRIMAXIN I.V., as supplied in single use infusion bottles, vials and MONOVIAL vials and reconstituted with the following

diluents (See **PREPARATION OF SOLUTION**), maintains satisfactory potency for 4 hours at room temperature or for 24 hours under refrigeration (5°C). Solutions of PRIMAXIN I.V. should not be frozen.

0.9% Sodium Chloride Injection

5% or 10% Dextrose Injection

5% Dextrose and 0.9% Sodium Chloride Injection

5% Dextrose Injection with 0.225% or 0.45% saline solution

5% Dextrose Injection with 0.15% potassium chloride solution

Mannitol 5% and 10%

PRIMAXIN I.V., as supplied in single dose ADD-Vantage vials and reconstituted with the following diluents (See **PREPARATION OF SOLUTION**), maintains satisfactory potency for 4 hours at room temperature.

0.9% Sodium Chloride Injection

5% Dextrose Injection

*Medical Officer's Comment: The proposed change is reasonable and improves the clarity of this section.*

#### D. HOW SUPPLIED

Current label reads: PRIMAXIN I.V. is supplied as a sterile powder mixture in vials and infusion bottles containing imipenem (anhydrous equivalent) and cilastatin sodium as follows:

Revised label reads: PRIMAXIN I.V. is supplied as a sterile powder in single dose containers including vials, infusion bottles, ADD-Vantage vials, and MONOVIAL vials containing imipenem (anhydrous equivalent) and cilastatin sodium as follows:

The following additional text has been added to the end of the section:

No. 3666 – 500 mg imipenem equivalent and 500 mg cilastatin equivalent and 20 mg sodium bicarbonate as a buffer

NDC 0006-3666-59 in trays of 25 MONOVIAL vials.

*Medical Officer's Comment: The proposed change and addition of text to this section is appropriate.*

Additional editorial changes:

#### A. ADVERSE REACTIONS

Editorial changes in this section consist of changes to punctuation and reformatting of Table titles.

#### B. PREPARATION OF SOLUTION

Editorial changes in this section consist of changes to punctuation.

*Medical Officer's Comment: The proposed changes are editorial and do not alter the meaning of the text.*

### III. Conclusions

The Applicant has appropriately revised labeling to reflect use of the approved MONOVIAL Vial. In addition, a number of editorial changes have been made, which are acceptable. It is recommended that the supplement be approved.

---

Jean Marie Mulinde, M.D.  
Medical Officer/HFD-520

NDA 50,587  
HFD-520/Division Files  
HFD-520/DepDivDir/LGavrilovich  
HFD-520/MedRev/JMulinde  
HFD-520/ProjMan/FLeSane

Concurrence Only:  
HFD-520/DivDir/GChikami  
HFD-520/MedTeamLead/ARakowsky  
HFD-520/ActingChemTeamLead/SPagay

/s/

-----  
Jean Mulinde  
7/7/00 08:22:33 AM  
MEDICAL OFFICER

Alexander Rakowsky  
7/7/00 09:31:43 AM  
MEDICAL OFFICER

Gary Chikami  
2/8/01 04:31:21 PM  
MEDICAL OFFICER

Maureen Dillon-Parker  
2/14/01 02:09:24 PM  
CSO