

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

21-043

FINAL PRINTED LABELING

MAXIMUM STRENGTH

RID[®]

MOUSSE

Easy To Apply
Non-drip Formula
Pleasant Scent
KILLS LICE

Pfizer

KEEP THIS AND ALL OTHERS OUT OF THE REACH OF CHILDREN. See back panel for additional warnings.
Distributed by: CONSUMER HEALTHCARE GROUP, PFEIZER INC., NEW YORK, NY 10017

Drug Facts

Active Ingredients
(calculated without propellant)
Piperonyl butoxide (4%), Pyrethrum extract*
(equivalent to 0.33% pyrethrins)

Purpose lice treatment

Uses treats head, pubic (crab), and body lice

Warnings

- For external use only
- Flammable: Keep away from fire or flame

Do not use near the eyes

Ask a doctor before use if you have an allergy to ragweed

When using this product

- keep out of eyes when rinsing hair
- close eyes tightly and do not open eyes until product is rinsed out
- protect eyes with washcloth, towel or other suitable method
- if product gets into the eyes, immediately flush with water
- do not use inside the nose, mouth, or vagina. Irritation may occur.
- do not inhale; use in a well ventilated area
- do not puncture or incinerate. Contents under pressure.

Stop use and ask a doctor if

- skin irritation or infection is present or develops
- inflammation of eyebrows or eyelashes occurs

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

NET WT. 5.5 OZ (156 g)

Drug Facts (continued)

Directions

- Important: Read warnings before using
- shake well before using
- holding the can upside down, apply RID[®] Mousse to dry hair or other affected area. Massage until all the hair is thoroughly wet with product. For best results against head lice, see complete Directions in the Consumer Information Insert.
- allow product to remain on the hair for 10 minutes but no longer
- wash area thoroughly with warm water and soap or shampoo
- a fine-toothed comb or a special lice/nit removing comb may be used to help remove dead lice or their eggs (nits) from hair
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice

Other Information

- store at 20°-25°C (68°-77°F)
- do not store at temperature above 43°C (110°F)
- keep in a cool place out of the sun

Inactive Ingredients cetearyl alcohol, isobutane, PEG-20 stearate, propane, propylene glycol, purified water, quaternium-52, SD Alcohol 3-C (26.5% w/w)

For Questions? call 1-800-RID LICE
(1-800-743-5423)

Visit us at <http://www.licerid.com>

*50% extract (not USP)

U.S. Pat. 6,783,202

SM 2512001P

EXP

LOT




ENLARGED FOR PROOFREADING

attach
task r

128 CODE

MA



Instruc

RID[®] MOUSSE

**MAXIMUM
STRENGTH**



Drug Facts

Active ingredients (calculated without propellant) Purpose
Piperonyl butoxide (4%) *Lice treatment*
Pyrethrum extract* (equivalent to 0.53% pyrethrin) *Lice treatment*

Uses: treats head, pubic (crab), and body lice

Warnings

- For external use only
- Flammable: Keep away from fire or flame

Do not use near the eyes

Ask a doctor before use if you have an allergy to pyrethroid

When using this product

- keep out of eyes when rinsing hair
- close eyes tightly and do not open eyes until product is rinsed out
- protect eyes with washcloth, towel or other suitable method
- if product gets into the eyes, immediately flush with water
- do not use inside the nose, mouth, or vagina. Irritation may occur.
- do not inhale; use in a well ventilated area
- do not puncture or incinerate. Contents under pressure.

Stop use and ask a doctor if

- skin irritation or infection is present or develops
- inflammation of eyebrows or eyelashes occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions


- Important: Read warnings before using
- shake well before using
- holding the can upside down, apply RID[®] Mousse to dry hair or other affected area. Massage until all the hair is thoroughly wet with product. For best results against head lice, see complete Directions in the Consumer Information insert.
- allow product to remain on the hair for 10 minutes but no longer
- wash area thoroughly with warm water and soap or shampoo
- a fine-toothed comb or a special comb/ removing comb may be used to help remove dead lice or their eggs (nits) from hair
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice

Other information

- store at 20°-25°C (68°-77°F)
- do not store at temperature above 43°C (110°F)
- keep in a cool place out of the sun

Inactive ingredients: cetylalcohol, isobutane, PEG-20 stearate, propane, propylene glycol, purified water, quaternium-15, SO Alcohol 3-C (25.5% w/w)

**Questions? call 1-800-RID LICE
(1-800-743-5423)**



Distributed By
CONSUMER HEALTH CARE GROUP,
PFIZER INC., NEW YORK, NY 10017

U.S. Pat. 5,783,202 SM 4012000SP
*50% extract (not USP)

EXP

LOT

Easy
Non-
Please

KIT

KEEP THIS AND A
See box

ENLARGED FOR PROOFREADING

11011

MAXIMUM STRENGTH



MOUSSE

Instrucciones En Español Incluidas

MAXIMUM STRENGTH



MOUSSE




Made By
Pfizer Health Care Group,
Inc., New York, NY 10017
L 5,783,202 SM 40120005P
Product of USA

MAXIMUM STRENGTH



MOUSSE



KILLS LICE

NET WT.
5.5 OZ (156 g)

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.
See back panel for additional warnings.

MAXIMUM STRENGTH



MOUSSE



Plus Patented
Egg Removal
Comb



Visit us at <http://www.liceoid.com>

ENLARGED FOR PROOFREADING

FORM 101 2100000 01/01/01 101