

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER:
NDA 20-958 / S-002

APPROVAL LETTER
w/ APPROVED LABELING

7/5/2001
 Jo-958
 SCF-002

REV	DATE	DESCRIPTION	BY
1	05/31/01	CARTON, SEAL, END	3-1/8 X 3-1/8 X 3-3/4

NOTES

- 1) DRAWING SHOWN PRINT SIDE UP
- 2) ALL DIMENSIONS IN INCHES
- 3) TOLERANCES UNLESS SPECIFIED FRACTIONS 1/32, ANGLES 1/2°
- 4) BASED ON PCI DWG. NO. L30126 (10-8-98)
- 5) ALL LIVE COPY & TEXT TO BE A MINIMUM OF 1/16" FROM ALL EDGES & CUTS.
- 6) LDT & EXP. DATE AREA
- 7) VISIONS BARCODE AREA
- 8) NO VARNISH AREA
- 9) ---- NOT PRINT




Labeling Format Information:

Fonts: Univers condensed, bold and bold italic.

Drug Facts:	9 pt	Leading:	6.5 pt
Header:	8 pt	Bullets:	5 pt
Subheader:	6 pt	Barlines:	2.5 pt
Body Text:	6 pt	Hairlines:	.5 pt
Drug Facts (continued):	8 pt		

Horizontal Scale: 94% Average Kerning: -4

JOB DESCRIPTION CTN PEPCID COMP BERRY CHEW TAB 5 FILE # 3003222 VERSION A-1 DATE May 31, 2001 DIE VINYL FC-C-752.00 PS IMAGE # M298/M298/ M299/M245/LIVE ART KEY CONTACT Tony Nucera COMPARED BY: _____ DESIGN PROGRAM ILLUS. ver. 8.0 MAXIMUM OF FIVE COLORS PER COMPONENT  THIS DESIGN BLOCK MUST BE SAVED WITH FILE AND PRINTED ON THERMAL.		PRINTED GRAPHIC ANALYST UPC # PART NUMBER PCNMPA NUMBER FCCI PACKAGING LOCATION COLOR STANDARD VALIDITY DATE	McNeil Johnson & Johnson - MERCK ARTWORK APPROVAL INITIAL BLOCK INITIALS _____ DATE _____ REGULATORY MEDICAL
---	--	---	--

- NOTES:
- 1) DRAWING SHOWN PRINT SIDE UP
 - 2) ALL DIMENSIONS IN INCHES
 - 3) TOLERANCES UNLESS SPECIFIED: FRACTIONS ±1/32, ANGLES ±1°
 - 4) BASED ON PCI DWG. NO. L3240IR3 (5-11-00)
 - 5) ALL LIVE COPY & TEXT TO BE A MINIMUM OF 1/16" FROM ALL SCORES & CUTS.
 - 6) [Cross-hatch symbol] NO PRINT AREA
 - 7) [Diagonal lines symbol] NO VARNISH/NO PRINT AREA
 - 8) [Diagonal lines symbol] NO PRINT AREA RESERVED FOR LOT & EXP DATE ONLINE PRINTING
 - 9) [Dashed line symbol] NO PRINT

REV	DATE	BY	DESCRIPTION
01			
02			
03			
04			
05			
06			
07			
08			
09			
10			

WHAT PATIENTS SHOULD KNOW ABOUT:

Pepcid Complete
Acid Reducer + Anacid

FOR PROFESSIONAL USE ONLY • NOT FOR RETAIL SALE

Works with Just One Tablet!

Relieves Acid Reflux
Reduces Stomach Acid

Relieves Heartburn Due To Acid Indigestion

25 *25 mg* *10 mg* flavored Chewable Tablets

HOW TO TAKE:

Take 1 tablet before breakfast and one additional tablet before dinner.

HOW TO TAKE:

Take 1 tablet before breakfast and one additional tablet before dinner.

CAUTIONS:

Do not use with other antacid products or acid reducers.

Do not use with other heartburn products or acid reducers.

Do not use with other antacid products or acid reducers.

Do not use with other heartburn products or acid reducers.

HOW TO TAKE:

Take 1 tablet before breakfast and one additional tablet before dinner.

CAUTIONS:

Do not use with other antacid products or acid reducers.

Do not use with other heartburn products or acid reducers.

Do not use with other antacid products or acid reducers.

Do not use with other heartburn products or acid reducers.

HOW TO TAKE:

Take 1 tablet before breakfast and one additional tablet before dinner.

CAUTIONS:

Do not use with other antacid products or acid reducers.

Do not use with other heartburn products or acid reducers.

Do not use with other antacid products or acid reducers.

Do not use with other heartburn products or acid reducers.


5-11-00
MINIMUM OF

REV. NO.		FC-D-706	McNeil <small>Pharmaceuticals, Inc.</small>
FORM		ARS	
REV. NO.		1.1	DISPENSIT
DATE		4/21/00	5-3/4 X 4-1/4 X 4-5/16
REVISED	BY	DATE	APPROVED
1	1		

Labeling Format Information:
Points: Univers condensed, bold and bold italic.
Drug Facts: 10 pt Leading: 8 pt
Header: 9 pt Bullets: 5 pt
Subheader: 6.5 pt Barlines: 2.5 pt
Body Text: 6.5 pt Hairlines: .5 pt
Drug Facts (continued): NA
Horizontal Scale: 94% Average Kerning: -4

DESCRIPTION DISP PEPCID COMP
FILE # 3003222
DATE May 31, 2001
M204, M209, M289
PS MADE # M300, M245, LIVE ART
REGISTRATION OF PVB, CO.
YELLOW CYAN SLA
THIS DESIGN BLOCK MUST BE SAVED W

DR LOT & EXP DATE ONLINE PRINTING



Drug Facts

Active Ingredients (by each chemically stable): Famotidine HCl, Ranitidine HCl, and Cimetidine HCl.


Caution: Do not take if you are allergic to famotidine or other H2 blockers.

Warnings: Do not use if you are allergic to famotidine or other H2 blockers. Do not use if you have trouble swallowing. Do not use if you are currently taking a prescription drug. Ask your doctor if you are currently taking a prescription drug. Ask your doctor if you are currently taking a prescription drug. Ask your doctor if you are currently taking a prescription drug.

Directions: Adults and children 12 years and over: Take 1 tablet 2 times a day with meals and at bedtime. Children 6 to 11 years: Take 1 tablet 2 times a day with meals and at bedtime. Children 2 to 5 years: Take 1/2 tablet 2 times a day with meals and at bedtime. Children under 2 years: Ask a doctor.

Other important information: Keep this medicine and warnings before use. Do not use if you are allergic to famotidine or other H2 blockers. Do not use if you have trouble swallowing. Do not use if you are currently taking a prescription drug. Ask your doctor if you are currently taking a prescription drug. Ask your doctor if you are currently taking a prescription drug.

Distributed by
Johnson & Johnson • MERCK
CONSUMER PHARMACEUTICALS CO.



Distributed by:
 Johnson & Johnson • MERCK
 CONSUMER PHARMACEUTICALS CO.
 FORT WASHINGTON, PA 19024 USA



WARNING - Do not use this product for more than 14 days.

If you are pregnant or trying to get pregnant, ask your doctor for advice. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

- Adults and children 12 years and over: - inhale 1 puff 3 times a day, 4 times a day, or as directed.
- Children 6 years of age and older: - inhale 1 puff 3 times a day, 4 times a day, or as directed.
- Children under 6 years of age: - use only with a spacer.

Other Information:

- Read the directions and warnings before use.
- Use only when you have symptoms of asthma.
- Do not use for more than 14 consecutive days.
- If you are using a spacer, see the instructions in the spacer's directions for use.
- For more information, call 1-800-458-5232, visit www.jnj.com, or visit www.merck.com.

Contains 100 puffs (2.5 mg) of albuterol sulfate.

Expiration or discontinuation: see instructions.

Leaving Format Information:

Form: Unit's condensed, bold and bold italic.

Drug Facts:

Font Size:	10 pt
Background:	5 pt
Subheader:	8.5 pt
Body Text:	8.5 pt
Header:	10 pt
Header:	8 pt
Header:	8 pt
Header:	5 pt
Header:	5 pt
Header:	5 pt
Header:	5 pt

Average Kerning: -4

Horizontal Scale: 94%

RESCRIPTION DISPENSER COMP BERRY CHEW TABS 3001 PROF

FILE #	300322	VERSION	E-1
DATE	May 31, 2001	ONE UNITS	FC-D-706-03
PS MADE #	WEG, N224, M29	KEY CONTACT	Tony Kozack
DESIGN #			
PROGRAM #			
SERIAL #			
REASON FOR CHANGE FROM COMMENTS			
YELLOW	CYAN	BLACK	OK BLUE
RED	RED	RED	RED

COMPARE BY: [Redacted]

FORMULA NUMBER: [Redacted]

PREPARED BY: [Redacted]

DATE: [Redacted]

EXPIRES: [Redacted]

REVISION: [Redacted]

REVISION DATE: [Redacted]

REVISION NUMBER: [Redacted]

REVISION DESCRIPTION: [Redacted]

REVISION ANALYST: [Redacted]

REVISION APPROVAL: [Redacted]



REVISION DATE: [Redacted]

REVISION BY: [Redacted]

Labeling Format Information:
 Fonts: Uppercase condensed, bold and bold
 Drugs Facts: 10 pt
 Header: 8.5 pt
 Subheader: 6.5 pt
 Body Text: 6 pt
 Footnote: 6 pt
 Horizontal Scale: 90%
 Average K:

JOB DESCRIPTION: CTN PEPCID COMP BERRY CHEW TAB 35			
FILE #	3002222	VERSION	C-1
DATE	May 31, 2001	DWG UNIT	FC-D-707.03
PS MAKE #	M299/LIVE.ART	KEY CONTACT	Tom Hoenes
DESIGN ILLUSTR.	PROGRAM	DESIGN ILLUSTR.	ver. 8.0
THIS DESIGN BLOCK MUST BE SAVED WITH FILE AND PRINTED ON THICK PAPER			
YELLOW	BLACK	CMAN	RED
DK BLUE	BLACK	CMAN	RED
DRAWING ANALYST			
DATE			
JOB #			
JOB NAME			
JOB NUMBER			
JOB UNIT			
JOB UNIT			
JOB UNIT			

<p>Peppid Complete</p> <p>Acid Reductant - Antacid</p> <p>Reduces Excess Stomach Acidity</p> <p>Relieves Heartburn Due to Acid Indigestion</p>	
<p>35 Berry Flavored Chewable Tablets</p>	
<p>Take-Along Packs</p>	
<p>Works with Just One Tablet!</p>	
<p>Contains Caffeine-Free Berry Flavored Antacid Tablets</p>	
<p>Relieves Heartburn Due to Acid Indigestion</p>	
<p>35 Chewable Tablets</p>	



Peppid Complete
 Acid Reductant - Antacid
 Reduces Excess Stomach Acidity
 Relieves Heartburn Due to Acid Indigestion

Take-Along Packs

Works with Just One Tablet!

Contains Caffeine-Free Berry Flavored Antacid Tablets
 Relieves Heartburn Due to Acid Indigestion

35 Chewable Tablets

Peppid Complete
 Acid Reductant - Antacid
 Reduces Excess Stomach Acidity
 Relieves Heartburn Due to Acid Indigestion

Take-Along Packs

Works with Just One Tablet!

Contains Caffeine-Free Berry Flavored Antacid Tablets
 Relieves Heartburn Due to Acid Indigestion

35 Chewable Tablets

Peppid Complete
 Acid Reductant - Antacid
 Reduces Excess Stomach Acidity
 Relieves Heartburn Due to Acid Indigestion

Take-Along Packs

Works with Just One Tablet!

Contains Caffeine-Free Berry Flavored Antacid Tablets
 Relieves Heartburn Due to Acid Indigestion

35 Chewable Tablets

Peppid Complete
 Acid Reductant - Antacid
 Reduces Excess Stomach Acidity
 Relieves Heartburn Due to Acid Indigestion

Take-Along Packs

Works with Just One Tablet!

Contains Caffeine-Free Berry Flavored Antacid Tablets
 Relieves Heartburn Due to Acid Indigestion

35 Chewable Tablets

XXXXXXXX


Complete

216

Do not use if the surface is scratched or worn.

	Very Toxic Caution: Avoid contact with skin, eyes, nose, mouth, and clothing. Wash thoroughly with soap and water. If swallowed, do not induce vomiting. Seek medical attention immediately.
	Flammable (Liquid) Caution: Keep away from heat, open flames, and other sources of ignition. Do not use in enclosed spaces.
	Corrosive Caution: Causes severe skin burns and eye damage. Avoid contact with skin and eyes. If contact occurs, wash immediately with plenty of water.
	Hazardous to the Environment Caution: Do not discharge into the environment. Dispose of in accordance with local, state, and federal regulations.

Distributed by
Johansen + Johnson • MERCK
CONSUMER PHARMACEUTICALS CO.
FORT WASHINGTON, PA 19034 USA



Using Format Information:

Font: Univers condensed, bold and bold italic

g: Factor: 10 pt
 Leader: 6.5 pt
 descender: 5 pt
 x-height: 2.5 pt
 ascender: 6.5 pt
 y: Factor: 5 pt
 g: Factor: 5 pt
 h: Factor: 5 pt
 k: Factor: 5 pt

Average Kerning: -20


- NOTES:
- 1) DRAWING SHOWN PRINT SIZE UP
 - 2) ALL DIMENSIONS IN INCHES
 - 3) TOLERANCES UNLESS SPECIFIED
 - 4) FRACTIONS 1/8", 3/16", 1/4", 3/8", 1/2", 5/8", 3/4", 7/8"
 - 5) BASED ON PRT. DIM. UNLESS OTHERWISE SPECIFIED (6-12-90)
 - 6) ALL LINE CORN. & TERT. TO BE A MINIMUM OF 1/16"
 - 7) FROM ALL SIDES & CUTS
 - 8) NO PRINT AREA RESERVED FOR LOT & EXP. DATE AREA
 - 9) NO VARNISH AREA

REV.	DATE	DESCRIPTION	BY	CHKD.
01	02/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
02	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
03	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
04	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
05	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
06	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
07	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
08	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
09	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
10	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
11	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
12	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
13	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
14	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
15	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
16	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
17	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
18	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
19	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
20	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
21	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
22	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
23	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
24	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
25	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
26	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
27	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
28	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
29	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
30	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
31	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
32	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
33	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
34	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
35	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
36	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
37	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
38	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
39	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
40	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
41	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
42	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
43	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
44	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
45	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
46	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
47	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
48	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
49	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
50	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
51	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
52	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
53	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
54	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
55	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
56	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
57	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
58	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
59	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
60	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
61	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
62	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
63	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
64	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
65	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
66	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
67	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
68	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
69	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
70	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
71	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
72	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
73	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
74	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
75	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
76	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
77	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
78	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
79	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
80	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
81	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
82	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
83	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
84	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
85	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
86	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
87	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
88	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
89	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
90	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
91	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
92	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
93	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
94	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
95	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
96	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
97	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
98	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
99	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL
100	03/27/00	REVISED TO ADD DIMENSIONS	ARL	JL

MONTELL
FC-D-707
ARL
JL
DISPENSIT
4-3/4 X 3-3/4 X 4-5/16

NOTES:
 1) ALL DIMENSIONS IN INCHES.
 2) TOLERANCES UNLESS SPECIFIED.
 3) FRACTIONS #1/64
 4) NO COLOR, NO COPY, NO BLEED

REV. NO.	DATE	REVISIONS
1	10/10/97	REVISED MEASUREMENTS
2		
3		
4		
5		
6		
7		
8		
9		
10		
11		
12		
13		
14		
15		
16		
17		
18		
19		
20		
21		
22		
23		
24		
25		
26		
27		
28		
29		
30		
31		
32		
33		
34		
35		
36		
37		
38		
39		
40		
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		



Pepcid Complete
Add Relief. A Complete.

DO NOT USE IF THE INDIVIDUAL WHOSE'S OWN IT IS.

WHAT YOU SHOULD KNOW ABOUT:

PACKAGE INSERT

A non prescription stomach medicine.

Pepcid Complete contains an acid reducer (ranitidine) with antacids. Antacids neutralize and temporarily relieve (but do not prevent) heartburn. Heartburn usually occurs in the upper stomach. Antacids do not prevent heartburn from returning. In Pepcid Complete, the acid reducer (ranitidine) is added to the antacids. This combination helps prevent heartburn from returning. In millions of people safety and effectiveness.

• Just 1 tablet relieves heartburn due to acid indigestion.

How to use PEPCID COMPLETE

Use PEPCID COMPLETE to relieve heartburn due to acid indigestion. Swallow 1 tablet with water. If you are taking other heartburn medicine, ask your doctor. Do not take more than 12 tablets each day.

TO RELIEVE SYMPTOMS
 Chew 1 tablet completely before swallowing.

Express acid in burning problem

It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the stomach may move up into the wrong places (the esophagus), causing burning pain and discomfort that heartburn with every swallow.

• Heartburn is caused by acid in the wrong places.


Use Pepcid Complete

- Do not lie flat for over one hour after eating.
- Do not drink alcohol, caffeine, spicy, acidic, or fatty foods.
- Do not eat chocolate, citrus fruits, or other foods that may worsen heartburn.
- Do not eat acidic foods, such as tomatoes, citrus fruits, and other foods that may worsen heartburn.
- Do not eat spicy foods, such as peppers, onions, and other foods that may worsen heartburn.
- Do not eat fatty foods, such as fried foods, pizza, and other foods that may worsen heartburn.
- Do not eat carbonated drinks, such as soda and beer.
- Do not eat minty foods, such as mints and gum.
- Do not eat salty foods, such as pickles and olives.
- Do not eat hard-to-digest foods, such as nuts and seeds.
- Do not eat high-fiber foods, such as bran cereal and high-fiber bread.

PEPCID COMPLETE
 Ranitidine HCl and Antacids
 Each 1 tablet completely relieves heartburn due to acid indigestion. Contains 12 tablets each in a blister pack.

Proven effective in clinical studies

In clinical studies, PEPCID COMPLETE was significantly better than ranitidine for heartburn relief.



Percent of heartburn episodes adequately relieved within 30 minutes from the time of dosing

Time	PEPCID COMPLETE	Ranitidine
0-15 min	63%	52%
16-30 min	67%	56%
31-45 min	68%	57%
46-60 min	69%	58%
61-75 min	69%	58%
76-90 min	69%	58%
91-105 min	69%	58%
106-120 min	69%	58%

PEPCID COMPLETE was significantly better than ranitidine for heartburn relief.

Warnings

Heartburn and acid indigestion are common, but you should stop taking PEPCID COMPLETE and see your doctor promptly if:

- You have trouble swallowing or stomach pain continues.
- You need to take this product for more than 14 days.
- You may have a serious condition that may need different treatment.

Always alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use with other famotidine products or acid reducers. Antacids may interact with certain prescription drugs. If you are presently taking a prescription drug, ask a doctor or pharmacist before use. This product should not be given to children under 13 years old, unless directed by a doctor.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Questions or comments?

Please call 1-800-755-4066 for more information and a guide on how to help control heartburn.

Developed for:
 Janssen Pharmaceutica, Inc.
 Division of Janssen-Cilag, Inc.
 One Laurent Street
 New Brunswick, NJ 07102 USA

DESCRIPTION: 8172 PEPCID COMP CHEW TAB 15 COUNT

VERSION: C-2

DATE: December 6, 2000

REV CONTRACT: 1500

COMPARED BY: [REDACTED]

PROGRAM TITLE, INC. NO.: [REDACTED]

REVISIONS: 0100

REVISIONS: 0200

REVISIONS: 0300

REVISIONS: 0400

REVISIONS: 0500

REVISIONS: 0600

REVISIONS: 0700

REVISIONS: 0800

REVISIONS: 0900

REVISIONS: 1000

REVISIONS: 1100

REVISIONS: 1200

REVISIONS: 1300

REVISIONS: 1400

REVISIONS: 1500

REVISIONS: 1600

REVISIONS: 1700

REVISIONS: 1800

REVISIONS: 1900

REVISIONS: 2000

REVISIONS: 2100

REVISIONS: 2200

REVISIONS: 2300

REVISIONS: 2400

REVISIONS: 2500

REVISIONS: 2600

REVISIONS: 2700

REVISIONS: 2800

REVISIONS: 2900

REVISIONS: 3000

Labeling Format Information: and bold face.

Drug Facts: 10 pt
Warnings: 8 pt
Directions: 8 pt
Other information: 8 pt
Contains: 8 pt
See also: 8 pt

Horizontal Scale: 84%

Average Kerning: 4

Janssen Pharmaceutica, Inc.
 One Laurent Street
 New Brunswick, NJ 07102 USA

PEPCID COMPLETE
 CHEWING TABLETS

Dist. by
J.S.J. MERCK CPC
Pt. Wash., PA 19034
USA

9000000

N EXP. XX/XX
LOT XXXXXX

Pepcid[®]
Complete
(Famotidine Salt Formulation)

1 Berry Flavored Chewable Tablet

Famotidine 30 mg
Calcium carbonate 600 mg
Magnesium hydroxide 165 mg

NOTES:

- 1) ALL DIMENSIONS IN INCHES.
- 2) PRINT REPEAT TOLERANCE NOT TO EXCEED ±1/32 OVER 12 IMPRESSIONS (35,508 4.031)
- 3) NO COPY AREA
- 4) LDT & EXP. DATE AREA

REV. NO.	DATE	REVISIONS	FIG. NO.	UNWIND-131	COMP. PRODUCTS CO. 2000 MILLERS RD. FORT WASHINGTON, PA. 19044
			DATE	3/31/00	CR POUCH MATERIAL UNWIND 2.959 L X 3.828 W

Pepcid[®] Complete

WITH **DOUBLE ACTION**

Acid Reducer + Antacid

Works with Just One Tablet!

Reduces AND Neutralizes Acid

Contains Calcium

Relieves Heartburn Due To Acid Indigestion

1 Chewable Tablet

Do not use if pouch is torn or tampered with.

Drug Facts

Active Ingredients (in each chewable tablet): Famotidine 10 mg, Calcium carbonate 800 mg, Magnesium hydroxide 165 mg.

Uses: Relieves heartburn associated with acid indigestion and sour stomach.

Warnings: Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.

Drug Facts (continued)

Do not use if you have trouble swallowing or with other famotidine products or acid reducers.

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if stomach pain continues or you need to take this product for more than 14 days.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

While folded on line, tear open at slit.

Drug Facts (continued)

Directions: Adults and children 12 years and over: Take one (1) white tablet whole slowly completely to relieve symptoms, chew tablet before swallowing. Do not use more than 2 chewable tablets in 24 hours. Children under 12 years: ask a doctor.

Other Information: Store at 25-30°C (77-86°F).

Contains: Famotidine, calcium carbonate, DCC and FF, croscarmellose sodium, hydroxypropyl methylcellulose, lactose, magnesium stearate, polyethylene glycol, and polyethylene glycol 400.

Questions or comments? Call toll-free 1-888-AS-8888.

150%OFSIZE

Pepcid[®] Complete

WITH **DOUBLE ACTION**

Acid Reducer + Antacid

Works with Just One Tablet!

Reduces AND Neutralizes Acid

Contains Calcium

Relieves Heartburn Due To Acid Indigestion

1 Chewable Tablet

Do not use if pouch is torn or tampered with.

Drug Facts

Active Ingredients (in each chewable tablet): Famotidine 10 mg, Calcium carbonate 800 mg, Magnesium hydroxide 165 mg.

Uses: Relieves heartburn associated with acid indigestion and sour stomach.

Warnings: Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.

Drug Facts (continued)

Do not use if you have trouble swallowing or with other famotidine products or acid reducers.

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if stomach pain continues or you need to take this product for more than 14 days.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

While folded on line, tear open at slit.

Drug Facts (continued)

Directions: Adults and children 12 years and over: Take one (1) white tablet whole slowly completely to relieve symptoms, chew tablet before swallowing. Do not use more than 2 chewable tablets in 24 hours. Children under 12 years: ask a doctor.

Other Information: Store at 25-30°C (77-86°F).

Contains: Famotidine, calcium carbonate, DCC and FF, croscarmellose sodium, hydroxypropyl methylcellulose, lactose, magnesium stearate, polyethylene glycol, and polyethylene glycol 400.

Questions or comments? Call toll-free 1-888-AS-8888.

Labeling Format Information:

Font: Univers condensed, bold and bold italic.	
Drug Facts:	8.5 pt
Header:	7 pt
Subheader:	6 pt
Body Text:	6 pt
Drug Facts (continued):	6 pt
Leadline:	6 pt
Subline:	5 pt
Bottomline:	4.5 pt
Headline:	5 pt

Horizontal Scale: 84-94% Average Naming: -4

DESCRIPTION	POUCHPEPCIDCOMPBERRYCHEWTABPROF
FILE #	30032ME VERSION
DATE	May01.200UNWIND-131.06 VINYL
PS IMAGE #	LIVEART KEY CONTACT: Tony Nucera
COMPARED BY:	
DESIGN PROGRAM:	ILLUSIO ver.
MAXIMUMS OF FIVE COLORS PER COMPONENT	
YELLOW BLACK RED	
THIS DESIGN BLOCK MUST BE SAVED WITH FILE AND PRINTED ON TERMINAL.	

PROVIDER	
GRAPHIC ANALYST	
UPC #	
PART NUMBER	
PCRNPA NUMBER	
FCC#	
PACKAGING LOCATION	
COLOR STAMPED	
TOLERANCE DATE	

McNeil Johnson & Johnson • MERCK	
ARTWORK APPROVAL INITIAL BLOCK	
INITIALS	DATE
REGULATORY	
MEDICAL	

NOTES:

- ALL DIMENSIONS IN DASHES
- TOLERANCES UNLESS SPECIFIED
- FRACTIONS 3/16"
- SHOWING DIMEN. POINTS UP
- NO POINT AREA
- PART NUMBER AREA MUST BE OPEN TO TOP FOR PART NUMBER UP OR FOR DIMEN. MAY MARK UP AREA
- NO POINT, NO COLOR & NO VARIATION AREA
- "LOT" AND "EXP" POINT AREA

REV. NO.	DATE	REVISIONS

DWG. NO.	LB-PS-848
REV. NO.	30
DATE	2/26/99

Johnson & Johnson MERCK
 Consumer Pharmaceuticals Co.
 P.O. Box 1187, Fort Washington, PA 19044

TITLE: BOOKLET LABEL FOR FACT 200CC BOTTLE

RESERVED FOR COUPON TEXT

EXCESS ACID-BURNING PROBLEM
 It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the stomach may move up into the wrong place (the esophagus), causing burning pain and discomfort that interferes with everyday activities.

Reflux—Caused by acid in the stomach
 Burning pain/discomfort in esophagus

A valve-like muscle called the lower esophageal sphincter (LES) is relaxed in an open position. Acid soaks up from stomach.

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat too much.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

50 Berry Flavored Chewable Tablets

Pepcid Complete
 Acid Reducer + Antacid
 350mg Famotidine
 100mg Ranitidine HCl
 50 Berry Flavored Chewable Tablets

Works with Just One Tablet

50 Berry Flavored Chewable Tablets

Drug Facts (continued)

Warnings:
 Always check Do not use if you are allergic to famotidine or other acid reducers.

Do not use:
 • If you have possible sensitivity with other famotidine products or acid reducers.

Take a doctor or pharmacist beforehand if you are presently taking a prescription drug. Famotidine may interact with certain prescription drugs.

How can you tell if it's done?
 • Check for heartburn relief. You need to take this product for more than 14 days.

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:
 • Adults and children 12 years and over:
 - Take one (1) tablet twice daily with water, as needed.
 - To relieve symptoms, chew 1 tablet before eating.
 - Do not use more than 2 chewable tablets in 24 hours.
 - At least 2 hours after last meal.

Other information:
 • Read the directions and warnings before use. • Keep it in its bottle. It contains important information.
 • Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F).

Contains Potentially Hazardous Ingredients: famotidine, ranitidine, HCl, sodium phosphate, sodium hydroxide, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose K100, hydroxypropyl methylcellulose K100, hydroxypropyl methylcellulose K100, hydroxypropyl methylcellulose K100.

Questions or comments? Call toll free 1-800-769-4000.

NON-PRESCRIPTION STOMACH MEDICINE

PEPCID COMPLETE combines an acid reducer (famotidine) with antacids (calcium carbonate) and magnesium hydroxide to relieve heartburn in two different ways: Acid reducers decrease the production of new stomach acid, blocking incoming acid that is already in the stomach. The antacids (calcium carbonate and magnesium hydroxide) help to neutralize the acid already in the stomach.

Shown actual size

• Just 1 tablet relieves heartburn due to acid indigestion.

RESERVED FOR COUPON TEXT

PROVEN EFFECTIVE IN CLINICAL STUDIES
 In clinical studies, PEPCID COMPLETE was significantly better than placebo tablets (tablets without medicine) in relieving heartburn.

Percent of heartburn episodes adequately relieved within 30 minutes

Treatment	Percent of episodes adequately relieved within 30 minutes
PEPCID COMPLETE	75
Placebo	45
Pepcid AC	55

Percent of heartburn episodes adequately relieved at 7 hours from the time of dosing

Treatment	Percent of episodes adequately relieved at 7 hours
PEPCID COMPLETE	70
Placebo	45
Pepcid AC	55

QUESTIONS OR COMMENTS?
 Please call 1-800-769-4000 for more information or call a pharmacist for help in buying generic medicines.

Distributed by: Johnson & Johnson MERCK
 CONSUMER PHARMACEUTICALS CO.
 FORT WASHINGTON, PA 19044
 © Registered trademark of Merck & Co., Inc.

Labeling Format Information:
 Fonts: Univers condensed, bold and bold italic.
 Drug Facts: 9 pt Leading: 6.5 pt
 Header: 8 pt Bullets: 6 pt
 Subheader: 8 pt Barlines: 2.5 pt
 Body Text: 8 pt Hairlines: 5 pt
 Drug Facts (continued): 8 pt
 Horizontal Scale: 94% Average Kerning: -4

JOB DESCRIPTION LBLPEPCIDCOMPBERRYCHEW7AB50

FILE # 3003222B-1 VERSION

DATE May 31 2001 LB-PS-848.01E VINYL

PS IMAGE # M300LIVEART KEY CONTACT Tony Hanco

DESIGNED BY: [Name]

COMPARED BY: [Name]

DESIGN PROGRAM ILLUSTRATOR

MAKING UP OF FIVE COLORES PER COMPONENT

YELLOW BLACK RED GREEN BLUE

THIS DESIGN IS TO BE USED AS A GUIDE WITH FILE AND SUBJECT TO TOLERANCE

PRINTER	ARTWORK APPROVAL
GRAPHIC ANALYST	PITIAL BLOCK
DATE #	INITIALS DATE
PART NUMBER	REGULATORY
FORMA NUMBER	MEDICAL
PECA	
PACKAGING LOCATION	
COLOR STANDARD	
VALIDITY DATE	

McNEIL Johnson & Johnson MERCK

**This is a representation of an electronic record that was signed electronically and
this page is the manifestation of the electronic signature.**

/s/

Linda Katz
11/6/01 03:28:29 PM