

**CENTER FOR DRUG EVALUATION AND RESEARCH**

*APPLICATION NUMBER:*

**50-784 / S-004, S-006**

**CLINICAL PHARMACOLOGY/  
BIOPHARMACEUTICS REVIEW(S)**

---

NDA#	50-784 (SE1-004)
DRUG NAME	Azithromycin (Zithromax®)
FORMULATION	500 mg film-coated tablets
SUBMISSION DATE	March 17, 2003
SUBMISSION TYPE	NDA Efficacy Supplement
SPONSOR	Pfizer, Inc., New York, NY 10017
MEDICAL DIVISION	Anti-Infective Drug Products
REVIEWER	Charles R. Bonapace, Pharm.D.
TEAM LEADER	Venkat R. Jarugula, Ph.D.

---

## CLINICAL PHARMACOLOGY/BIOPHARMACEUTICS REVIEW

### BACKGROUND:

NDA 50-784 was approved on May 24, 2002 for azithromycin 500 mg tablets once-daily for 3 days to treat acute bacterial exacerbations of chronic obstructive pulmonary disease (AECB) in adults due to susceptible strains of *Haemophilus influenzae*, *Moraxella catarrhalis*, and *Streptococcus pneumoniae*. The sponsor submitted the current efficacy supplement (SE1-004) for a new indication of acute bacterial sinusitis for Zithromax® (azithromycin) 500 mg tablets. This efficacy supplement provides clinical safety and efficacy data for the addition of acute bacterial sinusitis due to *Haemophilus influenzae*, *Moraxella catarrhalis*, and *Streptococcus pneumoniae* using the currently approved dosing regimen (azithromycin 500 mg once-daily for 3 days) in adults.

The sponsor performed two clinical efficacy studies (Study 066-1036 and Study 066-1057) to support the safety and efficacy of the azithromycin 3-day dosing regimen in adults. Both studies used the currently marketed formulation of azithromycin 500 mg tablets. No Clinical Pharmacology studies were submitted to support the efficacy supplement.

**Study 066-1036 (n=941):** This study was a Phase 3, randomized, multicenter, double-blind, comparative trial to evaluate the clinical efficacy of azithromycin vs. comparator in patients 18 years of age and older with clinically documented acute bacterial infection of the maxillary sinuses. Patients were randomized to receive azithromycin 500 mg once-daily for 3 days, azithromycin 500 mg once-daily for 6 days, or amoxicillin 500 mg/clavulanate 125 mg TID for 10 days.

**Study 066-1057 (n=539):** This study was a Phase 3, multicenter, open-label, randomized study to evaluate the bacteriologic efficacy of azithromycin in patients 18 years of age and older with clinically and radiologically documented acute bacterial infection of the maxillary sinuses. Patients were randomized to receive azithromycin 500 mg once-daily for 3 days or azithromycin 500 mg once-daily for 6 days.

The sponsor is also filing labeling supplements for NDA 50-710 (azithromycin for oral suspension) and NDA 50-711 (azithromycin 250 mg tablets) since they are included in the same label as the 500 mg tablets. Azithromycin for oral suspension is approved for patients  $\geq 6$  months of age for acute otitis media (10 mg/kg on day 1, then 5 mg/kg/day for 4 days OR 10 mg/kg/day for 3 days OR 30 mg/kg single dose), community acquired pneumonia (10 mg/kg on day 1, then 5 mg/kg/day for 4 days), and patients  $\geq 2$  yrs of age for pharyngitis/tonsillitis (12 mg/kg/day for 5 days).

The sponsor's proposed labeling includes the use of a 3-day regimen of azithromycin for oral suspension in the treatment of pediatric subjects  $\geq 6$  months of age with acute bacterial sinusitis at 10 mg/kg/day (30 mg/kg total). The pediatric claim is based on the safety and efficacy that has been established in the present NDA supplement in adults with acute bacterial sinusitis and on the fact that the 3-day pediatric dosing has been shown to be safe and effective in treating acute otitis media. The microbiology of acute bacterial sinusitis in children is similar to that seen in adults, with the primary pathogens being *H. influenzae*, *M. catarrhalis*, and *S. pneumoniae*. The same pathogens are the most frequent cause of bacterial acute otitis media in children.

The sponsor did not submit pharmacokinetic data comparing azithromycin 500 mg/day  $\times$  3 days in adults to 10 mg/kg/day  $\times$  3 days in pediatrics. However, the sponsor previously performed studies assessing the pharmacokinetics of azithromycin in healthy adults following the administration of 500 mg/day  $\times$  3 days (proposed dosage for the treatment of acute bacterial sinusitis) and infected pediatrics following the administration of 10 mg/kg/day  $\times$  3 days (proposed dosage for the treatment of acute bacterial sinusitis), 12 mg/kg/day  $\times$  5 days, or 20 mg/kg/day  $\times$  3 days. In the pediatric studies, the pharmacokinetics were only assessed after the last dose to minimize blood loss. The reviewer compared the AUC<sub>0-24</sub> obtained on day 1 and day 3 in adults to the AUC<sub>0-24</sub> obtained on day 3 or day 5 in pediatrics. A comparison of the studies and the mean (CV %) AUC<sub>0-24</sub> values are shown in the table below.

Parameter	066-087 (500 mg/day $\times$ 3 days)	AZM-F-93-004 (500 mg/day $\times$ 3 days)	066-220 (10 mg/kg/day $\times$ 3 days)	066-683 (12 mg/kg/day $\times$ 5 days)	066-095 (20 mg/kg/day $\times$ 3 days)
No. of subjects	12	12	16	17	12
Age	Adults	>18 yrs	6 months - 10 yrs	4 - 13 yrs	3 - 14 yrs
Population	Healthy	Healthy	Infected	Infected	Infected
Status	Fasted	Fasted	Fasted	Low-fat meal	Low-fat meal
Day 1 AUC <sub>0-24</sub> ( $\mu\text{g}\cdot\text{hr}/\text{mL}$ )	2.58 (33%)	1.85 (32%)	ND	ND	ND
Day 3 or 5 AUC <sub>0-24</sub> ( $\mu\text{g}\cdot\text{hr}/\text{mL}$ )	3.99 (31%)	3.09 (32%)	2.35 (81%)	3.94 (48%)	7.92 (36%)
AUC <sub>0-24</sub> adjusted to 10 mg/kg	---	---	2.35	3.28	3.96

ND - plasma concentrations were not obtained following the first dose

Following the administration of azithromycin 500 mg/day  $\times$  3 days to healthy adults, the mean AUC<sub>0-24</sub> on day 3 ranged from 3.09 to 3.99  $\mu\text{g}\cdot\text{hr}/\text{mL}$ . In comparison, the mean AUC<sub>0-24</sub> values (adjusted for a 10 mg/kg dose) on day 3 or day 5 ranged from 2.35 to 3.96  $\mu\text{g}\cdot\text{hr}/\text{mL}$  and were similar to the values obtained from adults. Thus, the AUC<sub>0-24</sub> resulting from the administration of 10 mg/kg/day for 3 days in pediatric patients is likely to be similar to the AUC<sub>0-24</sub> resulting from the administration of 500 mg/day  $\times$  3 days in adults.


The AUC<sub>0-24</sub> obtained after the last dose in pediatric subjects receiving 10 mg/kg/day (Study 066-220) was 24% to 41% lower than adults receiving 500 mg/day. However, there was a large degree of inter-subject variability in the studies reflected by the large coefficient of variation. The AUC<sub>0-24</sub> obtained after the last dose in pediatric subjects receiving 20 mg/kg/day for 3 days (Study 066-095) was 3.37-fold the values obtained from pediatric subjects receiving 10 mg/kg/day for 3 days. The current label states that food increases the C<sub>max</sub> of azithromycin for oral suspension by 56% but does not effect the AUC.

In Study 066-220, three subjects were missing one or more plasma concentrations that may have resulted in an inaccurate estimation of the AUC<sub>0-24</sub>. The mean (CV%) AUC<sub>0-24</sub> after the last dose of azithromycin

excluding these three subjects was 2.69 (72%). Thus, the  $AUC_{0-24}$  values based on this study may be only 13% to 33% lower in pediatric subjects receiving 10 mg/kg/day compared to adults receiving 500 mg/day.

One subject enrolled in Study 066-220 was between 6 months and 2 yrs of age, 7 subjects were between 2-5 yrs of age, and 8 subjects were between 5-10 yrs of age. There appeared to be differences in the pharmacokinetics of azithromycin between subjects based on their age. The individual and mean azithromycin concentration-time profile for each age category is shown in Figure 1.

**Figure 1. Individual (left) and mean (right) concentration-time profiles of azithromycin 10 mg/kg × 3 days by age group (from Study 066-220)**




NOTE: solid red lines, 5-10 yrs of age; broken blue lines, 2-4 yrs of age; broken green line, <2 yrs of age

Although the plasma concentrations of azithromycin were less in the <2 yrs of age group, the data are based on one subject. Because the impact of age on the pharmacokinetics of azithromycin has not been previously assessed, the sponsor was requested to perform a pharmacokinetic study addressing the impact of age on the pharmacokinetics of azithromycin as a Phase 4 commitment to NDA 50-710 (S-009 and S-010) approved December 14, 2001. The sponsor submitted the results from study A066-1049 to satisfy the Phase 4 commitment. The results from study A0661049 were reviewed by Dr. Charles R. Bonapace and are reported in the review dated January 7, 2004.

In study A066-1049, 32 pediatric subjects with infections were enrolled into four groups based on age: 6 months to 24 months (group 1), >24 months to <6 yrs (group 2), 6 yrs to <12 yrs (group 3), and 12 yrs to 16 yrs (group 4). All subjects received a single 10 mg/kg dose of intravenous azithromycin (maximum 500 mg) infused over 60 min. The mean serum azithromycin concentration-time profiles were similar between groups (see Figure 2).

Figure 2. Mean serum azithromycin concentration-time profiles following administration of a single 10 mg/kg (maximum 500 mg) intravenous dose



**RECOMMENDATIONS:**

This submission was reviewed by the Office of Clinical Pharmacology and Biopharmaceutics, Division of Pharmaceutical Evaluation III and found to be acceptable from a clinical pharmacology point of view.

Please forward the attached labeling changes to the reviewing medical officer and the sponsor.

**/S/**

Charles R. Bonapace, Pharm.D.  
Office of Clinical Pharmacology/Biopharmaceutics  
Division of Pharmaceutical Evaluation III

**/S/**

RD/FT Initialed by Venkat R. Jarugula, Ph.D., \_\_\_\_\_  
Team Leader

cc:  
Division File: NDA 50-784  
HFD-520 (CSO/Milstein)  
HFD-520 (MO/Alexander, Moledina)  
HFD-880 (Division File, Lazor, Selen, Jarugula, Bonapace)  
CDR (Clin. Pharm./Biopharm.)

25

(Version March 17, 2003)

**ZITHROMAX<sup>®</sup>**  
**(azithromycin tablets)**  
**and**  
**(azithromycin for oral suspension)**

**DESCRIPTION**

ZITHROMAX<sup>®</sup> (azithromycin tablets and azithromycin for oral suspension) contain the active ingredient azithromycin, an azalide, a subclass of macrolide antibiotics, for oral administration.


Azithromycin has the chemical name (2*R*,3*S*,4*R*,5*R*,8*R*, 10*R*,11*R*,12*S*,13*S*,14*R*)-

13-[(2,6-dideoxy-3-*C*-methyl-3-*O*-methyl- $\alpha$ -*L*-ribo-hexopyranosyl)

oxy]-2-ethyl-3,4,10-trihydroxy-3,5,6,8,10,12,14-heptamethyl-11-[[[3,4,6-trideoxy-3-

(dimethylamino)- $\beta$ -*D*-xylo-hexopyranosyl]oxy]-1-oxa-6-azacyclopentadecan-15-one.

Azithromycin is derived from erythromycin; however, it differs chemically from erythromycin in that a methyl-substituted nitrogen atom is incorporated into the lactone ring. Its molecular formula is C<sub>38</sub>H<sub>72</sub>N<sub>2</sub>O<sub>12</sub>, and its molecular weight is 749.00. Azithromycin has the following structural formula:



Azithromycin, as the dihydrate, is a white crystalline powder with a molecular formula of C<sub>38</sub>H<sub>72</sub>N<sub>2</sub>O<sub>12</sub>•2H<sub>2</sub>O and a molecular weight of 785.0.

ZITHROMAX<sup>®</sup> is supplied for oral administration as film-coated, modified capsular shaped tablets containing azithromycin dihydrate equivalent to either 250 mg or 500 mg azithromycin and the following inactive ingredients: dibasic calcium phosphate anhydrous, pregelatinized

starch, sodium croscarmellose, magnesium stearate, sodium lauryl sulfate, hydroxypropyl methylcellulose, lactose, titanium dioxide, triacetin and D&C Red #30 aluminum lake.

ZITHROMAX<sup>®</sup> for oral suspension is supplied in bottles containing azithromycin dihydrate powder equivalent to 300 mg, 600 mg, 900 mg, or 1200 mg azithromycin per bottle and the following inactive ingredients: sucrose; sodium phosphate, tribasic, anhydrous; hydroxypropyl cellulose; xanthan gum; FD&C Red #40; and spray dried artificial cherry, creme de vanilla and banana flavors. After constitution, each 5 mL of suspension contains 100 mg or 200 mg of azithromycin.

## CLINICAL PHARMACOLOGY

### Pharmacokinetics

Following oral administration of a single 500 mg dose (two 250 mg tablets) to 36 fasted healthy male volunteers, the mean (SD) pharmacokinetic parameters were  $AUC_{0-72} = 4.3 (1.2) \mu\text{g}\cdot\text{h}/\text{mL}$ ;  $C_{\text{max}} = 0.5 (0.2) \mu\text{g}/\text{mL}$ ;  $T_{\text{max}} = 2.2 (0.9)$  hours.

With a regimen of 500 mg (two 250 mg capsules\*) on day 1, followed by 250 mg daily (one 250 mg capsule) on days 2 through 5, the pharmacokinetic parameters of azithromycin in plasma in healthy young adults (18-40 years of age) are portrayed in the chart below.  $C_{\text{min}}$  and  $C_{\text{max}}$  remained essentially unchanged from day 2 through day 5 of therapy.

Pharmacokinetic Parameters (Mean)	Total n=12	
	Day 1	Day 5
$C_{\text{max}}$ ( $\mu\text{g}/\text{mL}$ )	0.41	0.24
$T_{\text{max}}$ (h)	2.5	3.2
$AUC_{0-24}$ ( $\mu\text{g}\cdot\text{h}/\text{mL}$ )	2.6	2.1
$C_{\text{min}}$ ( $\mu\text{g}/\text{mL}$ )	0.05	0.05
Urinary Excret. (% dose)	4.5	6.5

\*Azithromycin 250 mg tablets are bioequivalent to 250 mg capsules in the fasted state. Azithromycin 250 mg capsules are no longer commercially available.

In a two-way crossover study, 12 adult healthy volunteers (6 males, 6 females) received 1,500 mg of azithromycin administered in single daily doses over either 5 days (two 250 mg tablets on day 1, followed by one 250 mg tablet on days 2-5) or 3 days (500 mg per day for days 1-3). Due to limited serum samples on day 2 (3-day regimen) and days 2-4 (5-day regimen), the serum concentration-time profile of each subject was fit to a 3-compartment model and the  $AUC_{0-\infty}$  for the fitted concentration profile was comparable between the 5-day and 3-day regimens.

Pharmacokinetic Parameter [mean (SD)]	3-Day Regimen		5-Day Regimen	
	Day 1	Day 3	Day 1	Day 5
$C_{max}$ (serum, $\mu\text{g/mL}$ )	0.44 (0.22)	0.54 (0.25)	0.43 (0.20)	0.24 (0.06)
Serum $AUC_{0-\infty}$ ( $\mu\text{g}\cdot\text{hr/mL}$ )	17.4 (6.2)*		14.9 (3.1)*	
Serum $T_{1/2}$	71.8 hr		68.9 hr	

\*Total AUC for the entire 3-day and 5-day regimens

Median azithromycin exposure ( $AUC_{0-288}$ ) in mononuclear (MN) and polymorphonuclear (PMN) leukocytes following either the 5-day or 3-day regimen was more than a 1000-fold and 800-fold greater than in serum, respectively. Administration of the same total dose with either the 5-day or 3-day regimen may be expected to provide comparable concentrations of azithromycin within MN and PMN leukocytes.

Two azithromycin 250 mg tablets are bioequivalent to a single 500 mg tablet.

#### Absorption

The absolute bioavailability of azithromycin 250 mg capsules is 38%.

In a two-way crossover study in which 12 healthy subjects received a single 500 mg dose of azithromycin (two 250 mg tablets) with or without a high fat meal, food was shown to increase  $C_{max}$  by 23% but had no effect on AUC.

When azithromycin suspension was administered with food to 28 adult healthy male subjects,  $C_{max}$  increased by 56% and AUC was unchanged.

The AUC of azithromycin was unaffected by co-administration of an antacid containing aluminum and magnesium hydroxide with azithromycin capsules; however, the  $C_{max}$  was reduced by 24%. Administration of cimetidine (800 mg) two hours prior to azithromycin had no effect on azithromycin absorption.

#### Distribution

The serum protein binding of azithromycin is variable in the concentration range approximating human exposure, decreasing from 51% at 0.02  $\mu\text{g/mL}$  to 7% at 2  $\mu\text{g/mL}$ .

Following oral administration, azithromycin is widely distributed throughout the body with an apparent steady-state volume of distribution of 31.1 L/kg. Greater azithromycin concentrations in tissues than in plasma or serum were observed. High tissue concentrations should not be interpreted to be quantitatively related to clinical efficacy. The antimicrobial activity of azithromycin is pH related and appears to be reduced with decreasing pH. However, the extensive distribution of drug to tissues may be relevant to clinical activity.

Selected tissue (or fluid) concentration and tissue (or fluid) to plasma/serum concentration ratios are shown in the following table:

**AZITHROMYCIN CONCENTRATIONS FOLLOWING A 500 mg DOSE (TWO 250 mg CAPSULES) IN ADULTS<sup>1</sup>**

TISSUE OR FLUID	TIME AFTER DOSE (h)	TISSUE OR FLUID CONCENTRATION ( $\mu\text{g/g}$ or $\mu\text{g/mL}$ )	CORRESPONDING PLASMA OR SERUM LEVEL ( $\mu\text{g/mL}$ )	TISSUE (FLUID) PLASMA (SERUM) RATIO
SKIN	72-96	0.4	0.012	35
LUNG	72-96	4.0	0.012	>100
SPUTUM*	2-4	1.0	0.64	2
SPUTUM**	10-12	2.9	0.1	30
TONSIL***	9-18	4.5	0.03	>100
TONSIL***	180	0.9	0.006	>100
CERVIX****	19	2.8	0.04	70

<sup>1</sup> Azithromycin tissue concentrations were originally determined using 250 mg capsules.

\*Sample was obtained 2-4 hours after the first dose.

\*\*Sample was obtained 10-12 hours after the first dose.

\*\*\*Dosing regimen of two doses of 250 mg each, separated by 12 hours.

\*\*\*\*Sample was obtained 19 hours after a single 500 mg dose.

The extensive tissue distribution was confirmed by examination of additional tissues and fluids (bone, ejaculum, prostate, ovary, uterus, salpinx, stomach, liver, and gallbladder). As there are no data from adequate and well-controlled studies of azithromycin treatment of infections in these additional body sites, the clinical importance of these tissue concentration data is unknown.

Following a regimen of 500 mg on the first day and 250 mg daily for 4 days, only very low concentrations were noted in cerebrospinal fluid (less than 0.01  $\mu\text{g/mL}$ ) in the presence of non-inflamed meninges.

#### Metabolism

*In vitro* and *in vivo* studies to assess the metabolism of azithromycin have not been performed.

#### Elimination

Plasma concentrations of azithromycin following single 500 mg oral and i.v. doses declined in a polyphasic pattern with a mean apparent plasma clearance of 630 mL/min and terminal elimination half-life of 68 hours. The prolonged terminal half-life is thought to be due to extensive uptake and subsequent release of drug from tissues.

Biliary excretion of azithromycin, predominantly as unchanged drug, is a major route of elimination. Over the course of a week, approximately 6% of the administered dose appears as unchanged drug in urine.

### Special Populations

#### Renal Insufficiency

Azithromycin pharmacokinetics were investigated in 42 adults (21 to 85 years of age) with varying degrees of renal impairment. Following the oral administration of a single 1,000 mg dose of azithromycin, mean  $C_{max}$  and  $AUC_{0-120}$  increased by 5.1% and 4.2%, respectively in subjects with mild to moderate renal impairment (GFR 10 to 80 mL/min) compared to subjects with normal renal function (GFR >80 mL/min). The mean  $C_{max}$  and  $AUC_{0-120}$  increased 61% and 35%, respectively in subjects with severe renal impairment (GFR <10 mL/min) compared to subjects with normal renal function (GFR >80 mL/min). (See **DOSAGE AND ADMINISTRATION.**)

#### Hepatic Insufficiency

The pharmacokinetics of azithromycin in subjects with hepatic impairment have not been established.

#### Gender

There are no significant differences in the disposition of azithromycin between male and female subjects. No dosage adjustment is recommended based on gender.

#### Geriatric Patients

When studied in healthy elderly subjects aged 65 to 85 years, the pharmacokinetic parameters of azithromycin in elderly men were similar to those in young adults; however, in elderly women, although higher peak concentrations (increased by 30 to 50%) were observed, no significant accumulation occurred.

#### Pediatric Patients

In two clinical studies, azithromycin for oral suspension was dosed at 10 mg/kg on day 1, followed by 5 mg/kg on days 2 through 5 to two groups of children/pediatric subjects (aged 1-5 years and 5-15 years, respectively). The mean pharmacokinetic parameters on day 5 were  $C_{max}=0.216 \mu\text{g/mL}$ ,  $T_{max}=1.9$  hours, and  $AUC_{0-24}=1.822 \mu\text{g}\cdot\text{hr/mL}$  for the 1- to 5-year-old group and were  $C_{max}=0.383 \mu\text{g/mL}$ ,  $T_{max}=2.4$  hours, and  $AUC_{0-24}=3.109 \mu\text{g}\cdot\text{hr/mL}$  for the 5- to 15-year-old group.

Two clinical studies were conducted in 68 children/pediatric patients aged 3-16 years to determine the pharmacokinetics and safety of azithromycin for oral suspension in children/pediatrics. Azithromycin was administered following a low-fat breakfast.

The first study consisted of 35 pediatric patients treated with 20 mg/kg/day (maximum daily dose 500 mg) for 3 days of whom 34 patients were evaluated for pharmacokinetics.

In the second study, 33 pediatric patients received doses of 12 mg/kg/day (maximum daily dose 500 mg) for 5 days of whom 31 patients were evaluated for pharmacokinetics.

In both studies, azithromycin concentrations were determined over a 24 hour period following the last daily dose. Patients weighing above 25.0 kg in the 3-day study or 41.7 kg in the 5-day

study received the maximum adult daily dose of 500 mg. Eleven patients (weighing 25.0 kg or less) in the first study and 17 patients (weighing 41.7 kg or less) in the second study received a total dose of 60 mg/kg. The following table shows pharmacokinetic data in the subset of children/pediatric patients who received a total dose of 60 mg/kg.

Pharmacokinetic Parameter [mean (SD)]	3-Day Regimen (20 mg/kg x 3 days)	5-Day Regimen (12 mg/kg x 5 days)
n	11	17
C <sub>max</sub> (µg/mL)	1.1 (0.4)	0.5 (0.4)
T <sub>max</sub> (hr)	2.7 (1.9)	2.2 (0.8)
AUC <sub>0-24</sub> (µg·hr/mL)	7.9 (2.9)	3.9 (1.9)

The similarity of the overall exposure (AUC<sub>0-∞</sub>) between the 3-day and 5-day regimens in pediatric patients is unknown.

Single dose pharmacokinetics in ~~children given~~ pediatric subjects administered doses of 30 mg/kg have not been studied. (See **DOSAGE AND ADMINISTRATION**.)

#### Drug-Drug Interactions

Drug interaction studies were performed with azithromycin and other drugs likely to be co-administered. The effects of co-administration of azithromycin on the pharmacokinetics of other drugs are shown in Table 1 and the effect of other drugs on the pharmacokinetics of azithromycin are shown in Table 2.

Co-administration of azithromycin at therapeutic doses had a modest effect on the pharmacokinetics of the drugs listed in Table 1. No dosage adjustment of drugs listed in Table 1 is recommended when co-administered with azithromycin.

Co-administration of azithromycin with efavirenz or fluconazole had a modest effect on the pharmacokinetics of azithromycin. Nelfinavir significantly increased the C<sub>max</sub> and AUC of azithromycin. No dosage adjustment of azithromycin is recommended when administered with drugs listed in Table 2. (See **PRECAUTIONS - Drug Interactions**.)

Table 1. Drug Interactions: Pharmacokinetic Parameters for Co-administered Drugs in the Presence of Azithromycin

Co-administered Drug	Dose of Co-administered Drug	Dose of Azithromycin	n	Ratio (with/without azithromycin) of Co-administered Drug Pharmacokinetic Parameters (90% CI); No Effect = 1.00	
				Mean C <sub>max</sub>	Mean AUC
Atorvastatin	10 mg/day × 8 days	500 mg/day PO on days 6-8	12	0.83 (0.63 to 1.08)	1.01 (0.81 to 1.25)
Carbamazepine	200 mg/day × 2 days, then 200 mg BID × 18 days	500 mg/day PO for days 16-18	7	0.97 (0.88 to 1.06)	0.96 (0.88 to 1.06)
Cetirizine	20 mg/day × 11 days	500 mg PO on day 7, then 250 mg/day on days 8-11	14	1.03 (0.93 to 1.14)	1.02 (0.92 to 1.13)
Didanosine	200 mg PO BID × 21 days	1,200 mg/day PO on days 8-21	6	1.44 (0.85 to 2.43)	1.14 (0.83 to 1.57)
Efavirenz	400 mg/day × 7 days	600 mg PO on day 7	14	1.04*	0.95*
Fluconazole	200 mg PO single dose	1,200 mg PO single dose	18	1.04 (0.98 to 1.11)	1.01 (0.97 to 1.05)
Indinavir	800 mg TID × 5 days	1,200 mg PO on day 5	18	0.96 (0.86 to 1.08)	0.90 (0.81 to 1.00)
Midazolam	15 mg PO on day 3	500 mg/day PO × 3 days	12	1.27 (0.89 to 1.81)	1.26 (1.01 to 1.56)
Nelfinavir	750 mg TID × 11 days	1,200 mg PO on day 9	14	0.90 (0.81 to 1.01)	0.85 (0.78 to 0.93)
Rifabutin	300 mg/day × 10 days	500 mg PO on day 1, then 250 mg/day on days 2-10	6	See footnote below	NA
Sildenafil	100 mg on days 1 and 4	500 mg/day PO × 3 days	12	1.16 (0.86 to 1.57)	0.92 (0.75 to 1.12)
Theophylline	4 mg/kg IV on days 1, 11, 25	500 mg PO on day 7, 250 mg/day on days 8-11	10	1.19 (1.02 to 1.40)	1.02 (0.86 to 1.22)
Theophylline	300 mg PO BID × 15 days	500 mg PO on day 6, then 250 mg/day on days 7-10	8	1.09 (0.92 to 1.29)	1.08 (0.89 to 1.31)
Triazolam	0.125 mg on day 2	500 mg PO on day 1, then 250 mg/day on day 2	12	1.06*	1.02*
Trimethoprim/ Sulfamethoxazole	160 mg/800 mg/day PO × 7 days	1,200 mg PO on day 7	12	0.85 (0.75 to 0.97)/ 0.90 (0.78 to 1.03)	0.87 (0.80 to 0.95)/ 0.96 (0.88 to 1.03)
Zidovudine	500 mg/day PO × 21 days	600 mg/day PO × 14 days	5	1.12 (0.42 to 3.02)	0.94 (0.52 to 1.70)
Zidovudine	500 mg/day PO × 21 days	1,200 mg/day PO × 14 days	4	1.31 (0.43 to 3.97)	1.30 (0.69 to 2.43)

NA - Not Available

\* - 90% Confidence interval not reported

Mean rifabutin concentrations one-half day after the last dose of rifabutin were 60 ng/mL when co-administered with azithromycin and 71 ng/mL when co-administered with placebo.

Table 2. Drug Interactions: Pharmacokinetic Parameters for Azithromycin in the Presence of Co-administered Drugs (See **PRECAUTIONS - Drug Interactions.**)

Co-administered Drug	Dose of Co-administered Drug	Dose of Azithromycin	n	Ratio (with/without co-administered drug) of Azithromycin Pharmacokinetic Parameters (90% CI); No Effect = 1.00	
				Mean C <sub>max</sub>	Mean AUC
Efavirenz	400 mg/day × 7 days	600 mg PO on day 7	14	1.22 (1.04 to 1.42)	0.92*
Fluconazole	200 mg PO single dose	1,200 mg PO single dose	18	0.82 (0.66 to 1.02)	1.07 (0.94 to 1.22)
Nelfinavir	750 mg TID × 11 days	1,200 mg PO on day 9	14	2.36 (1.77 to 3.15)	2.12 (1.80 to 2.50)
Rifabutin	300 mg/day × 10 days	500 mg PO on day 1, then 250 mg/day on days 2-10	6	See footnote below	NA

NA – Not available

\* - 90% Confidence interval not reported

Mean azithromycin concentrations one day after the last dose were 53 ng/mL when coadministered with 300 mg daily rifabutin and 49 ng/mL when coadministered with placebo.

**Microbiology:** Azithromycin acts by binding to the 50S ribosomal subunit of susceptible microorganisms and, thus, interfering with microbial protein synthesis. Nucleic acid synthesis is not affected.

Azithromycin concentrates in phagocytes and fibroblasts as demonstrated by *in vitro* incubation techniques. Using such methodology, the ratio of intracellular to extracellular concentration was >30 after one hour incubation. *In vivo* studies suggest that concentration in phagocytes may contribute to drug distribution to inflamed tissues.

Azithromycin has been shown to be active against most isolates of the following microorganisms, both *in vitro* and in clinical infections as described in the **INDICATIONS AND USAGE** section.

**Aerobic and facultative gram-positive microorganisms**

- Staphylococcus aureus*
- Streptococcus agalactiae*
- Streptococcus pneumoniae*
- Streptococcus pyogenes*

NOTE: Azithromycin demonstrates cross-resistance with erythromycin-resistant gram-positive strains. Most strains of *Enterococcus faecalis* and methicillin-resistant staphylococci are resistant to azithromycin.

**Aerobic and facultative gram-negative microorganisms**

*Haemophilus ducreyi*  
*Haemophilus influenzae*  
*Moraxella catarrhalis*  
*Neisseria gonorrhoeae*

**“Other” microorganisms**

*Chlamydia pneumoniae*  
*Chlamydia trachomatis*  
*Mycoplasma pneumoniae*

Beta-lactamase production should have no effect on azithromycin activity.

The following *in vitro* data are available, **but their clinical significance is unknown.**

At least 90% of the following microorganisms exhibit an *in vitro* minimum inhibitory concentration (MIC) less than or equal to the susceptible breakpoints for azithromycin. However, the safety and effectiveness of azithromycin in treating clinical infections due to these microorganisms have not been established in adequate and well-controlled trials.

**Aerobic and facultative gram-positive microorganisms**

Streptococci (Groups C, F, G)  
Viridans group streptococci

**Aerobic and facultative gram-negative microorganisms**

*Bordetella pertussis*  
*Legionella pneumophila*

**Anaerobic microorganisms**

*Peptostreptococcus* species  
*Prevotella bivia*

**“Other” microorganisms**

*Ureaplasma urealyticum*

**Susceptibility Testing Methods:**

When available, the results of *in vitro* susceptibility test results for antimicrobial drugs used in resident hospitals should be provided to the physician as periodic reports which describe the susceptibility profile of nosocomial and community-acquired pathogens. These reports may differ from susceptibility data obtained from outpatient use, but could aid the physician in selecting the most effective antimicrobial.

**Dilution techniques:**

Quantitative methods are used to determine antimicrobial minimum inhibitory concentrations (MICs). These MICs provide estimates of the susceptibility of bacteria to antimicrobial compounds. The MICs should be determined using a standardized procedure. Standardized

procedures are based on a dilution method<sup>1,3</sup> (broth or agar) or equivalent with standardized inoculum concentrations and standardized concentrations of azithromycin powder. The MIC values should be interpreted according to criteria provided in Table 1.

**Diffusion techniques:**

Quantitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. One such standardized procedure<sup>2,3</sup> requires the use of standardized inoculum concentrations. This procedure uses paper disks impregnated with 15- $\mu$ g azithromycin to test the susceptibility of microorganisms to azithromycin. The disk diffusion interpretive criteria are provided in Table 1.

**Table 1. Susceptibility Interpretive Criteria for Azithromycin  
Susceptibility Test Result Interpretive Criteria**

Pathogen	Minimum Inhibitory Concentrations ( $\mu$ g/mL)			Disk Diffusion (zone diameters in mm)		
	S	I	R <sup>a</sup>	S	I	R <sup>a</sup>
<i>Haemophilus</i> spp.	$\leq 4$	--	--	$\geq 12$	--	--
<i>Staphylococcus aureus</i>	$\leq 2$	4	$\geq 8$	$\geq 18$	14-17	$\leq 13$
Streptococci including <i>S. pneumoniae</i> <sup>b</sup>	$\leq 0.5$	1	$\geq 2$	$\geq 18$	14-17	$\leq 13$

<sup>a</sup> The current absence of data on resistant strains precludes defining any category other than "susceptible." If strains yield MIC results other than susceptible, they should be submitted to a reference laboratory for further testing.

<sup>b</sup> Susceptibility of streptococci including *S. pneumoniae* to azithromycin and other macrolides can be predicted by testing erythromycin.

No interpretive criteria have been established for testing *Neisseria gonorrhoeae*. This species is not usually tested.

A report of "susceptible" indicates that the pathogen is likely to be inhibited if the antimicrobial compound reaches the concentrations usually achievable. A report of "intermediate" indicates that the result should be considered equivocal, and, if the microorganism is not fully susceptible to alternative, clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where high dosage of drug can be used. This category also provides a buffer zone which prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of "resistant" indicates that the pathogen is not likely to be inhibited if the antimicrobial compound reaches the concentrations usually achievable; other therapy should be selected.

*QUALITY CONTROL:*

Standardized susceptibility test procedures require the use of quality control microorganisms to control the technical aspects of the test procedures. Standard azithromycin powder should provide the following range of values noted in Table 2. Quality control microorganisms are specific strains of organisms with intrinsic biological properties. QC strains are very stable strains which will give a standard and repeatable susceptibility pattern. The specific strains used for microbiological quality control are not clinically significant.

**Table 2. Acceptable Quality Control Ranges for Azithromycin**

<b>QC Strain</b>	<b>Minimum Inhibitory Concentrations (<math>\mu\text{g/mL}</math>)</b>	<b>Disk Diffusion (zone diameters in mm)</b>
<i>Haemophilus influenzae</i> ATCC 49247	1.0-4.0	13-21
<i>Staphylococcus aureus</i> ATCC 29213	0.5-2.0	
<i>Staphylococcus aureus</i> ATCC 25923		21-26
<i>Streptococcus pneumoniae</i> ATCC 49619	0.06-0.25	19-25

**INDICATIONS AND USAGE**

ZITHROMAX<sup>®</sup> (azithromycin) is indicated for the treatment of patients with mild to moderate infections (pneumonia: see **WARNINGS**) caused by susceptible strains of the designated microorganisms in the specific conditions listed below. As recommended dosages, durations of therapy and applicable patient populations vary among these infections, please see **DOSAGE AND ADMINISTRATION** for specific dosing recommendations.

**Adults:**

**Acute bacterial exacerbations of chronic obstructive pulmonary disease** due to *Haemophilus influenzae*, *Moraxella catarrhalis* or *Streptococcus pneumoniae*.

**Acute bacterial sinusitis** due to *Haemophilus influenzae*, *Moraxella catarrhalis* or *Streptococcus pneumoniae*.

**Community-acquired pneumonia** due to *Chlamydia pneumoniae*, *Haemophilus influenzae*, *Mycoplasma pneumoniae* or *Streptococcus pneumoniae* in patients appropriate for oral therapy.

**NOTE: Azithromycin should not be used in patients with pneumonia who are judged to be inappropriate for oral therapy because of moderate to severe illness or risk factors such as any of the following:**

**patients with cystic fibrosis,  
patients with nosocomially acquired infections,  
patients with known or suspected bacteremia,  
patients requiring hospitalization,  
elderly or debilitated patients, or  
patients with significant underlying health problems that may compromise their ability to respond to their illness (including immunodeficiency or functional asplenia).**

**Pharyngitis/tonsillitis** caused by *Streptococcus pyogenes* as an alternative to first-line therapy in individuals who cannot use first-line therapy.

NOTE: Penicillin by the intramuscular route is the usual drug of choice in the treatment of *Streptococcus pyogenes* infection and the prophylaxis of rheumatic fever. ZITHROMAX<sup>®</sup> is often effective in the eradication of susceptible strains of *Streptococcus pyogenes* from the nasopharynx. Because some strains are resistant to ZITHROMAX<sup>®</sup>, susceptibility tests should be performed when patients are treated with ZITHROMAX<sup>®</sup>. Data establishing efficacy of azithromycin in subsequent prevention of rheumatic fever are not available.

**Uncomplicated skin and skin structure infections** due to *Staphylococcus aureus*, *Streptococcus pyogenes*, or *Streptococcus agalactiae*. Abscesses usually require surgical drainage.

**Urethritis and cervicitis** due to *Chlamydia trachomatis* or *Neisseria gonorrhoeae*.

**Genital ulcer disease** in men due to *Haemophilus ducreyi* (chancroid). Due to the small number of women included in clinical trials, the efficacy of azithromycin in the treatment of chancroid in women has not been established.

ZITHROMAX<sup>®</sup>, at the recommended dose, should not be relied upon to treat syphilis. Antimicrobial agents used in high doses for short periods of time to treat non-gonococcal urethritis may mask or delay the symptoms of incubating syphilis. All patients with sexually-transmitted urethritis or cervicitis should have a serologic test for syphilis and appropriate cultures for gonorrhea performed at the time of diagnosis. Appropriate antimicrobial therapy and follow-up tests for these diseases should be initiated if infection is confirmed.

Appropriate culture and susceptibility tests should be performed before treatment to determine the causative organism and its susceptibility to azithromycin. Therapy with ZITHROMAX<sup>®</sup> may be initiated before results of these tests are known; once the results become available, antimicrobial therapy should be adjusted accordingly.

**Children:** (See **PRECAUTIONS—Pediatric Use** and **CLINICAL STUDIES IN PEDIATRIC PATIENTS.**)

**Acute otitis media** caused by *Haemophilus influenzae*, *Moraxella catarrhalis* or *Streptococcus pneumoniae*. (For specific dosage recommendation, see **DOSAGE AND ADMINISTRATION.**)

**Acute bacterial sinusitis** caused by *Haemophilus influenzae*, *Moraxella catarrhalis* or *Streptococcus pneumoniae*. (For specific dosage recommendation, see **DOSAGE AND ADMINISTRATION.**)

**Community-acquired pneumonia** due to *Chlamydia pneumoniae*, *Haemophilus influenzae*, *Mycoplasma pneumoniae* or *Streptococcus pneumoniae* in patients appropriate for oral therapy. (For specific dosage recommendation, see **DOSAGE AND ADMINISTRATION.**)

**NOTE:** Azithromycin should not be used in pediatric patients with pneumonia who are judged to be inappropriate for oral therapy because of moderate to severe illness or risk factors such as any of the following:

- patients with cystic fibrosis,
- patients with nosocomially acquired infections,
- patients with known or suspected bacteremia,
- patients requiring hospitalization, or
- patients with significant underlying health problems that may compromise their ability to respond to their illness (including immunodeficiency or functional asplenia).

**Pharyngitis/tonsillitis** caused by *Streptococcus pyogenes* as an alternative to first-line therapy in individuals who cannot use first-line therapy. (For specific dosage recommendation, see **DOSAGE AND ADMINISTRATION.**)

**NOTE:** Penicillin by the intramuscular route is the usual drug of choice in the treatment of *Streptococcus pyogenes* infection and the prophylaxis of rheumatic fever. ZITHROMAX<sup>®</sup> is often effective in the eradication of susceptible strains of *Streptococcus pyogenes* from the nasopharynx. Because some strains are resistant to ZITHROMAX<sup>®</sup>, susceptibility tests should be performed when patients are treated with ZITHROMAX<sup>®</sup>. Data establishing efficacy of azithromycin in subsequent prevention of rheumatic fever are not available.

Appropriate culture and susceptibility tests should be performed before treatment to determine the causative organism and its susceptibility to azithromycin. Therapy with ZITHROMAX<sup>®</sup> may be initiated before results of these tests are known; once the results become available, antimicrobial therapy should be adjusted accordingly.

## CONTRAINDICATIONS

ZITHROMAX<sup>®</sup> is contraindicated in patients with known hypersensitivity to azithromycin, erythromycin or any macrolide antibiotic.

## WARNINGS

Serious allergic reactions, including angioedema, anaphylaxis, and dermatologic reactions including Stevens Johnson Syndrome and toxic epidermal necrolysis have been reported rarely in patients on azithromycin therapy. Although rare, fatalities have been reported. (See **CONTRAINDICATIONS**.) Despite initially successful symptomatic treatment of the allergic symptoms, when symptomatic therapy was discontinued, the allergic symptoms **recurred soon thereafter in some patients without further azithromycin exposure**. These patients required prolonged periods of observation and symptomatic treatment. The relationship of these episodes to the long tissue half-life of azithromycin and subsequent prolonged exposure to antigen is unknown at present.

If an allergic reaction occurs, the drug should be discontinued and appropriate therapy should be instituted. Physicians should be aware that reappearance of the allergic symptoms may occur when symptomatic therapy is discontinued.

**In the treatment of pneumonia, azithromycin has only been shown to be safe and effective in the treatment of community-acquired pneumonia due to *Chlamydia pneumoniae*, *Haemophilus influenzae*, *Mycoplasma pneumoniae* or *Streptococcus pneumoniae* in patients appropriate for oral therapy. Azithromycin should not be used in patients with pneumonia who are judged to be inappropriate for oral therapy because of moderate to severe illness or risk factors such as any of the following: patients with cystic fibrosis, patients with nosocomially acquired infections, patients with known or suspected bacteremia, patients requiring hospitalization, elderly or debilitated patients, or patients with significant underlying health problems that may compromise their ability to respond to their illness (including immunodeficiency or functional asplenia).**

**Pseudomembranous colitis has been reported with nearly all antibacterial agents and may range in severity from mild to life-threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhea subsequent to the administration of antibacterial agents.**

Treatment with antibacterial agents alters the normal flora of the colon and may permit overgrowth of clostridia. Studies indicate that a toxin produced by *Clostridium difficile* is a primary cause of "antibiotic-associated colitis."

After the diagnosis of pseudomembranous colitis has been established, therapeutic measures should be initiated. Mild cases of pseudomembranous colitis usually respond to discontinuation of the drug alone. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation, and treatment with an antibacterial drug clinically effective against *Clostridium difficile* colitis.

## PRECAUTIONS

**General:** Because azithromycin is principally eliminated via the liver, caution should be exercised when azithromycin is administered to patients with impaired hepatic function. Due to the limited data in subjects with GFR <10 mL/min, caution should be exercised when prescribing azithromycin in these patients. (See **CLINICAL PHARMACOLOGY - Special Populations - Renal Insufficiency.**)

The following adverse events have been reported with macrolide products: ventricular arrhythmias, including ventricular tachycardia and *torsade de pointes*, in individuals with prolonged QT intervals.

There has been a spontaneous report from the post-marketing experience of a patient with previous history of arrhythmias who experienced *torsade de pointes* and subsequent myocardial infarction following a course of azithromycin therapy.

### Information for Patients:

ZITHROMAX<sup>®</sup> tablets and oral suspension can be taken with or without food.

Patients should also be cautioned not to take aluminum- and magnesium-containing antacids and azithromycin simultaneously.

The patient should be directed to discontinue azithromycin immediately and contact a physician if any signs of an allergic reaction occur.

### Drug Interactions:

Co-administration of nelfinavir at steady-state with a single oral dose of azithromycin resulted in increased azithromycin serum concentrations. Although a dose adjustment of azithromycin is not recommended when administered in combination with nelfinavir, close monitoring for known side effects of azithromycin, such as liver enzyme abnormalities and hearing impairment, is warranted. (See **ADVERSE REACTIONS.**)

Azithromycin did not affect the prothrombin time response to a single dose of warfarin. However, prudent medical practice dictates careful monitoring of prothrombin time in all patients treated with azithromycin and warfarin concomitantly. Concurrent use of macrolides and warfarin in clinical practice has been associated with increased anticoagulant effects.

Drug interaction studies were performed with azithromycin and other drugs likely to be co-administered. (See **CLINICAL PHARMACOLOGY-Drug-Drug Interactions.**) When used in therapeutic doses, azithromycin had a modest effect on the pharmacokinetics of atorvastatin,

carbamazepine, cetirizine, didanosine, efavirenz, fluconazole, indinavir, midazolam, rifabutin, sildenafil, theophylline (intravenous and oral), triazolam, trimethoprim/sulfamethoxazole or zidovudine. Co-administration with efavirenz, or fluconazole had a modest effect on the pharmacokinetics of azithromycin. No dosage adjustment of either drug is recommended when azithromycin is coadministered with any of the above agents.

Interactions with the drugs listed below have not been reported in clinical trials with azithromycin; however, no specific drug interaction studies have been performed to evaluate potential drug-drug interaction. Nonetheless, they have been observed with macrolide products. Until further data are developed regarding drug interactions when azithromycin and these drugs are used concomitantly, careful monitoring of patients is advised:

Digoxin—elevated digoxin concentrations.

Ergotamine or dihydroergotamine—acute ergot toxicity characterized by severe peripheral vasospasm and dysesthesia.

Terfenadine, cyclosporine, hexobarbital and phenytoin concentrations.

**Laboratory Test Interactions:** There are no reported laboratory test interactions.

**Carcinogenesis, Mutagenesis, Impairment of Fertility:** Long-term studies in animals have not been performed to evaluate carcinogenic potential. Azithromycin has shown no mutagenic potential in standard laboratory tests: mouse lymphoma assay, human lymphocyte clastogenic assay, and mouse bone marrow clastogenic assay. No evidence of impaired fertility due to azithromycin was found.

**Pregnancy:** Teratogenic Effects. Pregnancy Category B: Reproduction studies have been performed in rats and mice at doses up to moderately maternally toxic dose concentrations (i.e., 200 mg/kg/day). These doses, based on a mg/m<sup>2</sup> basis, are estimated to be 4 and 2 times, respectively, the human daily dose of 500 mg. In the animal studies, no evidence of harm to the fetus due to azithromycin was found. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, azithromycin should be used during pregnancy only if clearly needed.

**Nursing Mothers:** It is not known whether azithromycin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when azithromycin is administered to a nursing woman.

**Pediatric Use:** (See **CLINICAL PHARMACOLOGY, INDICATIONS AND USAGE**, and **DOSAGE AND ADMINISTRATION**.)

Acute Otitis Media (total dosage regimen: 30 mg/kg, see **DOSAGE AND ADMINISTRATION**): Safety and effectiveness in the treatment of children with otitis media under 6 months of age have not been established.

Acute Bacterial Sinusitis (dosage regimen: 10 mg/kg on Days 1-3): Safety and effectiveness in the treatment of children with acute bacterial sinusitis under 6 months of age have not been established.

Community-Acquired Pneumonia (dosage regimen: 10 mg/kg on Day 1 followed by 5 mg/kg on Days 2-5): Safety and effectiveness in the treatment of children with community-acquired pneumonia under 6 months of age have not been established. Safety and effectiveness for pneumonia due to *Chlamydia pneumoniae* and *Mycoplasma pneumoniae* were documented in pediatric clinical trials. Safety and effectiveness for pneumonia due to *Haemophilus influenzae* and *Streptococcus pneumoniae* were not documented bacteriologically in the pediatric clinical trial due to difficulty in obtaining specimens. Use of azithromycin for these two microorganisms is supported, however, by evidence from adequate and well-controlled studies in adults.

Pharyngitis/Tonsillitis (dosage regimen: 12 mg/kg on Days 1-5): Safety and effectiveness in the treatment of children with pharyngitis/tonsillitis under 2 years of age have not been established.

**Studies evaluating the use of repeated courses of therapy have not been conducted. (See CLINICAL PHARMACOLOGY and ANIMAL TOXICOLOGY.)**

**Geriatric Use:** Pharmacokinetic parameters in older volunteers (65-85 years old) were similar to those in younger volunteers (18-40 years old) for the 5-day therapeutic regimen. Dosage adjustment does not appear to be necessary for older patients with normal renal and hepatic function receiving treatment with this dosage regimen. (See **CLINICAL PHARMACOLOGY.**)

In multiple-dose clinical trials of oral azithromycin, 9% of patients were at least 65 years of age (458/4949) and 3% of patients (144/4949) were at least 75 years of age. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in response between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

ZITHROMAX<sup>®</sup> 250 mg tablets contain 0.9 mg of sodium per tablet.

ZITHROMAX<sup>®</sup> 500 mg tablets contain 1.8 mg of sodium per tablet.

ZITHROMAX<sup>®</sup> for oral suspension 100 mg/5 mL contains 3.7 mg of sodium per 5 mL of constituted solution.

ZITHROMAX<sup>®</sup> for oral suspension 200 mg/5 mL contains 7.4 mg of sodium per 5 mL of constituted solution.

#### **ADVERSE REACTIONS**

In clinical trials, most of the reported side effects were mild to moderate in severity and were reversible upon discontinuation of the drug. Potentially serious side effects of angioedema and cholestatic jaundice were reported rarely. Approximately 0.7% of the patients (adults and children) from the 5-day multiple-dose clinical trials discontinued ZITHROMAX<sup>®</sup> (azithromycin) therapy because of treatment-related side effects. In adults given 500 mg/day for 3 days, the discontinuation rate due to treatment-related side effects was 0.6%.

In clinical trials in children given 30 mg/kg, either as a single dose or over 3 days, discontinuation from the trials due to treatment-related side effects was approximately 1%. (See **DOSAGE AND ADMINISTRATION.**) Most of the side effects leading to discontinuation were related to the gastrointestinal tract, e.g., nausea, vomiting, diarrhea, or abdominal pain. (See **CLINICAL STUDIES IN PEDIATRIC PATIENTS.**)

**Clinical:**

**Adults:**

*Multiple-dose regimens:* Overall, the most common treatment-related side effects in adult patients receiving multiple-dose regimens of ZITHROMAX<sup>®</sup> were related to the gastrointestinal system with diarrhea/loose stools (4-5%), nausea (3%) and abdominal pain (2-3%) being the most frequently reported.

No other treatment-related side effects occurred in patients on the multiple-dose regimens of ZITHROMAX<sup>®</sup> with a frequency greater than 1%. Side effects that occurred with a frequency of 1% or less included the following:

**Cardiovascular:** Palpitations, chest pain.

**Gastrointestinal:** Dyspepsia, flatulence, vomiting, melena and cholestatic jaundice.

**Genitourinary:** Monilia, vaginitis and nephritis.

**Nervous System:** Dizziness, headache, vertigo and somnolence.

**General:** Fatigue.

**Allergic:** Rash, pruritus, photosensitivity and angioedema.

*Single 1-gram dose regimen:* Overall, the most common side effects in patients receiving a single-dose regimen of 1 gram of ZITHROMAX<sup>®</sup> were related to the gastrointestinal system and were more frequently reported than in patients receiving the multiple-dose regimen.

Side effects that occurred in patients on the single one-gram dosing regimen of ZITHROMAX<sup>®</sup> with a frequency of 1% or greater included diarrhea/loose stools (7%), nausea (5%), abdominal pain (5%), vomiting (2%), dyspepsia (1%) and vaginitis (1%).

*Single 2-gram dose regimen:* Overall, the most common side effects in patients receiving a single 2-gram dose of ZITHROMAX<sup>®</sup> were related to the gastrointestinal system. Side effects that occurred in patients in this study with a frequency of 1% or greater included nausea (18%), diarrhea/loose stools (14%), vomiting (7%), abdominal pain (7%), vaginitis (2%), dyspepsia (1%) and dizziness (1%). The majority of these complaints were mild in nature.

**Children:**

*Single and Multiple-dose regimens:* The types of side effects in children were comparable to those seen in adults, with different incidence rates for the dosage regimens recommended in children.

**Acute Otitis Media:** For the recommended total dosage regimen of 30 mg/kg, the most frequent side effects ( $\geq 1\%$ ) attributed to treatment were diarrhea, abdominal pain, vomiting, nausea and

rash. (See **DOSAGE AND ADMINISTRATION** and **CLINICAL STUDIES IN PEDIATRIC PATIENTS**.)

The incidence, based on dosing regimen, is described in the table below:

Dosage Regimen	Diarrhea, %	Abdominal Pain, %	Vomiting, %	Nausea, %	Rash, %
1-day	4.3%	1.4%	4.9%	1.0%	1.0%
3-day	2.6%	1.7%	2.3%	0.4%	0.6%
5-day	1.8%	1.2%	1.1%	0.5%	0.4%

**Community-Acquired Pneumonia:** For the recommended dosage regimen of 10 mg/kg on Day 1 followed by 5 mg/kg on Days 2-5, the most frequent side effects attributed to treatment were diarrhea/loose stools, abdominal pain, vomiting, nausea and rash.

The incidence is described in the table below:

Dosage Regimen	Diarrhea/Loose stools, %	Abdominal Pain, %	Vomiting, %	Nausea, %	Rash, %
5-day	5.8%	1.9%	1.9%	1.9%	1.6%

**Pharyngitis/tonsillitis:** For the recommended dosage regimen of 12 mg/kg on Days 1-5, the most frequent side effects attributed to treatment were diarrhea, vomiting, abdominal pain, nausea and headache.

The incidence is described in the table below:

Dosage Regimen	Diarrhea, %	Abdominal Pain, %	Vomiting, %	Nausea, %	Rash, %	Headache, %
5-day	5.4%	3.4%	5.6%	1.8%	0.7%	1.1%

With any of the treatment regimens, no other treatment-related side effects occurred in children treated with ZITHROMAX<sup>®</sup> with a frequency greater than 1%. Side effects that occurred with a frequency of 1% or less included the following:

**Cardiovascular:** Chest pain.

**Gastrointestinal:** Dyspepsia, constipation, anorexia, enteritis, flatulence, gastritis, jaundice, loose stools and oral moniliasis.

**Hematologic and Lymphatic:** Anemia and leukopenia.

**Nervous System:** Headache (otitis media dosage), hyperkinesia, dizziness, agitation, nervousness and insomnia.

**General:** Fever, face edema, fatigue, fungal infection, malaise and pain.

**Allergic:** Rash and allergic reaction.

**Respiratory:** Cough increased, pharyngitis, pleural effusion and rhinitis.

**Skin and Appendages:** Eczema, fungal dermatitis, pruritus, sweating, urticaria and vesiculobullous rash.

**Special Senses:** Conjunctivitis.

**Post-Marketing Experience:**

Adverse events reported with azithromycin during the post-marketing period in adult and/or pediatric patients for which a causal relationship may not be established include:

**Allergic:** Arthralgia, edema, urticaria and angioedema.

**Cardiovascular:** Arrhythmias including ventricular tachycardia and hypotension.

**Gastrointestinal:** Anorexia, constipation, dyspepsia, flatulence, vomiting/diarrhea rarely resulting in dehydration, pseudomembranous colitis, pancreatitis, oral candidiasis and rare reports of tongue discoloration.

**General:** Asthenia, paresthesia, fatigue, malaise and anaphylaxis (rarely fatal).

**Genitourinary:** Interstitial nephritis and acute renal failure and vaginitis.

**Hematopoietic:** Thrombocytopenia.

**Liver/Biliary:** Abnormal liver function including hepatitis and cholestatic jaundice, as well as rare cases of hepatic necrosis and hepatic failure, some of which have resulted in death.

**Nervous System:** Convulsions, dizziness/vertigo, headache, somnolence, hyperactivity, nervousness, agitation and syncope.

**Psychiatric:** Aggressive reaction and anxiety.

**Skin/Appendages:** Pruritus, rarely serious skin reactions including erythema multiforme, Stevens Johnson Syndrome and toxic epidermal necrolysis.

**Special Senses:** Hearing disturbances including hearing loss, deafness and/or tinnitus and rare reports of taste perversion.

**Laboratory Abnormalities:****Adults:**

Clinically significant abnormalities (irrespective of drug relationship) occurring during the clinical trials were reported as follows: with an incidence of greater than 1%: decreased hemoglobin, hematocrit, lymphocytes, neutrophils and blood glucose; elevated serum creatine phosphokinase, potassium, ALT, GGT, AST, BUN, creatinine, blood glucose, platelet count, lymphocytes, neutrophils and eosinophils; with an incidence of less than 1%: leukopenia, neutropenia, decreased sodium, potassium, platelet count; elevated monocytes, basophils, bicarbonate, serum alkaline phosphatase, bilirubin, LDH and phosphate. The majority of subjects with elevated serum creatinine also had abnormal values at baseline.

When follow-up was provided, changes in laboratory tests appeared to be reversible.

In multiple-dose clinical trials involving more than 5000 patients, four patients discontinued therapy because of treatment-related liver enzyme abnormalities and one because of a renal function abnormality.

**Children:****One, Three and Five Day Regimens**

Laboratory data collected from comparative clinical trials employing two 3-day regimens (30 mg/kg or 60 mg/kg in divided doses over 3 days), or two 5-day regimens (30 mg/kg or 60 mg/kg

in divided doses over 5 days) were similar for regimens of azithromycin and all comparators combined, with most clinically significant laboratory abnormalities occurring at incidences of 1-5%. Laboratory data for patients receiving 30 mg/kg as a single dose were collected in one single center trial. In that trial, an absolute neutrophil count between 500-1500 cells/mm<sup>3</sup> was observed in 10/64 patients receiving 30 mg/kg as a single dose, 9/62 patients receiving 30 mg/kg given over 3 days, and 8/63 comparator patients. No patient had an absolute neutrophil count <500 cells/mm<sup>3</sup>. (See **DOSAGE AND ADMINISTRATION**.)

In multiple-dose clinical trials involving approximately 4700 pediatric patients, no patients discontinued therapy because of treatment-related laboratory abnormalities.

**DOSAGE AND ADMINISTRATION** (See **INDICATIONS AND USAGE** and **CLINICAL PHARMACOLOGY**.)

**Adults:**

Infection*	Recommended Dose/Duration of Therapy
Community-acquired pneumonia (mild severity) Pharyngitis/tonsillitis (second line therapy) Skin/skin structure (uncomplicated)	500 mg as a single dose on Day 1, followed by 250 mg once daily on Days 2 through 5.
Acute bacterial exacerbations of chronic obstructive pulmonary disease (mild to moderate)	500 mg QD x 3 days OR 500 mg as a single dose on Day 1, followed by 250 mg once daily on Days 2 through 5.
Acute bacterial sinusitis	500 mg QD x 3 days
Genital ulcer disease (chancroid)	One single 1 gram dose
Non-gonococcal urethritis and cervicitis	One single 1 gram dose
Gonococcal urethritis and cervicitis	One single 2 gram dose
* <b>DUE TO THE INDICATED ORGANISMS</b> (See <b>INDICATIONS AND USAGE</b> .)	

ZITHROMAX<sup>®</sup> tablets can be taken with or without food.

**Renal Insufficiency:**

No dosage adjustment is recommended for subjects with renal impairment (GFR ≤80 mL/min). The mean AUC<sub>0-120</sub> was similar in subjects with GFR 10-80 mL/min compared to subjects with normal renal function, whereas it increased 35% in subjects with GFR <10 mL/min compared to subjects with normal renal function. Caution should be exercised when azithromycin is administered to subjects with severe renal impairment. (See **CLINICAL PHARMACOLOGY, Special Populations, Renal Insufficiency**.)

**Hepatic Insufficiency:**

The pharmacokinetics of azithromycin in subjects with hepatic impairment have not been established. No dose adjustment recommendations can be made in patients with impaired hepatic function (See **CLINICAL PHARMACOLOGY, Special Populations, Hepatic Insufficiency**.)

No dosage adjustment is recommended based on age or gender. (See **CLINICAL PHARMACOLOGY, Special Populations.**)

**Children:**

ZITHROMAX<sup>®</sup> for oral suspension can be taken with or without food.

**Acute Otitis Media:** The recommended dose of ZITHROMAX<sup>®</sup> for oral suspension for the treatment of children with acute otitis media is 30 mg/kg given as a single dose or 10 mg/kg once daily for 3 days or 10 mg/kg as a single dose on the first day followed by 5 mg/kg/day on Days 2 through 5. (See chart below.)

**Acute Bacterial Sinusitis:** The recommended dose of ZITHROMAX<sup>®</sup> for oral suspension for the treatment of children with acute bacterial sinusitis is 10 mg/kg once daily for 3 days. (See chart below.)

**Community-Acquired Pneumonia:** The recommended dose of ZITHROMAX<sup>®</sup> for oral suspension for the treatment of children with community-acquired pneumonia is 10 mg/kg as a single dose on the first day followed by 5 mg/kg on Days 2 through 5. (See chart below.)

**PEDIATRIC DOSAGE GUIDELINES FOR OTITIS MEDIA, ACUTE BACTERIAL SINUSITIS AND COMMUNITY-ACQUIRED PNEUMONIA  
(Age 6 months and above, see PRECAUTIONS—Pediatric Use.)  
Based on Body Weight**

OTITIS MEDIA AND COMMUNITY-ACQUIRED PNEUMONIA: 5-Day Regimen*							
Dosing Calculated on 10 mg/kg/day Day 1 and 5 mg/kg/day Days 2 to 5.							
Weight		100 mg/5 mL		200 mg/5 mL		Total mL per Treatment Course	Total mg per Treatment Course
Kg	Lbs.	Day 1	Days 2-5	Day 1	Days 2-5		
5	11	2.5 mL (½ tsp)	1.25 mL (¼ tsp)			7.5 mL	150 mg
10	22	5 mL (1 tsp)	2.5 mL (½ tsp)			15 mL	300 mg
20	44			5 mL (1 tsp)	2.5 mL (½ tsp)	15 mL	600 mg
30	66			7.5 mL (1½ tsp)	3.75 mL (¾ tsp)	22.5 mL	900 mg
40	88			10 mL (2 tsp)	5 mL (1 tsp)	30 mL	1200 mg
50 and above	110 and above			12.5 mL (2½ tsp)	6.25 mL (1¼ tsp)	37.5 mL	1500 mg

\* Effectiveness of the 3-day or 1-day regimen in children with community-acquired pneumonia has not been established.

<b>OTITIS MEDIA AND ACUTE BACTERIAL SINUSITIS: (3-Day Regimen)</b>					
<b>Dosing Calculated on 10 mg/kg/day</b>					
<b>Weight</b>		<b>100 mg/5 mL</b>	<b>200 mg/5 mL</b>	<b>Total mL per Treatment Course</b>	<b>Total mg per Treatment Course</b>
<b>Kg</b>	<b>Lbs.</b>	<b>Day 1-3</b>	<b>Day 1-3</b>		
5	11	2.5 mL (1/2 tsp)		7.5 mL	150 mg
10	22	5 mL (1 tsp)		15 mL	300 mg
20	44		5 mL (1 tsp)	15 mL	600 mg
30	66		7.5 mL (1 ½ tsp)	22.5 mL	900 mg
40	88		10 mL (2 tsp)	30 mL	1200 mg
50 and above	110 and above		12.5 mL (2 ½ tsp)	37.5 mL	1500 mg

<b>OTITIS MEDIA: (1-Day Regimen)</b>				
<b>Dosing Calculated on 30 mg/kg as a single dose</b>				
<b>Weight</b>		<b>200 mg/5 mL</b>	<b>Total mL per Treatment Course</b>	<b>Total mg per Treatment Course</b>
<b>Kg</b>	<b>Lbs.</b>	<b>Day 1</b>		
5	11	3.75 mL (3/4 tsp)	3.75 mL	150 mg
10	22	7.5 mL (1 ½ tsp)	7.5 mL	300 mg
20	44	15 mL (3 tsp)	15 mL	600 mg
30	66	22.5 mL (4 ½ tsp)	22.5 mL	900 mg
40	88	30 mL (6 tsp)	30 mL	1200 mg
50 and above	110 and above	37.5 mL (7 ½ tsp)	37.5 mL	1500 mg

The safety of re-dosing azithromycin in children who vomit after receiving 30 mg/kg as a single dose has not been established. In clinical studies involving 487 patients with acute otitis media given a single 30 mg/kg dose of azithromycin, eight patients who vomited within 30 minutes of dosing were re-dosed at the same total dose.

**Pharyngitis/Tonsillitis:** The recommended dose of ZITHROMAX® for children with pharyngitis/tonsillitis is 12 mg/kg once daily for 5 days. (See chart below.)

**PEDIATRIC DOSAGE GUIDELINES FOR PHARYNGITIS/TONSILLITIS**  
**(Age 2 years and above, see PRECAUTIONS—Pediatric Use.)**  
**Based on Body Weight**

<b>PHARYNGITIS/TONSILLITIS: (5-Day Regimen)</b>				
<b>Dosing Calculated on 12 mg/kg/day for 5 days.</b>				
<b>Weight</b>		<b>200 mg/5 mL</b>	<b>Total mL per Treatment Course</b>	<b>Total mg per Treatment Course</b>
<b>Kg</b>	<b>Lbs.</b>	<b>Day 1-5</b>		
8	18	2.5 mL (½ tsp)	12.5 mL	500 mg
17	37	5 mL (1 tsp)	25 mL	1000 mg
25	55	7.5 mL (1½ tsp)	37.5 mL	1500 mg
33	73	10 mL (2 tsp)	50 mL	2000 mg
40	88	12.5 mL (2½ tsp)	62.5 mL	2500 mg

Constituting instructions for ZITHROMAX® Oral Suspension, 300, 600, 900, 1200 mg bottles. The table below indicates the volume of water to be used for constitution:

<u>Amount of water to be added</u>	<u>Total volume after constitution (azithromycin content)</u>	<u>Azithromycin concentration after constitution</u>
9 mL (300 mg)	15 mL (300 mg)	100 mg/5 mL
9 mL (600 mg)	15 mL (600 mg)	200 mg/5 mL
12 mL (900 mg)	22.5 mL (900 mg)	200 mg/5 mL
15 mL (1200 mg)	30 mL (1200 mg)	200 mg/5 mL

Following constitution, and for use with the oral syringe, the supplied Press in Bottle Adapter should be inserted into the neck of the bottle then sealed with the original closure. Shake well before each use. Oversized bottle provides shake space. Keep tightly closed.

Use only the dosing device provided to measure the correct amount of suspension. (See **HOW SUPPLIED**.) The dosing device may need to be filled multiple times to provide the complete dose prescribed. Rinse the device with water after the complete daily dose has been administered.

After mixing, store suspension at 5° to 30°C (41° to 86°F) and use within 10 days. Discard after full dosing is completed.

## HOW SUPPLIED

ZITHROMAX<sup>®</sup> 250 mg tablets are supplied as pink modified capsular shaped, engraved, film-coated tablets containing azithromycin dihydrate equivalent to 250 mg of azithromycin. ZITHROMAX<sup>®</sup> 250 mg tablets are engraved with "PFIZER" on one side and "306" on the other. These are packaged in bottles and blister cards of 6 tablets (Z-PAKS<sup>®</sup>) as follows:

Bottles of 30	NDC 0069-3060-30
Boxes of 3 (Z-PAKS <sup>®</sup> of 6)	NDC 0069-3060-75
Unit Dose package of 50	NDC 0069-3060-86

ZITHROMAX<sup>®</sup> 500 mg tablets are supplied as pink modified capsular shaped, engraved, film-coated tablets containing azithromycin dihydrate equivalent to 500 mg of azithromycin. ZITHROMAX<sup>®</sup> 500 mg tablets are engraved with "Pfizer" on one side and "ZTM500" on the other. These are packaged in bottles and blister cards of 3 tablets (TRI-PAKS<sup>™</sup>) as follows:

Bottles of 30	NDC 0069-3070-30
Boxes of 3 (TRI-PAKS <sup>™</sup> of 3 tablets)	NDC 0069-3070-75
Unit Dose package of 50	NDC 0069-3070-86

ZITHROMAX<sup>®</sup> tablets should be stored between 15° to 30°C (59° to 86°F).

ZITHROMAX<sup>®</sup> for oral suspension after constitution contains a flavored suspension. ZITHROMAX<sup>®</sup> for oral suspension is supplied to provide 100 mg/5 mL or 200 mg/5 mL suspension in bottles with accompanying calibrated dosing device as follows:

<u>Azithromycin contents per bottle</u>	<u>NDC</u>
300 mg	0069-3110-19
600 mg	0069-3120-19
900 mg	0069-3130-19
1200 mg	0069-3140-19

See **DOSAGE AND ADMINISTRATION** for constitution instructions with each bottle type.

Storage: Store dry powder below 30°C (86°F). Store constituted suspension between 5° to 30°C (41° to 86°F) and discard when full dosing is completed.

## CLINICAL STUDIES (See **INDICATIONS AND USAGE** and **Pediatric Use**.)

### Pediatric Patients

From the perspective of evaluating pediatric clinical trials, Days 11-14 were considered on-therapy evaluations because of the extended half-life of azithromycin. Day 11-14 data are provided for clinical guidance. Day 24-32 evaluations were considered the primary test of cure endpoint.

## Acute Otitis Media

### Safety and efficacy using azithromycin 30 mg/kg given over 5 days

#### Protocol 1

In a double-blind, controlled clinical study of acute otitis media performed in the United States, azithromycin (10 mg/kg on Day 1 followed by 5 mg/kg on Days 2-5) was compared to amoxicillin/clavulanate potassium (4:1). For the 553 patients who were evaluated for clinical efficacy, the clinical success rate (i.e., cure plus improvement) at the Day 11 visit was 88% for azithromycin and 88% for the control agent. For the 521 patients who were evaluated at the Day 30 visit, the clinical success rate was 73% for azithromycin and 71% for the control agent.

In the safety analysis of the above study, the incidence of treatment-related adverse events, primarily gastrointestinal, in all patients treated was 9% with azithromycin and 31% with the control agent. The most common side effects were diarrhea/loose stools (4% azithromycin vs. 20% control), vomiting (2% azithromycin vs. 7% control), and abdominal pain (2% azithromycin vs. 5% control).

#### Protocol 2

In a non-comparative clinical and microbiologic trial performed in the United States, where significant rates of beta-lactamase producing organisms (35%) were found, 131 patients were evaluable for clinical efficacy. The combined clinical success rate (i.e., cure and improvement) at the Day 11 visit was 84% for azithromycin. For the 122 patients who were evaluated at the Day 30 visit, the clinical success rate was 70% for azithromycin.

Microbiologic determinations were made at the pre-treatment visit. Microbiology was not reassessed at later visits. The following presumptive bacterial/clinical cure outcomes (i.e., clinical success) were obtained from the evaluable group:

#### Presumed Bacteriologic Eradication

	Day 11	Day 30
	Azithromycin	Azithromycin
<i>S. pneumoniae</i>	61/74 (82%)	40/56 (71%)
<i>H. influenzae</i>	43/54 (80%)	30/47 (64%)
<i>M. catarrhalis</i>	28/35 (80%)	19/26 (73%)
<i>S. pyogenes</i>	11/11 (100%)	7/7
Overall	177/217 (82%)	97/137 (73%)

In the safety analysis of this study, the incidence of treatment-related adverse events, primarily gastrointestinal, in all patients treated was 9%. The most common side effect was diarrhea (4%).

#### Protocol 3

In another controlled comparative clinical and microbiologic study of otitis media performed in the United States, azithromycin was compared to amoxicillin/clavulanate potassium (4:1). This study utilized two of the same investigators as Protocol 2 (above), and these two investigators enrolled 90% of the patients in Protocol 3. For this reason, Protocol 3 was not considered to be

an independent study. Significant rates of beta-lactamase producing organisms (20%) were found. Ninety-two (92) patients were evaluable for clinical and microbiologic efficacy. The combined clinical success rate (i.e., cure and improvement) of those patients with a baseline pathogen at the Day 11 visit was 88% for azithromycin vs. 100% for control; at the Day 30 visit, the clinical success rate was 82% for azithromycin vs. 80% for control.

Microbiologic determinations were made at the pre-treatment visit. Microbiology was not reassessed at later visits. At the Day 11 and Day 30 visits, the following presumptive bacterial/clinical cure outcomes (i.e., clinical success) were obtained from the evaluable group:

Presumed Bacteriologic Eradication

	Day 11		Day 30	
	Azithromycin	Control	Azithromycin	Control
<i>S. pneumoniae</i>	25/29 (86%)	26/26 (100%)	22/28 (79%)	18/22 (82%)
<i>H. influenzae</i>	9/11 (82%)	9/9	8/10 (80%)	6/8
<i>M. catarrhalis</i>	7/7	5/5	5/5	2/3
<i>S. pyogenes</i>	2/2	5/5	2/2	4/4
Overall	43/49 (88%)	45/45 (100%)	37/45 (82%)	30/37 (81%)

In the safety analysis of the above study, the incidence of treatment-related adverse events, primarily gastrointestinal, in all patients treated was 4% with azithromycin and 31% with the control agent. The most common side effect was diarrhea/loose stools (2% azithromycin vs. 29% control).

**Safety and efficacy using azithromycin 30 mg/kg given over 3 days**

Protocol 4

In a double-blind, controlled, randomized clinical study of acute otitis media in children from 6 months to 12 years of age, azithromycin (10 mg/kg per day for 3 days) was compared to amoxicillin/clavulanate potassium (7:1) in divided doses q12h for 10 days. Each child received active drug and placebo matched for the comparator.

For the 366 patients who were evaluated for clinical efficacy at the Day 12 visit, the clinical success rate (i.e., cure plus improvement) was 83% for azithromycin and 88% for the control agent. For the 362 patients who were evaluated at the Day 24-28 visit, the clinical success rate was 74% for azithromycin and 69% for the control agent.

In the safety analysis of the above study, the incidence of treatment-related adverse events, primarily gastrointestinal, in all patients treated was 10.6% with azithromycin and 20.0% with the control agent. The most common side effects were diarrhea/loose stools (5.9% azithromycin vs. 14.6% control), vomiting (2.1% azithromycin vs. 1.1% control), and rash (0.0% azithromycin vs. 4.3% control).

**Safety and efficacy using azithromycin 30 mg/kg given as a single dose**

Protocol 5

A double blind, controlled, randomized trial was performed at nine clinical centers. Infants and children from 6 months to 12 years of age were randomized 1:1 to treatment with either azithromycin (given at 30 mg/kg as a single dose on Day 1) or amoxicillin/clavulanate potassium (7:1), divided q12h for 10 days. Each child received active drug, and placebo matched for the comparator.

Clinical response (Cure, Improvement, Failure) was evaluated at End of Therapy (Day 12-16) and Test of Cure (Day 28-32). Safety was evaluated throughout the trial for all treated subjects. For the 321 subjects who were evaluated at End of Treatment, the clinical success rate (cure plus improvement) was 87% for azithromycin, and 88% for the comparator. For the 305 subjects who were evaluated at Test of Cure, the clinical success rate was 75% for both azithromycin and the comparator.

In the safety analysis, the incidence of treatment-related adverse events, primarily gastrointestinal, was 16.8% with azithromycin, and 22.5% with the comparator. The most common side effects were diarrhea (6.4% with azithromycin vs. 12.7% with the comparator), vomiting (4% with each agent), rash (1.7% with azithromycin vs. 5.2% with the comparator) and nausea (1.7% with azithromycin vs. 1.2% with the comparator).

#### Protocol 6

In a non-comparative clinical and microbiological trial, 248 patients from 6 months to 12 years of age with documented acute otitis media were dosed with a single oral dose of azithromycin (30 mg/kg on Day 1).

For the 240 patients who were evaluable for clinical modified Intent-to-Treat (MITT) analysis, the clinical success rate (i.e., cure plus improvement) at Day 10 was 89% and for the 242 patients evaluable at Day 24-28, the clinical success rate (cure) was 85%.

#### Presumed Bacteriologic Eradication

	Day 10	Day 24-28
<i>S. pneumoniae</i>	70/76 (92%)	67/76 (88%)
<i>H. influenzae</i>	30/42 (71%)	28/44 (64%)
<i>M. catarrhalis</i>	10/10 (100%)	10/10 (100%)
Overall	110/128 (86%)	105/130 (81%)

In the safety analysis of this study, the incidence of treatment-related adverse events, primarily gastrointestinal, in all the subjects treated was 12.1%. The most common side effects were vomiting (5.6%), diarrhea (3.2%), and abdominal pain (1.6%).

#### Pharyngitis/Tonsillitis

In three double-blind controlled studies, conducted in the United States, azithromycin (12 mg/kg once a day for 5 days) was compared to penicillin V (250 mg three times a day for 10 days) in the treatment of pharyngitis due to documented Group A  $\beta$ -hemolytic streptococci (GABHS or *S. pyogenes*). Azithromycin was clinically and microbiologically statistically superior to

penicillin at Day 14 and Day 30 with the following clinical success (i.e., cure and improvement) and bacteriologic efficacy rates (for the combined evaluable patient with documented GABHS):

Three U.S. Streptococcal Pharyngitis Studies  
Azithromycin vs. Penicillin V  
EFFICACY RESULTS

	Day 14	Day 30
<b>Bacteriologic Eradication:</b>		
Azithromycin	323/340 (95%)	255/330 (77%)
Penicillin V	242/332 (73%)	206/325 (63%)
<b>Clinical Success (Cure plus improvement):</b>		
Azithromycin	336/343 (98%)	310/330 (94%)
Penicillin V	284/338 (84%)	241/325 (74%)

Approximately 1% of azithromycin-susceptible *S. pyogenes* isolates were resistant to azithromycin following therapy.

The incidence of treatment-related adverse events, primarily gastrointestinal, in all patients treated was 18% on azithromycin and 13% on penicillin. The most common side effects were diarrhea/loose stools (6% azithromycin vs. 2% penicillin), vomiting (6% azithromycin vs. 4% penicillin), and abdominal pain (3% azithromycin vs. 1% penicillin).

### Adult Patients

#### Acute Bacterial Exacerbations of Chronic Obstructive Pulmonary Disease

In a randomized, double-blind controlled clinical trial of acute exacerbation of chronic bronchitis (AECB), azithromycin (500 mg once daily for 3 days) was compared with clarithromycin (500 mg twice daily for 10 days). The primary endpoint of this trial was the clinical cure rate at Day 21-24. For the 304 patients analyzed in the modified intent to treat analysis at the Day 21-24 visit, the clinical cure rate for 3 days of azithromycin was 85% (125/147) compared to 82% (129/157) for 10 days of clarithromycin.

The following outcomes were the clinical cure rates at the Day 21-24 visit for the bacteriologically evaluable patients by pathogen:

Pathogen	Azithromycin (3 Days)	Clarithromycin (10 Days)
<i>S. pneumoniae</i>	29/32 (91%)	21/27 (78%)
<i>H. influenzae</i>	12/14 (86%)	14/16 (88%)
<i>M. catarrhalis</i>	11/12 (92%)	12/15 (80%)

In the safety analysis of this study, the incidence of treatment-related adverse events, primarily gastrointestinal, were comparable between treatment arms (25% with azithromycin and 29% with clarithromycin). The most common side effects were diarrhea, nausea and abdominal pain

with comparable incidence rates for each symptom of 5-9% between the two treatment arms. (See **ADVERSE REACTIONS**.)

### **Acute Bacterial Sinusitis**

In a randomized, double-blind, double-dummy controlled clinical trial of acute bacterial sinusitis, azithromycin (500 mg once daily for 3 days) was compared with amoxicillin/clavulanate (500/125 mg tid for 10 days). Clinical response assessments were made at Day 10 and Day 28. The primary endpoint of this trial was prospectively defined as the clinical cure rate at Day 28. For the 594 patients analyzed in the modified intent to treat analysis at the Day 10 visit, the clinical cure rate for 3 days of azithromycin was 88% (268/303) compared to 85% (248/291) for 10 days of amoxicillin/clavulanate. For the 586 patients analyzed in the modified intent to treat analysis at the Day 28 visit, the clinical cure rate for 3 days of azithromycin was 71.5% (213/298) compared to 71.5% (206/288), with a 97.5% confidence interval of -8.4 to 8.3, for 10 days of amoxicillin/clavulanate.

In this comparative trial, 99% of patients receiving treatment with azithromycin and 82% of patients receiving treatment with amoxicillin/clavulanate completed active study therapy as prescribed (p<0.0001.)

In the safety analysis of this study, the overall incidence of treatment-related adverse events, primarily gastrointestinal, was lower in the azithromycin treatment arm (31%) than in the amoxicillin/clavulanate arm (51%). The most common side effects were diarrhea (17% in the azithromycin arm vs. 32% in the amoxicillin/clavulanate arm), and nausea (7% in the azithromycin arm vs. 12% in the amoxicillin/clavulanate arm). (See **ADVERSE REACTIONS**).

In an open label, noncomparative study requiring baseline transantral sinus punctures the following outcomes were the clinical success rates at the Day 7 and Day 28 visits for the modified intent to treat patients administered 500 mg of azithromycin once daily for 3 days with the following pathogens:

<b>Pathogen</b>	<b>Azithromycin (500 mg per day for 3 Days)</b>	
	<b>Day 7</b>	<b>Day 28</b>
<i>S. pneumoniae</i>	23/26 (88%)	21/25 (84%)
<i>H. influenzae</i>	28/32 (87%)	24/32 (75%)
<i>M. catarrhalis</i>	14/15 (93%)	13/14 (93%)

The overall incidence of treatment-related adverse events in the noncomparative study was 21% in modified intent to treat patients treated with azithromycin at 500 mg once daily for 3 days with the most common side effects being diarrhea (9%), abdominal pain (4%) and nausea (3%). (See **ADVERSE REACTIONS**).

An azithromycin regimen of 500 mg for 6 days was evaluated in the double-blind and open studies described above. In the double blind comparative study, this regimen showed equivalent efficacy to amoxicillin/clavulanate (500/125 mg tid for 10 days), with a toleration profile comparable to 500 mg azithromycin administered for 3 days. In the open label study, clinical response rates by pathogen for the modified intent to treat patients administered 500 mg of azithromycin once daily for 6 days were:

Pathogen	Azithromycin (500 mg per day for 6 Days)	
	Day 7	Day 28
	<i>S. pneumoniae</i>	39/42 (93%)
<i>H. influenzae</i>	27/28 (96%)	24/28 (86%)
<i>M. catarrhalis</i>	19/19 (100%)	15/19 (79%)

#### ANIMAL TOXICOLOGY

Phospholipidosis (intracellular phospholipid accumulation) has been observed in some tissues of mice, rats, and dogs given multiple doses of azithromycin. It has been demonstrated in numerous organ systems (e.g., eye, dorsal root ganglia, liver, gallbladder, kidney, spleen, and pancreas) in dogs treated with azithromycin at doses which, expressed on the basis of mg/m<sup>2</sup>, are approximately equal to the recommended adult human dose, and in rats treated at doses approximately one-sixth of the recommended adult human dose. This effect has been shown to be reversible after cessation of azithromycin treatment. Phospholipidosis has been observed to a similar extent in the tissues of neonatal rats and dogs given daily doses of azithromycin ranging from 10 days to 30 days. Based on the pharmacokinetic data, phospholipidosis has been seen in the rat (30 mg/kg dose) at observed C<sub>max</sub> value of 1.3 µg/mL (six times greater than the observed C<sub>max</sub> of 0.216 µg/mL at the pediatric dose of 10 mg/kg). Similarly, it has been shown in the dog (10 mg/kg dose) at observed C<sub>max</sub> value of 1.5 µg/mL (seven times greater than the observed same C<sub>max</sub> and drug dose in the studied pediatric population). On a mg/m<sup>2</sup> basis, 30 mg/kg dose in the neonatal rat (135 mg/m<sup>2</sup>) and 10 mg/kg dose in the neonatal dog (79 mg/m<sup>2</sup>) are approximately 0.5 and 0.3 times, respectively, the recommended dose in the pediatric patients with an average body weight of 25 kg. Phospholipidosis, similar to that seen in the adult animals, is reversible after cessation of azithromycin treatment. The significance of these findings for animals and for humans is unknown.

#### REFERENCES:

1. National Committee for Clinical Laboratory Standards, *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically* – Fifth Edition. Approved Standard NCCLS Document M7-A5, Vol. 20, No. 2 (ISBN 1-56238-394-9). NCCLS, 940 West Valley Road, Suite 1400, Wayne, PA 19087-1898, January 2000.
2. National Committee for Clinical Laboratory Standards, *Performance Standards for Antimicrobial Disk Susceptibility Tests* - Seventh Edition. Approved Standard NCCLS Document M2-A7, Vol. 20, No. 1 (ISBN 1-56238-393-0). NCCLS, 940 West Valley Road, Suite 1400, Wayne, PA 19087-1898, January 2000.

3. National Committee for Clinical Laboratory Standards. *Performance Standards for Antimicrobial Susceptibility Testing* – Eleventh Informational Supplement. NCCLS Document M100-S11, Vol. 21, No. 1 (ISBN 1-56238-426-0). NCCLS, 940 West Valley Road, Suite 1400, Wayne, PA 19087-1898, January 2001.

**Rx only**

Licensed from Pliva

©2003 PFIZER INC

*Distributed by:*



**Pfizer Labs**

Division of Pfizer Inc, NY, NY 10017

70-5179-00-1.1

Revised January 2003

-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/  
-----

Charles Bonapace  
1/13/04 12:46:47 PM  
BIOPHARMACEUTICS

Venkateswar Jarugula  
1/13/04 01:21:49 PM  
BIOPHARMACEUTICS

4 pages redacted from this section of  
the approval package consisted of draft labeling