

CENTER FOR DRUG EVALUATION AND RESEARCH

APPLICATION NUMBER:
ANDA 76-387

APPROVED LABELING



CLOTRIMAZOLE TROCHE
(clotrimazole lozenges)



JUL 19

10001777/02

ROXANE LABORATORIES, INC.

CLOTRIMAZOLE TROCHE
(clotrimazole lozenges)


FOR TOPICAL ORAL ADMINISTRATION

R_x only

DESCRIPTION

Each Clotrimazole Troche (clotrimazole lozenges) contains 10 mg clotrimazole [1-(o-chloro- α,α -diphenylbenzyl)imidazole], a synthetic antifungal agent, for topical use in the mouth.

Structural Formula:



C₂₂H₁₇ClN₂

The troche dosage form is a large, slowly dissolving tablet (lozenge) containing 10 mg of clotrimazole dispersed in dextrates, microcrystalline cellulose, modified cellulose gum, povidone, and magnesium stearate.

CLINICAL PHARMACOLOGY

Clotrimazole is a broad-spectrum antifungal agent that inhibits the growth of pathogenic yeasts by altering the permeability of cell membranes. The action of clotrimazole is fungistatic at concentrations of drug up to 20 mcg/mL and may be fungicidal *in vitro* against *Candida albicans* and other species of the genus *Candida* at higher concentrations. No single-step or multiple-step resistance to clotrimazole has developed during successive passages of *Candida albicans* in the laboratory; however, individual organism tolerance has been observed during successive passages in the laboratory. Such *in vitro* tolerance has resolved once the organism has been removed from the antifungal environment.

After oral administration of a 10 mg clotrimazole troche to healthy volunteers, concentrations sufficient to inhibit most species of *Candida* persist in saliva for up to three hours following the approximately 30 minutes needed for a troche to dissolve. The long term persistence of drug in saliva appears to be related to the slow release of clotrimazole from the oral mucosa to which the drug is apparently bound. Repetitive dosing at three hour intervals maintains salivary levels above the minimum inhibitory concentrations of most strains of *Candida*; however, the relationship between *in vitro* susceptibility of pathogenic fungi to clotrimazole and prophylaxis or cure of infections in humans has not been established.

In another study, the mean serum concentrations were 4.98 \pm 3.7 and 3.23 \pm 1.4 nanograms/mL of clotrimazole at 30 and 60 minutes, respectively, after administration as a troche.

INDICATIONS AND USAGE

Clotrimazole Troches (clotrimazole lozenges) are indicated for the local treatment of oropharyngeal candidiasis. The diagnosis should be confirmed by a KOH smear and/or culture prior to treatment.

Clotrimazole Troches (clotrimazole lozenges) are also indicated prophylactically to reduce the incidence of oropharyngeal candidiasis in patients immunocompromised by conditions that include chemotherapy, radiotherapy, or steroid therapy utilized in the treatment of leukemia, solid tumors, or renal transplantation. There are no data from adequate and well-controlled trials to establish the safety and efficacy of this product for prophylactic use in patients immunocompromised by etiologies other than those listed in the previous sentence. (See **DOSAGE AND ADMINISTRATION**)

CONTRAINDICATIONS

Clotrimazole Troches (clotrimazole lozenges) are contraindicated in patients who are hypersensitive to any of its components.

WARNING

Clotrimazole Troches (clotrimazole lozenges) are not indicated for the treatment of systemic mycoses including systemic candidiasis.

PRECAUTIONS

Abnormal liver function tests have been reported in patients treated with clotrimazole troches; elevated SGOT levels were reported in about 15% of patients in the clinical trials. In most cases the elevations were minimal and it was often impossible to distinguish effects of clotrimazole from those of other therapy and the underlying disease (malignancy in most cases). Periodic assessment of hepatic function is advisable particularly in patients with pre-existing hepatic impairment.

Since patients must be instructed to allow each troche to dissolve slowly in the mouth in order to achieve maximum effect of the medication, they must be of such an age and physical and/or mental condition to comprehend such instructions.

Carcinogenesis: An 18 month dosing study with clotrimazole in rats has not revealed any carcinogenic effect.

Use in Pregnancy: Pregnancy Category C:

Clotrimazole has been shown to be embryotoxic in rats and mice when given in doses 100 times the adult human dose (in mg/kg), possibly secondary to maternal toxicity. The drug was not teratogenic in mice, rabbits, and rats when given in doses up to 200,180, and 100 times the human dose.

Clotrimazole given orally to mice from nine weeks before mating through weaning at a dose 120 times the human dose was associated with impairment of mating, decreased number of viable young, and decreased survival to weaning. No effects were observed at 60 times the human dose. When the drug was given to rats during a similar time period at 50 times the human dose, there was a slight decrease in the number of pups per litter and decreased pup viability.

There are no adequate and well controlled studies in pregnant women. Clotrimazole troches should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

PEDIATRIC USE

Safety and effectiveness of clotrimazole in children below the age of 3 years have not been established; therefore, its use in such patients is not recommended.

The safety and efficacy of the prophylactic use of clotrimazole troches in children have not been established.

GERIATRIC USE

Clinical studies of clotrimazole did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

ADVERSE REACTIONS

Abnormal liver function tests have been reported in patients treated with clotrimazole troches; elevated SGOT levels were reported in about 15% of patients in the clinical trials (See **PRECAUTIONS**).

Nausea, vomiting, unpleasant mouth sensations and pruritus have also been reported with the use of the troche.

OVERDOSAGE

No data available.

DRUG ABUSE AND DEPENDENCE

No data available.

DOSAGE AND ADMINISTRATION

Clotrimazole Troches (clotrimazole lozenges) are administered only as a lozenge that must be slowly dissolved in the mouth. The recommended dose is one troche five times a day for fourteen consecutive days. Only limited data are available on the safety and effectiveness of the clotrimazole troche after prolonged administration; therefore, therapy should be limited to short term use, if possible.

For prophylaxis to reduce the incidence of oropharyngeal candidiasis in patients immunocompromised by conditions that include chemotherapy, radiotherapy, or steroid therapy utilized in the treatment of leukemia, solid tumors, or renal transplantation, the recommended dose is one troche three times daily for the duration of chemotherapy or until steroids are reduced to maintenance levels.

HOW SUPPLIED

Clotrimazole Troches (clotrimazole lozenges) are white, flat face tablets that are plain on one side and embossed with "54/552" on the other side.

0054-4146-22	10 mg white tablet, bottle of 70
0054-4146-23	10 mg white tablet, bottle of 140
0054-4146-29	10 mg white tablet, bottle of 500
0054-8146-22	10 mg white tablet, 7 x 10 unit dose tablets

Store below 86°F (30°C).
Avoid freezing.

10001777/02

Revised December 2002
© RLI, 2002



ENLARGED TO 110%
BY FOIA STAFF

DOSAGE: Each tablet must be slowly dissolved in the mouth.

See accompanying literature for complete information.

Dispense in tight containers (USP).



Store below 86°F (30°C). Avoid freezing.

NDC 0054-4146-22 70 Tablets

EXP. LOT

**CLOTRIMAZOLE
TROCHE**
(clotrimazole lozenges)

10 mg




Roxane
Laboratories, Inc.
Columbus, Ohio 43216

10001775/01
© RLI, 2002

NDC 0054-4146-23 140 Tablets

**CLOTRIMAZOLE
TROCHE**
(clotrimazole lozenges)

10 mg



LOT
EXP.

See accompanying literature for complete information.

Dispense in tight containers (USP).

Store below 86°F (30°C).
Avoid freezing.

Roxane
Laboratories, Inc.
Columbus, Ohio 43216

10001773/01

© RLI, 2002

NDC 0054-4146-29

500 Tablets

**CLOTRIMAZOLE
TROCHE**
(clotrimazole lozenges)

10 mg

Each troche contains
10 mg of clotrimazole (100 mcg of
clotrimazole base) and 0.5 mg of
DOSAGE: Each troche is to be dissolved
in the mouth.

APPROVED

See accompanying literature for
complete information.

Dispense in tight containers (USP).

Store below 86°F (30°C). Avoid
freezing.

10001774/01



Roxane
Laboratories, Inc.
Columbus, Ohio 43216

© RLI, 2002



LOT
EXP.

**CLOTRIMAZOLE
TROCHE/LOZENGES**
Tablet **10** mg

LOT XXXXXX EXP. XXXXXX
N 0054-8146-22 Roxane • Cols., OH



**CLOTRIMAZOLE
TROCHE/LOZENGES**
Tablet **10** mg

LOT XXXXXX EXP. XXXXXX
N 0054-8146-22 Roxane • Cols., OH



**CLOTRIMAZOLE
TROCHE/LOZENGES**
Tablet **10** mg

LOT XXXXXX EXP. XXXXXX
N 0054-8146-22 Roxane • Cols., OH



**CLOTRIMAZOLE
TROCHE/LOZENGES**
Tablet **10** mg

LOT XXXXXX EXP. XXXXXX
N 0054-8146-22 Roxane • Cols., OH



**CLOTRIMAZOLE
TROCHE/LOZENGES**
Tablet **10** mg

LOT XXXXXX EXP. XXXXXX
N 0054-8146-22 Roxane • Cols., OH



**CLOTRIMAZOLE
TROCHE/LOZENGES**
Tablet **10** mg

LOT XXXXXX EXP. XXXXXX
N 0054-8146-22 Roxane • Cols., OH



**CLOTRIMAZOLE
TROCHE/LOZENGES**
Tablet **10** mg

LOT XXXXXX EXP. XXXXXX
N 0054-8146-22 Roxane • Cols., OH



**CLOTRIMAZOLE
TROCHE/LOZENGES**
Tablet **10** mg

LOT XXXXXX EXP. XXXXXX
N 0054-8146-22 Roxane • Cols., OH



**CLOTRIMAZOLE
TROCHE/LOZENGES**
Tablet **10** mg

LOT XXXXXX EXP. XXXXXX
N 0054-8146-22 Roxane • Cols., OH



**CLOTRIMAZOLE
TROCHE/LOZENGES**
Tablet **10** mg

LOT XXXXXX EXP. XXXXXX
N 0054-8146-22 Roxane • Cols., OH



LOT
EXP.

NDC 0054-8146-22 7 x 10 Unit-Dose
Tablets

CLOTRIMAZOLE TROCHE (clotrimazole lozenges)

10 mg

Each troche contains 10 mg of clotrimazole (1 mg of diphenhydramine HCl) and 10 mg of polyethylene glycol.

POSAGE: Swallow (do not chew) slowly dissolved in the mouth. See accompanying literature for complete information.


This drug is for oral use only. It is not for injection or other use. Use as directed. See accompanying literature for complete information.

For information call 1-800-391-0111.

Store below 86°F (30°C).
Avoid freezing.
Rx only.



Roxane
Laboratories, Inc.
Columbus, Ohio 43216



CLOTRIMAZOLE TROCHE (clotrimazole lozenges)

10 mg

NDC 0054-8146-22 7 x 10 Unit-Dose
Tablets

CLOTRIMAZOLE TROCHE (clotrimazole lozenges)

10 mg

Each troche contains 10 mg of clotrimazole (1 mg of diphenhydramine HCl) and 10 mg of polyethylene glycol.

POSAGE: Swallow (do not chew) slowly dissolved in the mouth. See accompanying literature for complete information.

This drug is for oral use only. It is not for injection or other use. Use as directed. See accompanying literature for complete information.

For information call 1-800-391-0111.

Store below 86°F (30°C).
Avoid freezing.
Rx only.



Roxane
Laboratories, Inc.
Columbus, Ohio 43216

CLOTRIMAZOLE TROCHE (clotrimazole lozenges)

10 mg

REDUCED TO 78%
BY FOIA STAFF