

**CENTER FOR DRUG EVALUATION AND RESEARCH**

**APPROVAL PACKAGE FOR:**

**APPLICATION NUMBER**

**21-332**

**Approved Labeling**

**SYMLIN®**  
**(pramlintide acetate) Injection**

**Rx only**

**WARNING**

SYMLIN is used with insulin and has been associated with an increased risk of insulin-induced severe hypoglycemia, particularly in patients with type 1 diabetes. When severe hypoglycemia associated with SYMLIN use occurs, it is seen within 3 hours following a SYMLIN injection. If severe hypoglycemia occurs while operating a motor vehicle, heavy machinery, or while engaging in other high-risk activities, serious injuries may occur. Appropriate patient selection, careful patient instruction, and insulin dose adjustments are critical elements for reducing this risk.

**DESCRIPTION**

SYMLIN® (pramlintide acetate) Injection is an antihyperglycemic drug for use in patients with diabetes treated with insulin. Pramlintide is a synthetic analog of human amylin, a naturally occurring neuroendocrine hormone synthesized by pancreatic beta cells that contributes to glucose control during the postprandial period. Pramlintide is provided as an acetate salt of the synthetic 37-amino acid polypeptide, which differs in amino acid sequence from human amylin by replacement with proline at positions 25 (alanine), 28 (serine), and 29 (serine).

The structural formula of pramlintide acetate is as shown:

Lys-Cys-Asn-Thr-Ala-Thr-Cys-Ala-Thr-Gln-Arg-Leu-Ala-Asn-Phe-Leu-Val-His-Ser-Ser-Asn-Asn-Phe-Gly-Pro-Ile-Leu-Pro-Pro-Thr-Asn-Val-Gly-Ser-Asn-Thr-Tyr-NH<sub>2</sub> acetate (salt) with a disulfide bridge between the two Cys residues.

Pramlintide acetate is a white powder that has a molecular formula of C<sub>171</sub>H<sub>267</sub>N<sub>51</sub>O<sub>53</sub>S<sub>2</sub>• x C<sub>2</sub>H<sub>4</sub>O<sub>2</sub> (3 ≤ x ≤ 8); the molecular weight is 3949.4. Pramlintide acetate is soluble in water.


SYMLIN is formulated as a clear, isotonic, sterile solution for subcutaneous (SC) administration. SYMLIN vials contain 0.6 mg/mL of pramlintide (as acetate), 2.25 mg/mL of metacresol as a preservative, D-mannitol as a tonicity modifier, and acetic acid and sodium acetate as pH modifiers. SYMLIN has a pH of approximately 4.0.

## CLINICAL PHARMACOLOGY

### Amylin Physiology

Amylin is co-located with insulin in secretory granules and co-secreted with insulin by pancreatic beta cells in response to food intake. Amylin and insulin show similar fasting and postprandial patterns in healthy individuals (Figure 1).

Figure 1: Secretion Profile of Amylin and Insulin in Healthy Adults



Amylin affects the rate of postprandial glucose appearance through a variety of mechanisms. Amylin slows gastric emptying (i.e., the rate at which food is released from the stomach to the small intestine) without altering the overall absorption of nutrients. In addition, amylin suppresses glucagon secretion (not normalized by insulin alone), which leads to suppression of endogenous glucose output from the liver. Amylin also regulates food intake due to centrally-mediated modulation of appetite.

In patients with insulin-using type 2 or type 1 diabetes, the pancreatic beta cells are dysfunctional or damaged, resulting in reduced secretion of both insulin and amylin in response to food.

### Mechanism of Action

SYMLIN, by acting as an amylinomimetic agent, has the following effects: 1) modulation of gastric emptying; 2) prevention of the postprandial rise in plasma glucagon; and 3) satiety leading to decreased caloric intake and potential weight loss.

**Gastric Emptying.** The gastric-emptying rate is an important determinant of the postprandial rise in plasma glucose. SYMLIN slows the rate at which food is released from the stomach to the small intestine following a meal and, thus, it reduces the initial

postprandial increase in plasma glucose. This effect lasts for approximately 3 hours following SYMLIN administration. SYMLIN does not alter the net absorption of ingested carbohydrate or other nutrients.

**Postprandial Glucagon Secretion.** In patients with diabetes, glucagon concentrations are abnormally elevated during the postprandial period, contributing to hyperglycemia. SYMLIN has been shown to decrease postprandial glucagon concentrations in insulin-using patients with diabetes.

**Satiety.** SYMLIN administered prior to a meal has been shown to reduce total caloric intake. This effect appears to be independent of the nausea that can accompany SYMLIN treatment.

### Pharmacokinetics

**Absorption.** The absolute bioavailability of a single SC dose of SYMLIN is approximately 30 to 40%. Subcutaneous administration of different doses of SYMLIN into the abdominal area or thigh of healthy subjects resulted in dose-proportionate maximum plasma concentrations ( $C_{max}$ ) and overall exposure (expressed as area under the plasma concentration curve or (AUC)) (Table 1).

**Table 1: Mean Pharmacokinetic Parameters Following Administration of Single SC Doses of SYMLIN**

SC Dose ( $\mu$ g)	AUC <sub>(0-∞)</sub> (pmol*min/L)	C <sub>max</sub> (pmol/L)	T <sub>max</sub> (min)	Elimination t <sub>1/2</sub> (min)
30	3750	39	21	55
60	6778	79	20	49
90	8507	102	19	51
120	11970	147	21	48

Injection of SYMLIN into the arm showed higher exposure with greater variability, compared with exposure after injection of SYMLIN into the abdominal area or thigh.

There was no strong correlation between the degree of adiposity as assessed by BMI or skin fold thickness measurements and relative bioavailability. Injections administered with 6.0-mm and 12.7-mm needles yielded similar bioavailability.

**Distribution.** SYMLIN does not extensively bind to blood cells or albumin (approximately 40% of the drug is unbound in plasma), and thus SYMLIN's pharmacokinetics should be insensitive to changes in binding sites.

**Metabolism and Elimination.** In healthy subjects, the half-life of SYMLIN is approximately 48 minutes. SYMLIN is metabolized primarily by the kidneys. Des-lys<sup>1</sup> pramlintide (2-37 pramlintide), the primary metabolite, has a similar half-life and is

biologically active both *in vitro* and *in vivo* in rats. AUC values are relatively constant with repeat dosing, indicating no bioaccumulation.

#### **Special Populations.**

**Renal Insufficiency:** Patients with moderate or severe renal impairment ( $Cl_{Cr} >20$  to  $\leq 50$  mL/min) did not show increased SYMLIN exposure or reduced SYMLIN clearance, compared to subjects with normal renal function. No studies have been done in dialysis patients.

**Hepatic Insufficiency:** Pharmacokinetic studies have not been conducted in patients with hepatic insufficiency. However, based on the large degree of renal metabolism (see Metabolism and Elimination), hepatic dysfunction is not expected to affect blood concentrations of SYMLIN.

**Geriatric:** Pharmacokinetic studies have not been conducted in the geriatric population. SYMLIN should only be used in patients known to fully understand and adhere to proper insulin adjustments and glucose monitoring. No consistent age-related differences in the activity of SYMLIN have been observed in the geriatric population (n=539 for patients 65 years of age or older in the clinical trials).

**Pediatric:** SYMLIN has not been evaluated in the pediatric population.

**Gender:** No study has been conducted to evaluate possible gender effects on SYMLIN pharmacokinetics. However, no consistent gender-related differences in the activity of SYMLIN have been observed in the clinical trials (n=2799 for male and n=2085 for female).

**Race/Ethnicity:** No study has been conducted to evaluate the effect of ethnicity on SYMLIN pharmacokinetics. However, no consistent differences in the activity of SYMLIN have been observed among patients of differing race/ethnicity in the clinical trials (n=4257 for white, n=229 for black, n=337 for Hispanic, and n=61 for other ethnic origins).


**Drug Interactions:** The effect of SYMLIN (120  $\mu$ g) on acetaminophen (1000 mg) pharmacokinetics as a marker of gastric-emptying was evaluated in patients with type 2 diabetes (n=24). SYMLIN did not significantly alter the AUC of acetaminophen. However, SYMLIN decreased acetaminophen  $C_{max}$  (about 29% with simultaneous co-administration) and increased the time to maximum plasma concentration or  $t_{max}$  (ranging from 48 to 72 minutes) dependent on the time of acetaminophen administration relative to SYMLIN injection. SYMLIN did not significantly affect acetaminophen  $t_{max}$  when acetaminophen was administered 1 to 2 hours before SYMLIN injection. However, the  $t_{max}$  of acetaminophen was significantly increased when acetaminophen was administered simultaneously with or up to 2 hours following SYMLIN injection (see PRECAUTIONS, Drug Interactions).

## 152 Pharmacodynamics

153 In clinical studies in patients with insulin-using type 2 and type 1 diabetes, SYMLIN  
 154 administration resulted in a reduction in mean postprandial glucose concentrations, reduced  
 155 glucose fluctuations, and reduced food intake. SYMLIN doses differ for insulin-using type 2  
 156 and type 1 patients (see DOSAGE AND ADMINISTRATION).

158 **Reduction in Postprandial Glucose Concentrations.** SYMLIN administered  
 159 subcutaneously immediately prior to a meal reduced plasma glucose concentrations  
 160 following the meal when used with regular insulin or rapid-acting insulin analogs (Figure 2).  
 161 **This reduction in postprandial glucose decreased the amount of short-acting insulin**  
 162 **required and limited glucose fluctuations based upon 24-hour glucose monitoring.**  
 163 When rapid-acting analog insulins were used, plasma glucose concentrations tended to rise  
 164 during the interval between 150 minutes following SYMLIN injection and the next meal (see  
 165 DOSAGE and ADMINISTRATION).

167 **Figure 2: Postprandial Plasma Glucose Profiles in Patients With Type 2 and Type 1**  
 168 **Diabetes Receiving SYMLIN and/or Insulin**



169  
 170

171 **Reduced Food Intake.** A single, subcutaneous dose of SYMLIN 120 µg (type 2) or  
 172 30 µg (type 1) administered 1 hour prior to an unlimited buffet meal was associated with  
 173 reductions in total caloric intake (placebo-subtracted mean changes of ~23% and 21%,  
 174 respectively), which occurred without decreases in meal duration.

175

176

**CLINICAL STUDIES**

177

178

179

180

181

182

A total of 5325 patients and healthy volunteers received SYMLIN in clinical studies. This includes 1688 with type 2 diabetes and 2375 with type 1 diabetes in short- and long-term controlled clinical trials, long-term uncontrolled clinical trials, and an open-label study in the clinical practice setting.

**Clinical Studies in Type 2 Diabetes**

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

The efficacy of a range of SYMLIN doses was evaluated in several placebo-controlled and open-label clinical trials in insulin-using patients with type 2 diabetes. Based on results obtained in these studies, the recommended dose of SYMLIN for patients with insulin-using type 2 diabetes is 120 µg administered immediately prior to major meals.

Two, long-term (26 to 52 week), randomized, double-blind, placebo-controlled studies of SYMLIN were conducted in patients with type 2 diabetes using fixed dose insulin to isolate the SYMLIN effect. Demographic and baseline characteristics for the 871 SYMLIN-treated patients are as follows: mean baseline HbA1c ranged from 9.0 to 9.4%, mean age was 56.4 to 59.1 years, mean duration of diabetes ranged from 11.5 to 14.4 years, and mean BMI ranged from 30.1 to 34.4 kg/m<sup>2</sup>. In both of these studies, SYMLIN or placebo was added to the participants' existing diabetes therapies, which included insulin with or without a sulfonylurea agent and/or metformin.

Table 2 summarizes the composite results across both studies for patients assigned to the 120-µg dose after 6 months of treatment.

**Table 2: Mean (SE) Change in HbA1c, Weight, and Insulin at 6 Months in the Double-Blind, Placebo-Controlled Studies in Patients With Insulin-using Type 2 Diabetes**

Variable	Placebo	SYMLIN (120 µg)
Baseline HbA1c (%)	9.3 (0.08)	9.1 (0.06)
Change in HbA1c at 6 Months Relative to Baseline (%)	-0.17 (0.07)	-0.57 (0.06)*
Placebo-Subtracted HbA1c Change at 6 Months (%)	NA	-0.40 (0.09)*
Baseline Weight (kg)	91.3 (1.2)	92.5 (1.2)
Change in Weight at 6 Months Relative to Baseline (kg)	+0.2 (0.2)	-1.5 (0.2)*
Placebo-Subtracted Weight Change at 6 Months (kg)	NA	-1.7 (0.3)*
Percent Change in Insulin Doses at 6 Months: Rapid/Short-Acting	+6.5 (2.7)	-3.0 (1.6)*
Percent Change in Insulin Doses at 6 Months: Long-Acting	+5.2 (1.4)	-0.2 (1.3)*

\* Statistically significant reduction compared with placebo (p-value < 0.05).

In a cohort of 145 patients who completed two years of SYMLIN treatment the baseline subtracted HbA1c and weight reductions were: -0.40% and -0.36 kg, respectively.

**Open-Label Study in the Clinical Practice Setting.** An open-label study of SYMLIN was conducted at the recommended dose of 120 µg in 166 patients with insulin-using type 2 diabetes who were unable to achieve glycemic targets using insulin alone. A flexible-dose insulin regimen was employed in these patients (see DOSAGE and ADMINISTRATION). In this study, patients adjusted their insulin regimen based on pre-and post-meal glucose monitoring. At baseline, mean HbA1c was 8.3%, mean age was 54.4 years, mean duration of diabetes was 13.3 years, and mean BMI was 38.6 kg/m<sup>2</sup>. SYMLIN was administered with major meals. SYMLIN plus insulin treatment for 6 months resulted in a baseline-subtracted mean HbA1c reduction of -0.56 ± 0.15 % and a baseline-subtracted mean weight reduction of -2.76 ± 0.34 kg. These changes were accomplished with reductions in doses of total, short-acting, and long-acting insulin (-6.4 ± 2.66, -10.3 ± 4.84, and -4.20 ± 2.42 %, respectively).

### Clinical Studies in Type 1 Diabetes

The efficacy of a range of SYMLIN doses was evaluated in several placebo-controlled and open-label clinical trials conducted in patients with type 1 diabetes. Based on results obtained in these studies, the recommended dose of SYMLIN for patients with type 1 diabetes is 30 µg or 60 µg administered immediately prior to major meals.

Three, long-term (26 to 52 week), randomized, double-blind, placebo-controlled studies of SYMLIN were conducted in patients with type 1 diabetes (N=1717). Two of these studies allowed only minimal insulin adjustments in order to isolate the SYMLIN effect; in the third study, insulin adjustments were made according to standard medical practice. Demographic and baseline characteristics for the 1179 SYMLIN-treated patients were as follows: mean baseline HbA1c range was 8.7 to 9.0%, mean age range was 37.3 to 41.9 years, mean duration of diabetes range was 15.5 to 19.2 years, and mean BMI range was 25.0 to 26.8 kg/m<sup>2</sup>. SYMLIN or placebo was added to existing insulin therapies.

**Table 3** summarizes the composite results across these studies for patients assigned to the 30 or 60 µg dose after 6 months of treatment.



**Table 3: Mean (SE) Change in HbA1c, Weight, and Insulin at 6 Months in the Double-Blind, Placebo-Controlled Studies in Patients With Type 1 Diabetes**

Variable	Placebo	SYMLIN (30 or 60 µg)
Baseline HbA1c (%)	9.0 (0.06)	8.9 (0.04)
Change in HbA1c at 6 Months Relative to Baseline (%)	-0.10 (0.05)	-0.43 (0.04)*
Placebo-Subtracted HbA1c Change at 6 Months (%)	NA	-0.33 (0.06)*
Baseline Weight (kg)	75.1 (0.6)	76.1 (0.5)
Change in Weight at 6 Months Relative to Baseline (kg)	+0.6 (0.1)	-1.1 (0.1)*
Placebo-Subtracted Weight Change at 6 Months (kg)	NA	-1.7 (0.1)*
Percent Change in Insulin Doses at 6 Months: Rapid/Short-Acting	+1.7 (3.3)	-3.6 (2.9)
Percent Change in Insulin Doses at 6 Months: Long-Acting	+2.5 (1.9)	+1.9 (1.3)

\* Statistically significant reduction compared with placebo (p-value < 0.05).

In a cohort of 73 patients who completed two years of SYMLIN treatment the baseline subtracted HbA1c and weight changes were: -0.35% and 0.60 kg, respectively.

**SYMLIN Dose-Titration Trial.** A dose-titration study of SYMLIN was conducted in patients with type 1 diabetes. Patients with relatively good baseline glycemic control (mean HbA1c = 8.1%) were randomized to receive either insulin plus placebo or insulin plus SYMLIN. Other baseline and demographics characteristics were: mean age of 41 years, mean duration of diabetes of 20 years, mean BMI of 28 kg/m<sup>2</sup>. SYMLIN was initiated at a dose of 15 µg and titrated upward at weekly intervals by 15-µg increments to doses of 30 µg or 60 µg, based on whether patients experienced nausea. Once a tolerated dose of either 30 µg or 60 µg was reached, the SYMLIN dose was maintained for the remainder of the study (SYMLIN was administered before major meals). During SYMLIN titration, the insulin dose (mostly the short/rapid acting insulin) was reduced by 30-50% in order to reduce the occurrence of hypoglycemia. Once a tolerated SYMLIN dose was reached, insulin dose adjustments were made according to standard clinical practice, based on pre- and post-meal blood glucose monitoring. By 6 months of treatment, patients treated with SYMLIN and insulin and patients treated with insulin and placebo had equivalent reductions in mean HbA1c ( $-0.47 \pm 0.07\%$  vs.  $-0.49 \pm 0.07\%$ , respectively); patients on SYMLIN lost weight ( $-1.33 \pm 0.31$  kg relative to baseline and  $-2.6$  kg relative to placebo plus insulin-treated patients). SYMLIN-treated patients used less total insulin ( $-11.7\%$  relative to baseline) and less short/rapid-acting insulin ( $-22.8\%$  relative to baseline).

**Open-Label Study in the Clinical Practice Setting.** An open-label study of SYMLIN was conducted in patients with type 1 diabetes who were unable to achieve glycemic targets using insulin alone. A flexible-dose insulin regimen was employed in these patients after SYMLIN titration was completed (see DOSAGE and ADMINISTRATION). In this study, patients adjusted their insulin regimen based on pre-and post-meal glucose monitoring. At baseline, mean HbA1c was 8.0%, mean age was 42.7 years, mean duration of diabetes was 21.2 years, and mean BMI was 28.6 kg/m<sup>2</sup>. SYMLIN daily dosage was 30 µg or 60 µg with major meals.

SYMLIN plus insulin reduced HbA1c and body weight from baseline at 6 months by a mean of 0.18% and 3.0 kg, respectively. These changes in glycemic control and body weight were achieved with reductions in doses of total, short-acting, and long-acting insulin ( $-12.0 \pm 1.36$ ,  $-21.7 \pm 2.81$ , and  $-0.4 \pm 1.59$  %, respectively).

## INDICATIONS AND USAGE

SYMLIN is given at mealtimes and is indicated for:

- Type 1 diabetes, as an adjunct treatment in patients who use mealtime insulin therapy and who have failed to achieve desired glucose control despite optimal insulin therapy.
- Type 2 diabetes, as an adjunct treatment in patients who use mealtime insulin therapy and who have failed to achieve desired glucose control despite optimal insulin therapy, with or without a concurrent sulfonylurea agent and/or metformin.

## CONTRAINDICATIONS

SYMLIN is contraindicated in patients with any of the following:

- a known hypersensitivity to SYMLIN or any of its components, including metacresol;
- a confirmed diagnosis of gastroparesis;
- hypoglycemia unawareness.

## WARNINGS

### Patient Selection

**Proper patient selection is critical to safe and effective use of SYMLIN.** Before initiation of therapy, the patient's HbA1c, recent blood glucose monitoring data, history of insulin-induced hypoglycemia, current insulin regimen, and body weight should be reviewed. SYMLIN therapy should only be considered in patients with insulin-using type 2 or type 1 diabetes who fulfill the following criteria:

- have failed to achieve adequate glycemic control despite individualized insulin management;
- are receiving ongoing care under the guidance of a health care professional skilled in the use of insulin and supported by the services of diabetes educator(s).

Patients meeting any of the following criteria should NOT be considered for SYMLIN therapy:

- poor compliance with current insulin regimen;
- poor compliance with prescribed self-blood glucose monitoring;
- have an HbA1c >9%;
- recurrent severe hypoglycemia requiring assistance during the past 6 months;
- presence of hypoglycemia unawareness;
- confirmed diagnosis of gastroparesis;
- require the use of drugs that stimulate gastrointestinal motility;
- pediatric patients.

**Hypoglycemia.** SYMLIN alone does not cause hypoglycemia. However, SYMLIN is indicated to be co-administered with insulin therapy and in this setting SYMLIN increases the risk of insulin-induced severe hypoglycemia, particularly in patients with type 1 diabetes. Severe hypoglycemia associated with SYMLIN occurs within the first 3 hours following a SYMLIN injection. If severe hypoglycemia occurs while operating a motor vehicle, heavy machinery, or while engaging in other high-risk activities, serious injuries may occur. Therefore, when introducing SYMLIN therapy, appropriate precautions need to be taken to avoid increasing the risk for insulin-induced severe hypoglycemia. These precautions include **frequent pre- and post-meal glucose monitoring combined with an initial 50% reduction in pre-meal doses of short-acting insulin (see DOSAGE and ADMINISTRATION).**

Symptoms of hypoglycemia may include hunger, headache, sweating, tremor, irritability, or difficulty concentrating. Rapid reductions in blood glucose concentrations may induce such symptoms regardless of glucose values. More severe symptoms of hypoglycemia include loss of consciousness, coma, or seizure.

Early warning symptoms of hypoglycemia may be different or less pronounced under certain conditions, such as long duration of diabetes; diabetic nerve disease; use of medications such as beta-blockers, clonidine, guanethidine, or reserpine; or intensified diabetes control.

The addition of any antihyperglycemic agent such as SYMLIN to an existing regimen of one or more anti-hyperglycemic agents (e.g., insulin, sulfonylurea), or other agents that can increase the risk of hypoglycemia may necessitate further insulin dose adjustments and particularly close monitoring of blood glucose.

The following are examples of substances that may increase the blood glucose-lowering effect and susceptibility to hypoglycemia: oral anti-diabetic products, ACE inhibitors,

diisopyramide, fibrates, fluoxetine, MAO inhibitors, pentoxifylline, propoxyphene, salicylates, and sulfonamide antibiotics.

Clinical studies employing a controlled hypoglycemic challenge have demonstrated that SYMLIN does not alter the counter-regulatory hormonal response to insulin-induced hypoglycemia. Likewise, in SYMLIN-treated patients, the perception of hypoglycemic symptoms was not altered with plasma glucose concentrations as low as 45 mg/dL.

## **PRECAUTIONS**

### **General:**

Hypoglycemia (See WARNINGS).

**Information for Patients:** Health care providers should inform patients of the potential risks and advantages of SYMLIN therapy. Health care providers should also inform patients about self-management practices including glucose monitoring, proper injection technique, timing of dosing, and proper storage of SYMLIN. In addition, the importance of adherence to meal planning, physical activity, recognition and management of hypoglycemia and hyperglycemia, and assessment of diabetes complications. Refer patients to the SYMLIN Medication Guide for additional information.

Instruct patients on handling of special situations such as intercurrent conditions (illness or stress), an inadequate or omitted insulin dose, inadvertent administration of increased insulin or SYMLIN dose, inadequate food intake or missed meals.

**SYMLIN and insulin should always be administered as separate injections and never be mixed.**

Women with diabetes should be advised to inform their healthcare professional if they are pregnant or contemplating pregnancy.

**Renal Impairment:** The dosing requirements for SYMLIN are not altered in patients with moderate or severe renal impairment ( $Cl_{Cr} >20$  to  $\leq 50$  mL/min). No studies have been done in dialysis patients (see CLINICAL PHARMACOLOGY; Special Populations).

**Hepatic Impairment:** Studies have not been performed in patients with hepatic impairment. However, hepatic dysfunction is not expected to affect blood concentrations of SYMLIN (see CLINICAL PHARMACOLOGY; Special Populations).

**Allergy: Local allergy.** Patients may experience redness, swelling, or itching at the site of injection. These minor reactions usually resolve in a few days to a few weeks. In some instances, these reactions may be related to factors other than SYMLIN, such as irritants in a skin cleansing agent or improper injection technique.

**Systemic Allergy.** In controlled clinical trials up to 12 months, potential systemic allergic reactions were reported in 65 (5)% of type 2 patients and 59 (5%) of type 1 SYMLIN-treated patients. Similar reactions were reported by 18 (4%) and 28 (5%) of placebo-treated type 2 and type 1 patients, respectively. No patient receiving SYMLIN was withdrawn from a trial due to a potential systemic allergic reaction.

#### **Drug Interactions**

Due to its effects on gastric emptying, SYMLIN therapy should not be considered for patients taking drugs that alter gastrointestinal motility (e.g., anticholinergic agents such as atropine) and agents that slow the intestinal absorption of nutrients (e.g.,  $\alpha$ -glucosidase inhibitors). Patients using these drugs have not been studied in clinical trials.

SYMLIN has the potential to delay the absorption of concomitantly administered oral medications. When the rapid onset of a concomitant orally administered agent is a critical determinant of effectiveness (such as analgesics), the agent should be administered at least 1 hour prior to or 2 hours after SYMLIN injection.

In clinical trials, the concomitant use of sulfonylureas or biguanides did not alter the adverse event profile of SYMLIN. No formal interaction studies have been performed to assess the effect of SYMLIN on the kinetics of oral antidiabetic agents.

#### **Mixing SYMLIN and Insulin**

The pharmacokinetic parameters of SYMLIN were altered when mixed with regular, NPH, and 70/30 premixed formulations of recombinant human insulin immediately prior to injection. **Thus, SYMLIN and insulin should not be mixed and must be administered separately.**

#### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

**Carcinogenesis.** A two-year carcinogenicity study was conducted in CD-1 mice with doses of 0.2, 0.5, and 1.2 mg/kg/day of SYMLIN (32, 67, and 159 times the exposure resulting from the maximum recommended human dose based on area under the plasma concentration curve or AUC, respectively). No drug-induced tumors were observed. A two-year carcinogenicity study was conducted in Sprague-Dawley rats with doses of 0.04, 0.2, and 0.5 mg/kg/day of SYMLIN (3, 9, and 25 times the exposure resulting from the maximum recommended human dose based on AUC, respectively). No drug-induced tumors were observed in any organ.

**Mutagenesis.** SYMLIN was not mutagenic in the Ames test and did not increase chromosomal aberration in the human lymphocytes assay. SYMLIN was not clastogenic in the *in vivo* mouse micronucleus test or in the chromosomal aberration assay utilizing Chinese hamster ovary cells.

**Impairment of Fertility.** Administration of 0.3, 1, or 3 mg/kg/day of SYMLIN (8, 17, and 82 times the exposure resulting from the maximum recommended human dose based on body surface area) had no significant effects on fertility in male or female rats. The

highest dose of 3 mg/kg/day resulted in dystocia in 8/12 female rats secondary to significant decreases in serum calcium levels.

### **Pregnancy**

**Teratogenic Effects: Pregnancy Category C.** No adequate and well-controlled studies have been conducted in pregnant women. Studies in perfused human placenta indicate that SYMLIN has low potential to cross the maternal/fetal placental barrier. Embryofetal toxicity studies with SYMLIN have been performed in rats and rabbits. Increases in congenital abnormalities (neural tube defect, cleft palate, exencephaly) were observed in fetuses of rats treated during organogenesis with 0.3 and 1.0 mg/kg/day (10 and 47 times the exposure resulting from the maximum recommended human dose based on AUC, respectively). Administration of doses up to 0.3 mg/kg/day SYMLIN (9 times maximum recommended dose based on AUC) to pregnant rabbits had no adverse effects in embryofetal development; however, animal reproduction studies are not always predictive of human response. SYMLIN should be used during pregnancy only if it is determined by the healthcare professional that the potential benefit justifies the potential risk to the fetus.

### **Nursing Mothers**

It is unknown whether SYMLIN is excreted in human milk. Many drugs, including peptide drugs, are excreted in human milk. Therefore, SYMLIN should be administered to nursing women only if it is determined by the health care professional that the potential benefit outweighs the potential risk to the infant.

### **Pediatric Use**

Safety and effectiveness of SYMLIN in pediatric patients have not been established.

### **Geriatric Use**

SYMLIN has been studied in patients ranging in age from 15 to 84 years of age, including 539 patients 65 years of age or older. The change in HbA1c values and hypoglycemia frequencies did not differ by age, but greater sensitivity in some older individuals cannot be ruled out. Thus, both SYMLIN and insulin regimens should be carefully managed to obviate an increased risk of severe hypoglycemia.

## **ADVERSE REACTIONS**

Adverse events (excluding hypoglycemia, discussed below) commonly associated with SYMLIN when co-administered with a fixed dose of insulin in the long-term, placebo-controlled trials in insulin-using type 2 patients and type 1 patients are presented in Table 4 and Table 5, respectively. The same adverse events were also shown in the open-label clinical practice study, which employed flexible insulin dosing.

**Table 4: Treatment-Emergent Adverse Events Occurring With  $\geq 5\%$  Incidence and Greater Incidence With SYMLIN Compared With Placebo in Long-Term,**

**Placebo-Controlled Trials. Incidence of the Same Events in the Open-Label Clinical Practice Study (Patients With Insulin-Using Type 2 Diabetes, 120 µg)**

	Long-Term, Placebo-Controlled Studies		Open-Label, Clinical Practice Study
	Placebo + Insulin (n(%)) (N=284)	SYMLIN + Insulin (n(%)) (N=292)	SYMLIN + Insulin (n(%)) (N=166)
Nausea	34 (12)	81 (28)	53 (30)
Headache	19 (7)	39 (13)	8 (5)
Anorexia	5 (2)	27 (9)	1 (<1)
Vomiting	12 (4)	24 (8)	13 (7)
Abdominal Pain	19 (7)	23 (8)	3 (2)
Fatigue	11 (4)	20 (7)	5 (3)
Dizziness	11 (4)	17 (6)	3 (2)
Coughing	12 (4)	18 (6)	4 (2)
Pharyngitis	7 (2)	15 (5)	6 (3)

**Table 5: Treatment-Emergent Adverse Events Occurring With ≥5% Incidence and Greater Incidence with SYMLIN Compared to Placebo in Long-Term, Placebo-Controlled Studies. Incidence of the Same Events in the Open-Label Clinical Practice Study (Patients With Type 1 Diabetes, 30 or 60 µg)**

	Long-Term, Placebo-Controlled Studies		Open-Label, Clinical Practice Study
	Placebo + Insulin (n(%)) (N=538)	SYMLIN + Insulin (n(%)) (N=716)	SYMLIN + Insulin (n(%)) (N=265)
Nausea	92 (17)	342 (48)	98 (37)
Anorexia	12 (2)	122 (17)	0 (0)
Inflicted Injury	55 (10)	97 (14)	20 (8)
Vomiting	36 (7)	82 (11)	18 (7)
Arthralgia	27 (5)	51 (7)	6 (2)
Fatigue	22 (4)	51 (7)	12 (4.5)
Allergic Reaction	28 (5)	41 (6)	1 (<1)
Dizziness	21 (4)	34 (5)	5 (2)

Most adverse events were gastrointestinal in nature. In patients with type 2 or type 1 diabetes, the incidence of nausea was higher at the beginning of SYMLIN treatment and decreased with time in most patients. The incidence and severity of nausea are reduced when SYMLIN is gradually titrated to the recommended doses (see DOSAGE and ADMINISTRATION).

### **Severe Hypoglycemia**

SYMLIN alone (without the concomitant administration of insulin) does not cause hypoglycemia. However, SYMLIN is indicated as an adjunct treatment in patients who use mealtime insulin therapy and co-administration of SYMLIN with insulin can increase the risk of insulin-induced hypoglycemia, particularly in patients with type 1 diabetes

(see Boxed Warning). The incidence of severe hypoglycemia during the SYMLIN clinical development program is summarized in Table 6 and Table 7.

**Table 6: Incidence and Event Rate of Severe Hypoglycemia in Long-Term, Placebo-Controlled and Open-Label, Clinical Practice Studies in Patients With Insulin-Using Type 2 Diabetes**

	Long-Term, Placebo-Controlled Studies (No Insulin Dose-Reduction During Initiation)				Open-Label, Clinical Practice Study (Insulin Dose-Reduction During Initiation)	
	Placebo + Insulin		SYMLIN + Insulin		SYMLIN + Insulin	
	0-3 Months (n=284)	>3-6 Months (n=251)	0-3 Months (n=292)	>3-6 Months (n=255)	0-3 Months (n=166)	>3-6 Months (n=150)
<b>Severe Hypoglycemia</b>						
<b>Patient-Ascertained*</b>						
Event Rate (event rate/patient year)	0.24	0.13	0.45	0.39	0.05	0.03
Incidence (%)	2.1	2.4	8.2	4.7	0.6	0.7
<b>Medically Assisted**</b>						
Event Rate (event rate/patient year)	0.06	0.07	0.09	0.02	0.05	0.03
Incidence (%)	0.7	1.2	1.7	0.4	0.6	0.7

\* **Patient-ascertained severe hypoglycemia:** Requiring the assistance of another individual (including aid in ingestion of oral carbohydrate); and/or requiring the administration of glucagon injection, intravenous glucose, or other medical intervention.

\*\* **Medically assisted severe hypoglycemia:** Requiring glucagon, IV glucose, hospitalization, paramedic assistance, emergency room visit, and/or assessed as an SAE by the investigator.

**Table 7: Incidence and Event Rate of Severe Hypoglycemia in Long-Term, Placebo-Controlled and Open-Label, Clinical Practice Studies in Patients With Type 1 Diabetes**

	Long-Term, Placebo-Controlled Studies (No Insulin Dose-Reduction During Initiation)				Open-Label, Clinical Practice Study (Insulin Dose-Reduction During Initiation)	
	Placebo + Insulin		SYMLIN + Insulin		SYMLIN + Insulin	
	0-3 Months (n=538)	>3-6 Months (n=470)	0-3 Months (n=716)	>3-6 Months (n=576)	0-3 Months (n=265)	>3-6 Months (n=213)
<b>Severe Hypoglycemia</b>						
<b>Patient-Ascertained*</b>						
Event Rate (event rate/patient year)	1.33	1.06	1.55	0.82	0.29	0.16
Incidence (%)	10.8	8.7	16.8	11.1	5.7	3.8
<b>Medically Assisted**</b>						
Event Rate (event rate/patient year)	0.19	0.24	0.50	0.27	0.10	0.04
Incidence (%)	3.3	4.3	7.3	5.2	2.3	0.9

\* **Patient-ascertained severe hypoglycemia:** Requiring the assistance of another individual (including aid in ingestion of oral carbohydrate); and/or requiring the administration of glucagon injection, intravenous glucose, or other medical intervention.

\*\* **Medically assisted severe hypoglycemia:** Requiring glucagon, IV glucose, hospitalization, paramedic assistance, emergency room visit, and/or assessed as an SAE by the investigator.



**534 OVERDOSAGE**

535 Single 10 mg doses of SYMLIN (83 times the maximum dose of 120 µg) were  
536 administered to three healthy volunteers. Severe nausea was reported in all three  
537 individuals and was associated with vomiting, diarrhea, vasodilatation, and dizziness. No  
538 hypoglycemia was reported. SYMLIN has a short half-life and in the case of overdose,  
539 supportive measures are indicated.

**541 DOSAGE AND ADMINISTRATION**

542 SYMLIN dosage differs depending on whether the patient has type 2 or type 1  
543 diabetes (see below). When initiating therapy with SYMLIN, initial insulin dose  
544 reduction is required in all patients (both type 2 and type 1) to reduce the risk of  
545 insulin-induced hypoglycemia. As this reduction in insulin can lead to glucose  
546 elevations, patients should be monitored at regular intervals to assess SYMLIN  
547 tolerability and the effect on blood glucose, so that **individualized** insulin adjustments  
548 can be initiated. If SYMLIN therapy is discontinued for any reason (e.g., surgery or  
549 illnesses), the same initiation protocol should be followed when SYMLIN therapy is  
550 re-instituted (see below).

**552 Initiation of SYMLIN therapy****553 Patients With Insulin-using Type 2 Diabetes**

554 In patients with insulin-using type 2 diabetes, SYMLIN should be initiated at a dose  
555 of 60 µg and increased to a dose of 120 µg as tolerated.

556 Patients should be instructed to:

- 557 • Initiate SYMLIN at 60 µg subcutaneously, immediately prior to major meals;
- 558 • Reduce preprandial, rapid-acting or short-acting insulin dosages, including fixed-mix  
559 insulins (70/30) by 50%;
- 560 • Monitor blood glucose frequently, including pre- and post-meals and at bedtime;
- 561 • Increase the SYMLIN dose to 120 µg when no clinically significant nausea has  
562 occurred for 3-7 days. **SYMLIN dose adjustments should be made only as**  
563 **directed by the health care professional.** If significant nausea persists at the 120 µg  
564 dose, the SYMLIN dose should be decreased to 60 µg;
- 565 • Adjust insulin doses to optimize glycemic control once the target dose of SYMLIN is  
566 achieved and nausea (if experienced) has subsided. **Insulin dose adjustments**  
567 **should be made only as directed by the health care professional;**
- 568 • Contact a health care professional skilled in the use of insulin to review SYMLIN and  
569 insulin dose adjustments at least once a week until a target dose of SYMLIN is  
570 achieved, SYMLIN is well-tolerated, and blood glucose concentrations are stable.

**572 Patients With Type 1 Diabetes**

573 In patients with type 1 diabetes, SYMLIN should be initiated at a dose of 15 µg and  
574 titrated at 15-µg increments to a maintenance dose of 30 µg or 60 µg as tolerated.

575 Patients should be instructed to:

- Initiate SYMLIN at a starting dose of 15 µg subcutaneously, immediately prior to major meals;
- Reduce preprandial, rapid-acting or short-acting insulin dosages, including fixed-mix insulins (e.g., 70/30) by 50%;
- Monitor blood glucose frequently, including pre- and post-meals and at bedtime;
- Increase the SYMLIN dose to the next increment (30 µg, 45 µg, or 60 µg) when no clinically significant nausea has occurred for at least 3 days. **SYMLIN dose adjustments should be made only as directed by the health care professional.** If significant nausea persists at the 45- or 60-µg dose level, the SYMLIN dose should be decreased to 30 µg. If the 30-µg dose is not tolerated, discontinuation of SYMLIN therapy should be considered;
- Adjust insulin doses to optimize glycemic control once the target dose of SYMLIN is achieved and nausea (if experienced) has subsided. **Insulin dose adjustments should be made only as directed by the health care professional;**
- Contact a health care professional skilled in the use of insulin to review SYMLIN and insulin dose adjustments at least once a week until a target dose of SYMLIN is achieved, SYMLIN is well-tolerated, and blood glucose concentrations are stable.

#### **Once Target Dose of SYMLIN Is Achieved in Type 2 or Type 1 Patients**

After a maintenance dose of SYMLIN is achieved, both insulin-using patients with type 2 diabetes and patients with type 1 diabetes should be instructed to:

- Adjust insulin doses to optimize glycemic control once the target dose of SYMLIN is achieved and nausea (if experienced) has subsided. **Insulin dose adjustments should be made only as directed by a health care professional;**
- Contact a health care professional in the event of recurrent nausea or hypoglycemia. An increased frequency of mild to moderate hypoglycemia should be viewed as a warning sign of increased risk for severe hypoglycemia.

#### **Administration**

SYMLIN should be administered subcutaneously immediately prior to each major meal (≥250 kcal or containing ≥30 g of carbohydrate).

To administer SYMLIN from vials, use a U-100 insulin syringe (preferably a 0.3 mL [0.3 cc] size) for optimal accuracy. If using a syringe calibrated for use with U-100 insulin, use the chart below (**Table 8**) to measure the microgram dosage in unit increments. **Do not mix SYMLIN with insulin.**

614 **Table 8: Conversion of SYMLIN Dose to Insulin Unit Equivalents**  
615

Dosage Prescribed (µg)	Increment Using a U-100 Syringe (Units)	Volume (cc or mL)
15	2.5	0.025
30	5.0	0.050
45	7.5	0.075
60	10.0	0.100
120	20.0	0.200

616

617 Each SYMLIN dose should be administered subcutaneously into the abdomen or thigh.

618 Injection sites should be rotated so that the same site is not used repeatedly. The

619 injection site selected should also be distinct from the site chosen for any concomitant

620 insulin injection. Each SYMLIN injection should be administered subcutaneously to

621 abdomen or thigh (administered into the arm is not recommended because of variable  
622 absorption).

623

- 624 • SYMLIN and insulin should always be administered as separate injections.
- 625 • SYMLIN should not be mixed with any type of insulin.
- 626 • If a SYMLIN dose is missed, do not give an additional injection.
- 627 • Always use a new syringe and needle to give SYMLIN and insulin injections.

628

629 **Discontinuation of Therapy**

630 SYMLIN therapy should be discontinued if any of the following occur:

- 631 • Recurrent unexplained hypoglycemia that requires medical assistance;
- 632 • Persistent clinically significant nausea;
- 633 • Noncompliance with self-monitoring of blood glucose concentrations;
- 634 • Noncompliance with insulin dose adjustments;
- 635 • Noncompliance with scheduled health care professional contacts or recommended  
636 clinic visits.

637

638 **Preparation and Handling**639 SYMLIN should be inspected visually for particulate matter or discoloration prior to  
640 administration whenever the solution and the container permit.

641

642 **HOW SUPPLIED**

643 SYMLIN is supplied as a sterile injection in 5 mL vials, containing 0.6 mg/mL  
644 pramlintide (as acetate), for use with a syringe. To administer SYMLIN from vials, use a  
645 U-100 insulin syringe (preferably a 0.3 mL [0.3 cc] size). If using a syringe calibrated  
646 for use with U-100 insulin, use the chart (Table 8) in the DOSAGE AND  
647 ADMINISTRATION section to measure the microgram dosage in unit increments. **Do**  
648 **not mix SYMLIN with insulin.**

649

650 SYMLIN Injection is available in the following package size:  
651 5 mL vials (NDC 66780-110)

652

653 **STORAGE**

654 **Unopened (not in-use) Vials:** Before use, SYMLIN vials should be refrigerated, 36°F to  
655 46°F (2°C to 8°C), and protected from light. Do not freeze. If a vial has been frozen or  
656 overheated, throw it away.

657

658 **Opened (in-use) Vials:** Opened vials in use (punctured) can be kept either refrigerated  
659 or at room temperature for up to 28 days as long as the temperature is not greater than  
660 77°F (25°C). Opened vials, whether or not refrigerated, must be used within 28 days.  
661 Discard after 28 days.

662

663 Storage conditions and stability are summarized in **Table 9:**

664

665 **Table 9: Storage Conditions and Stability of SYMLIN (5 mL Vials)**

666

	Unopened (not in-use) Refrigerated	Open (in-use) Refrigerated or Room Temperature
5 mL Vial	Until Expiration Date	Use Within 28 days

667

668

669 Vials are manufactured for:  
670 Amylin Pharmaceuticals, Inc.  
671 San Diego, CA 92121 USA  
672 1-800-349-8919  
673 <http://www.symlin.com>

674

675 **Rx only**

676

677

## Medication Guide

### **SYMLIN® (SĪM-lĭn) (pramlintide acetate) Injection**

Read the Medication Guide that comes with SYMLIN before you start using it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your doctor about your medical condition or treatment.

#### **What is the most important information I should know about SYMLIN?**

- SYMLIN is used with insulin to lower blood sugar, especially high blood sugar that happens after meals.
- SYMLIN is given at mealtimes. The use of SYMLIN does not replace your daily insulin but may lower the amount of insulin you need, especially before meals.
- Even when SYMLIN is carefully added to your mealtime insulin therapy, your blood sugar may drop too low, especially if you have type 1 diabetes. If this low blood sugar (severe hypoglycemia) happens, it is seen within 3 hours after a SYMLIN injection. Severe low blood sugar makes it hard to think clearly, drive a car, use heavy machinery or do other risky activities where you could hurt yourself or others.
- SYMLIN should only be used by people with type 1 and type 2 diabetes who:
  - already use their insulin as prescribed, but still need better blood sugar control.
  - will follow their doctor's instructions exactly.
  - will follow up with their doctor often.
  - will test their blood sugar levels before and after every meal, and at bedtime.
  - understand how to adjust SYMLIN and insulin doses.

#### **What is SYMLIN?**

SYMLIN is an injectable medicine for adults with type 1 and type 2 diabetes to control blood sugar. SYMLIN slows down the movement of food through your stomach. This affects how fast sugar enters your blood after eating. SYMLIN is always used with insulin to help lower blood sugar during the 3 hours after meals.

#### **Who should not use SYMLIN?**

##### **Do not use SYMLIN if you:**

- cannot tell when your blood sugar is low (hypoglycemia unawareness).
- have a stomach problem called gastroparesis. This is when your stomach does not empty as fast as it should.
- are allergic to SYMLIN or any ingredients in SYMLIN. See the end of this Medication Guide for a complete list of ingredients.

37 SYMLIN has not been studied in children.

38 **What should I tell my doctor before starting SYMLIN?**

39 **Tell your doctor about all of your medical conditions including if you:**

- 40 • **are pregnant or planning to become pregnant.** It is not known if SYMLIN can harm your  
41 unborn baby. You and your doctor will decide how to best control your blood sugar levels  
42 during pregnancy.
- 43 • **are breastfeeding.** It is not known if SYMLIN passes into your milk and if it can harm your  
44 baby. You and your doctor will decide the best way to feed your baby if you are using  
45 SYMLIN.

46 **Keep a list of all the medicines you take. Tell your doctor about all the medicines you take**  
47 **including prescription and non-prescription medicines, vitamins, and herbal supplements.**  
48 SYMLIN can slow down how other medicines pass through your stomach and may affect how  
49 much of them get into your body. Therefore, you may have to change the times you take certain  
50 medicines.

51 **How should I use SYMLIN?**

- 52 • **You must use SYMLIN exactly as prescribed. The amount of SYMLIN you use will**  
53 **depend on whether you have type 1 or type 2 diabetes.** You and your doctor will decide if  
54 you can use SYMLIN.
- 55 • **Never mix SYMLIN and insulin.** You must use different syringes for SYMLIN and insulin  
56 because insulin can affect SYMLIN when the two are mixed together.
- 57 • **Injecting SYMLIN is similar to injecting insulin. Inject SYMLIN under the skin**  
58 **(subcutaneously) of your stomach area (abdomen) or upper leg (thigh).** Inject SYMLIN  
59 at a site that is more than 2 inches away from your insulin injection. Allow SYMLIN to  
60 warm to room temperature before injecting. Use a U-100 insulin syringe (best to use 0.3 mL  
61 [0.3 cc] size) to draw-up and inject SYMLIN. Always use a new syringe and needle for each  
62 SYMLIN injection.
- 63 • **The dose of SYMLIN that your doctor prescribes should be one in the table below. Use this**  
64 **table to match your SYMLIN dose to insulin syringe units:**

<b>Find Your Dose in micrograms (µg)</b>	<b>Draw Up This Amount in U-100 Insulin Syringe (units)</b>
15	2.5
30	5.0
45	7.5
60	10.0
120	20.0

65

- 66 • **Do not use SYMLIN if the liquid in the vial looks cloudy.**
- 67 • **If you take more than your prescribed dose of SYMLIN, you may get nauseous or vomit, and**  
68 **you may not be able to eat the amount of food you usually eat. Pay careful attention to the**

amount of insulin you use at this time as you may be at more risk for low blood sugar. Contact your doctor for guidance.

- If you miss or forget a dose of SYMLIN, wait until the next meal and take your usual dose of SYMLIN at that meal. Do not take more than your usual dose of SYMLIN.

#### Using SYMLIN and insulin with Type 2 Diabetes (see the Table above)

1. Start SYMLIN at 60 µg injected under your skin, just before major meals. A major meal must have at least 250 calories or 30 grams of carbohydrate.
2. Reduce your rapid-acting or short-acting insulin doses before meals by **50 percent**, including fixed-mix insulins such as 70/30. This means half of the dose you usually use.
3. You must check your blood sugar before and after every meal and at bedtime.
4. Increase your dose of SYMLIN to 120 µg on your doctor's instructions if you have not had any nausea for 3 days or more.
5. Tell your doctor right away if you have nausea with the 120 µg dose. Your doctor will tell you how to adjust your dose of SYMLIN.
6. Your doctor may make changes to your insulin doses to better control your blood sugar once you are using the 120 µg dose of SYMLIN. All insulin changes should be directed by your doctor.

#### Using SYMLIN and insulin with Type 1 Diabetes (see the Table above)

1. Start SYMLIN at 15 µg injected under your skin, just before major meals. A major meal must have at least 250 calories or 30 grams of carbohydrate.
2. When starting SYMLIN, reduce your rapid-acting or short-acting insulin doses before meals by **50 percent**, including fixed-mix insulins such as 70/30. This means half of the dose you usually use. All insulin changes should be directed by your doctor.
3. You must check your blood sugar before and after every meal and at bedtime.
4. Increase your dose of SYMLIN to 30 µg on your doctor's instructions if you have not had any nausea for 3 days or more. If you have nausea with SYMLIN at 30 µg, call your doctor right away. Your doctor may decide that you should stop SYMLIN.
5. Increase your dose of SYMLIN to 45 µg on your doctor's instructions if you have not had any nausea for 3 days or more while using the 30 µg dose.
6. Increase your dose of SYMLIN to 60 µg on your doctor's instructions if you have not had any nausea for 3 days or more while using the 45 µg dose.
7. Call your doctor right away if you are bothered with nausea on the 45 µg or 60 µg dose. Your doctor may decide that you should reduce SYMLIN to the 30 µg dose.
8. Your doctor may make changes to your insulin doses to better control your blood sugar once you are on a dose of SYMLIN that is right for you. All insulin changes should be directed by your doctor.

105 **Staying on SYMLIN**

- 106 • Once you reach your recommended dose of SYMLIN, talk to your doctor about changing  
107 your insulin doses to better control your blood sugar. You may have to increase your  
108 long-acting insulin to prevent high blood sugar (hyperglycemia) between meals. **Insulin**  
109 **changes should be directed by your doctor based on blood sugar testing.**
- 110 • Call your doctor if nausea or low blood sugar continues while on your recommended dose of  
111 SYMLIN. Low blood sugar that happens often is a warning sign of possible severe low  
112 blood sugar, especially if you have type 1 diabetes.
- 113 • **If you stop taking SYMLIN for any reason, such as surgery or illness, call your doctor.**  
114 **SYMLIN should be restarted as described above in “How should I use SYMLIN?”**

115 **When should I not use SYMLIN?**

116 **Do not use SYMLIN if:**

- 117 • Your blood sugar is too low.
- 118 • You do not plan to eat. Do not inject SYMLIN if you skip a meal.
- 119 • You plan to eat a meal with less than 250 calories or 30 grams of carbohydrate.
- 120 • You are sick and can't eat your usual meal.
- 121 • You are having surgery or a medical test where you cannot eat.
- 122 • You are pregnant or breastfeeding and have not talked to your doctor.
- 123 Talk to your doctor if you have any of these conditions.

124 **What should I avoid while taking SYMLIN?**

- 125 • Do not drive or operate dangerous machinery until you know how SYMLIN affects your  
126 blood sugar. Low blood sugar makes it hard to think clearly, drive a car, use heavy  
127 machinery or do other risky activities where you could hurt yourself or others. Discuss with  
128 your doctor what activities you should avoid.
- 129 • Alcohol may increase the risk of low blood sugar.
- 130 • **Your doctor will tell you which medicines you can take while using SYMLIN. Do not**  
131 **take other medicines that slow stomach emptying.**
- 132 Always have fast-acting sugar (such as hard candy, glucose tablets, juice) or glucagon available  
133 to treat low blood sugar.

134 **What are the possible side effects of SYMLIN?**

135 **Low blood sugar (hypoglycemia)**

- 136 • SYMLIN is used with insulin to lower your blood sugar, but your blood sugar may  
137 drop too low, especially if you have type 1 diabetes. See “What is the most important  
138 information I should know about SYMLIN?”



- When starting SYMLIN, reduce your doses of insulin before meals as recommended by your doctor to reduce the chance of low blood sugar. You and your doctor should talk about a plan to treat low blood sugar. You should have fast-acting sugar (such as hard candy, glucose tablets, juice) or glucagon with you at all times. Call your doctor if you have low blood sugar more often than normal or severe low blood sugar.

**Your chance for low blood sugar is higher if you:**

- do not reduce your insulin dose before meals at the beginning of SYMLIN treatment, as directed by your doctor.
- use more SYMLIN or insulin than prescribed by your doctor.
- change your insulin dose without checking your blood sugar.
- eat less food than your usual meal.
- are sick and cannot eat.
- are more active than usual.
- have a low blood sugar level before eating.
- drink alcohol.

**Nausea:** Nausea is the most common side effect with SYMLIN. Mild nausea is more likely during the first weeks after starting SYMLIN and usually does not last long. It is very important to start SYMLIN at a low dose and increase it as directed by your doctor. See "How should I use SYMLIN?" If nausea continues or bothers you, call your doctor right away.

**Other Side Effects:** SYMLIN also may cause the following side effects: decreased appetite, vomiting, stomach pain, tiredness, dizziness, or indigestion.

SYMLIN also can cause reactions at the injection site including redness, minor bruising, or pain. Follow the directions under "How should I use SYMLIN?" to reduce the chance of an injection site reaction.

Tell your doctor if you have any side effects that bother you or that do not go away.

These are not all the side effects with SYMLIN. Ask your doctor or pharmacist for more information.

**How should I store SYMLIN?**

- Store SYMLIN vials in the refrigerator until you open them.
- Opened vials can be refrigerated or kept at room temperature for up to 28 days. Any opened vial should be thrown away after 28 days, even if it still has medicine in it.
- Throw away any vial that is out-of-date, has been frozen, heated above room temperature (77°F/25°C) or left at room temperature for more than 28 days.
- **Keep SYMLIN and all medicines out of the reach of children.**

**General information about the safe and effective use of SYMLIN**

Medicines are sometimes prescribed for conditions other than those described in a Medication Guide. Do not use SYMLIN for a condition for which it was not prescribed. Do not give SYMLIN to other people, even if they have the same symptoms that you have. It may harm them.

This Medication Guide summarizes the most important information about SYMLIN. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about SYMLIN that is written for health professionals.

More information on SYMLIN can be found at <http://www.symmlin.com>.

SYMLIN Customer Service is available 24 hours a day at 1-800-349-8919.

**What are the ingredients in SYMLIN?**

**Active ingredient:** pramlintide acetate

**Inactive ingredients:** metacresol, D-mannitol, acetic acid, and sodium acetate.

---

*This draft Medication Guide is pending approval by the U.S. Food and Drug Administration.*

Literature issued March 2005.

Manufactured for Amylin Pharmaceuticals, Inc.  
San Diego CA 92121, USA  
1-800-349-8919  
<http://www.symmlin.com>

Copyright © 2005, Amylin Pharmaceuticals, Inc. All rights reserved.