

Patient Guide to ((REVLIMID REMS[®] logo)) Risk Evaluation And Mitigation Strategy (REMS) Program

This guide provides you important information about:

- The risks of REVLIMID[®] (lenalidomide)
 - Birth defects (deformed babies) or death of an unborn baby
 - Low white blood cells (neutropenia) and low platelets (thrombocytopenia)
 - Blood clots in your arteries (heart attacks and strokes), veins (deep vein thrombosis) and lungs (pulmonary embolism). To reduce this increased risk, most people who take REVLIMID will also take a blood thinner medicine

- The REVLIMID REMS[®] Program
 - What females who can get pregnant need to know:
 - Birth control options
 - What females who can not get pregnant need to know
 - What males need to know

((REVLIMID logo))

Table of contents

Risks of REVLIMID [®] (lenalidomide).....	3
What is the REVLIMID REMS [®] program?	3
How to receive your first prescription for REVLIMID	4
What do all patients need to know about the REVLIMID REMS [®] program?.....	6
What do females who can get pregnant need to know about the REVLIMID REMS [®] program?.....	7
What do females who can not get pregnant need to know about the REVLIMID REMS [®] program?.....	12
What do males need to know about the REVLIMID REMS [®] program?.....	14
Mandatory confidential patient surveys.....	16
Warning to patients taking REVLIMID.....	18

Risks of REVLIMID® (lenalidomide)

REVLIMID is similar to the medicine thalidomide (THALOMID®). Thalidomide can cause severe life-threatening birth defects. If REVLIMID is used during pregnancy, it can cause birth defects or death to unborn babies. REVLIMID must not be used by pregnant females and females who are able to get pregnant. Females who are able to get pregnant must avoid pregnancy while taking REVLIMID.

REVLIMID causes low white blood cells (neutropenia) and low platelets (thrombocytopenia) in most patients.

REVLIMID causes a higher chance for blood clots in your arteries (heart attacks and strokes), veins (deep vein thrombosis) and lungs (pulmonary embolism). To reduce this increased risk, most people who take REVLIMID will also take a blood thinner medicine.

What is the REVLIMID REMS® program?

To avoid serious risks to unborn babies, REVLIMID is only available under a restricted distribution program called the “REVLIMID Risk Evaluation and Mitigation Strategy (REMS)” (formerly known as the RevAssist® program). Only certified prescribers can prescribe REVLIMID and only certified pharmacies can dispense REVLIMID. In order to receive REVLIMID, patients must be enrolled in the REVLIMID REMS® program and agree to follow the requirements.

For more information about REVLIMID and the REVLIMID REMS® program, please visit www.CelgeneRiskManagement.com, or call the Celgene Customer Care Center toll-free at **1-888-423-5436**.

How to receive your first prescription for REVLIMID® (lenalidomide)

For Females:

Counseling Your healthcare provider will counsel you on why and how you and your partner should prevent pregnancy. Your healthcare provider will also inform you on not to share the drug, not to donate blood, and about appropriate contraceptive use. You should be instructed not to extensively handle or open REVLIMID capsules
Pregnancy Test #1 If you can get pregnant, you must take a pregnancy test within 10-14 days before getting a REVLIMID prescription
Pregnancy Test #2 If you can get pregnant, you must take another pregnancy test within 24 hours before getting a REVLIMID prescription
Enrollment You and your healthcare provider will then complete and submit the REVLIMID® (lenalidomide) Patient-Physician Agreement Form
Complete Mandatory Confidential Survey You and your healthcare provider will each complete a survey. Visit www.CelgeneRiskManagement.com or call 1-888-423-5436 and press 1 to take your survey
Prescription Your healthcare provider will send your prescription to a certified pharmacy
Pharmacy Call The certified pharmacy will contact you to provide counseling on the serious risks and safe-use conditions of REVLIMID before you receive your prescription. They will also coordinate delivery of REVLIMID to you
Receive REVLIMID REVLIMID will be shipped with a Medication Guide to the address you provide. A signature

will be required to receive this shipment

For each of your following prescriptions, pregnancy tests will be required depending on your ability to get pregnant. For specific pregnancy test requirements, please refer to page 10 of this guide. For full detailed information about the REVLIMID REMS[®] program requirements, please visit www.CelgeneRiskManagement.com or review the Patient Guide to REVLIMID REMS[®] Program.

For Males:

<p>Counseling</p> <p>Your healthcare provider will counsel you on why and how you and your partner should prevent pregnancy. Your healthcare provider will also inform you on not to share the drug, not to donate blood or sperm, and about appropriate contraceptive use. You should be instructed not to extensively handle or open REVLIMID capsules</p>
<p>Enrollment</p> <p>You and your healthcare provider will then complete and submit the Patient-Physician Agreement Form</p>
<p>Complete Mandatory Confidential Survey</p> <p>You will not have to take a survey for your first prescription, but will have to for the following ones. Visit www.CelgeneRiskManagement.com or call 1-888-423-5436 and press 1 to take your survey</p>
<p>Prescription</p> <p>Your healthcare provider will send your prescription to a certified pharmacy</p>
<p>Pharmacy Call</p> <p>The certified pharmacy will contact you to provide counseling on the serious risks and safe-use conditions of REVLIMID before you receive your prescription. They will also coordinate delivery of REVLIMID to you</p>
<p>Receive REVLIMID</p> <p>REVLIMID will be shipped with a Medication Guide to the address you provide. A signature will be required to receive this shipment</p>

For each of your following prescriptions, you will need to follow a similar process.

For full detailed information about the REVLIMID REMS[®] program requirements, please visit www.CelgeneRiskManagement.com or review the Patient Guide to REVLIMID REMS[®] Program.

What do all patients need to know about the REVLIMID REMS[®] program?

General guidelines

- This medicine is **only** for you. **Do not share it with anyone** even if they have symptoms like yours. It may harm them and can cause birth defects
- REVLIMID[®] (lenalidomide) must be kept out of the reach of children
- Do not open or unnecessarily handle REVLIMID capsules
- Keep REVLIMID in a cool, dry place
- Do **not** donate blood while you are taking REVLIMID, during breaks (dose interruptions), and for 4 weeks after stopping REVLIMID
- Unused REVLIMID capsules should be returned for disposal to Celgene by calling **1-888-423-5436** or to your REVLIMID prescriber, or to the pharmacy that dispensed the REVLIMID to you

What do females who can get pregnant need to know about the REVLIMID REMS® program?

A. Before taking REVLIMID® (lenalidomide)

- You must sign the REVLIMID® (lenalidomide) Patient-Physician Agreement Form that says you understand that REVLIMID should not be used during pregnancy, and that you agree not to become pregnant while taking REVLIMID
- If there is **any** chance that you can get pregnant, you must agree to use **at the same time** at least 1 highly effective method and at least 1 additional effective method of birth control **every time** you have sex with a male starting at least 4 weeks **before** taking REVLIMID
- Your healthcare provider must give you a pregnancy test 10 to 14 days before you receive your first prescription for REVLIMID, and again within 24 hours before you receive your first prescription for REVLIMID. If you are pregnant, you cannot take REVLIMID
- You will have pregnancy tests before starting REVLIMID and while taking REVLIMID, even if you agree not to have sex with a male
- Before your healthcare provider can write your prescription for REVLIMID, you must take part in a mandatory confidential survey. The survey will make sure that you understand and can follow information designed to prevent serious risks to unborn babies
- Before dispensing REVLIMID® (lenalidomide), your REVLIMID REMS® certified pharmacy will contact you to discuss treatment
- Your healthcare provider will talk with you about your birth control options

1. Choose at least 1 highly effective method and at least 1 additional effective method of birth control. Talk to your healthcare provider about the following acceptable birth control methods. See below.

Reliable Methods of Birth Control to Use Together

Highly effective birth control methods	Additional effective birth control methods
Intrauterine device (IUD)	
Hormonal methods (birth control pills, hormonal patches, injections, vaginal rings, or implants) +	Male latex or synthetic condom
Tubal ligation (having your tubes tied)	Diaphragm
Partner's vasectomy (tying of the tubes to prevent the passing of sperm)	Cervical cap

2. Use the 2 methods of birth control at the same time

- **Remember:** You must use at the same time at least 1 highly effective method and at least 1 additional effective method of birth control every time you have sex with a male. However, your healthcare provider may recommend that you use 2 different methods instead for medical reasons

What do females who can get pregnant need to know about the REVLIMID REMS[®] program? (continued)

- Talk to your healthcare provider to make sure that other medicines or dietary supplements you are taking do not interfere with your hormonal birth control methods
- **Remember, not having sex is the only birth control method that is 100% effective**

3. Unacceptable methods of birth control

- Progesterone-only “mini-pills”
- IUD Progesterone T
- Female condoms
- Natural family planning (rhythm method) or breastfeeding
- Fertility awareness
- Withdrawal
- Cervical shield*

4. Take pregnancy tests

- You must have a pregnancy test performed by your healthcare provider 10 to 14 days before receiving your first prescription for REVLIMID[®] (lenalidomide) and again within 24 hours before receiving your first prescription for REVLIMID. Both pregnancy tests must have a negative result

* A cervical shield should not be confused with a cervical cap, which is an effective secondary form of contraception.

What do females who can get pregnant need to know about the REVLIMID REMS® program? (continued)

B. While taking REVLIMID® (lenalidomide)

- If you are able to get pregnant, you must continue (including during breaks [dose interruptions]) to use **at the same time** at least 1 highly effective method and at least 1 additional effective method of birth control **every time** you have sex with a male
- **Remember, not having sex is the only birth control method that is 100% effective**
- You must talk to your healthcare provider before changing any birth control methods you have already agreed to use
- You will have a pregnancy test performed by your healthcare provider:
 - Every week during the first 4 weeks of treatment, then
 - Every 4 weeks if your menstrual cycles are regular, or
 - Every 2 weeks if your cycles are irregular
 - If you miss your period or have unusual menstrual bleeding, or
 - If your medication is not dispensed within 7 days of taking the pregnancy test
- If you had sex with a male without using birth control or if you think your birth control has failed, stop taking REVLIMID® (lenalidomide) immediately and call your healthcare provider right away. Your healthcare provider will discuss your options, which may include emergency birth control. Do not wait until your next appointment to tell your healthcare provider if you miss your menstrual period or if you think you may be pregnant
- If you get pregnant, or think you may be pregnant, you must immediately stop taking REVLIMID. Contact your healthcare provider immediately to discuss your pregnancy. If you do not have an obstetrician, your healthcare provider will refer you to one for care and counseling. If your healthcare provider is not available, call the Celgene Customer Care Center at **1-888-423-5436**
- You must not breastfeed a baby while you are taking REVLIMID
- In order to continue receiving REVLIMID, you must take part in a mandatory confidential survey every month. You must also continue to discuss your treatment with your REVLIMID REMS® healthcare provider. To take the survey, please visit www.CelgeneRiskManagement.com, access the Celgene REMS mobile app, or call the Celgene Customer Care Center at **1-888-423-5436**

C. After you have stopped taking REVLIMID® (lenalidomide)

- You must continue to use **at the same time** at least 1 highly effective method and at least 1 additional effective method of birth control **every time** you have sex with a male:

- For at least 4 weeks after stopping REVLIMID, or
- Do not have any sex with a male for 4 weeks after stopping REVLIMID

See also “General guidelines” on page 5 for requirements for all patients.

What do females who can not get pregnant need to know about the REVLIMID REMS[®] program?

A. Before taking REVLIMID[®] (lenalidomide)

- You must sign the REVLIMID[®] (lenalidomide) Patient-Physician Agreement Form that says you are currently not pregnant and are not able to get pregnant. This means that:
 - You have been in natural menopause for at least 2 years, or
 - You have had both ovaries and/or uterus removed
- For females who have not started their period (menstruation) and are under the age of 18, a parent or legal guardian must sign the REVLIMID[®] (lenalidomide) Patient-Physician Agreement Form that says the patient is not pregnant, is not able to get pregnant, and/or will not be having sex with a male for at least 4 weeks before starting REVLIMID
- Before your healthcare provider can write your prescription for REVLIMID, you must take part in a mandatory confidential survey. The survey will make sure that you understand and can follow information designed to prevent serious risks to unborn babies
- Before dispensing REVLIMID[®] (lenalidomide), your REVLIMID REMS[®] certified pharmacy will contact you to discuss treatment

B. While taking REVLIMID

- In order to continue receiving REVLIMID, you must take part in a mandatory confidential survey every six months. You must also continue to discuss your treatment with your REVLIMID REMS[®] healthcare provider. To take the survey, please visit www.CelgeneRiskManagement.com, access the Celgene REMS mobile app, or call the Celgene Customer Care Center at **1-888-423-5436**

See also “General guidelines” on page 5 for requirements for all patients.

What do males need to know about the REVLIMID REMS[®] program?

- You must use a latex or synthetic condom, **every time** you have sex with a female who is able to get pregnant, even if you have had a successful vasectomy (tying of the tubes to prevent the passing of sperm)

A. Before taking REVLIMID[®] (lenalidomide)

- You must sign the REVLIMID[®] (lenalidomide) Patient-Physician Agreement Form. You must agree that while taking REVLIMID you will use a latex or synthetic condom **every time** you have sex with a female who is pregnant or who is able to get pregnant, even if you have had a successful vasectomy (tying of the tubes to prevent the passing of sperm)
- Before dispensing REVLIMID, your REVLIMID REMS[®] certified pharmacy will contact you to discuss treatment

B. While taking REVLIMID

- You must use a latex or synthetic condom, **every time** (including during breaks [dose interruptions]) you have sex with a female who is pregnant or who is able to get pregnant, even if you have had a successful vasectomy (tying of the tubes to prevent the passing of sperm)
- **Remember, not having sex is the only birth control method that is 100% effective**
- You must tell your healthcare provider right away if you had sex with a female without using a latex or synthetic condom, or if you think for any reason that your partner may be pregnant, or if your partner is pregnant. If your healthcare provider is not available, call the Celgene Customer Care Center at **1-888-423-5436**
- You must **not** donate sperm while taking REVLIMID[®] (lenalidomide) (including during breaks [dose interruptions])
- In order to continue receiving REVLIMID, you must take part in a mandatory confidential survey every month. You must also continue to discuss your treatment with your REVLIMID REMS[®] healthcare provider. To take the survey, please visit **www.CelgeneRiskManagement.com**, access the Celgene REMS mobile app, or call the Celgene Customer Care Center at **1-888-423-5436**

C. After you have stopped taking REVLIMID

- For 4 weeks after receiving your last dose of REVLIMID, you must use a latex or synthetic condom **every time** you have sex with a female who is pregnant or who is able to get pregnant, even if you have had a successful vasectomy (tying of the tubes to prevent the passing of sperm)
- You must **not** donate sperm for 4 weeks after stopping REVLIMID

See also “General guidelines” on page 5 for requirements for all patients.

Mandatory confidential patient surveys

As a patient who is enrolled in the REVLIMID REMS[®] program for REVLIMID[®] (lenalidomide), you will need to complete a brief mandatory confidential survey as outlined below.

Adult females who can get pregnant

- Initial survey before first prescription
- Monthly

Adult females who can not get pregnant

- Initial survey before first prescription
- Every six months

Female children

- Initial survey before first prescription
- Monthly

Males

- No initial survey
- Monthly

Mandatory confidential survey process

- When your healthcare provider tells you to take the survey, go to the patient Mandatory Confidential Survey section of **www.CelgeneRiskManagement.com**, the Celgene REMS mobile app, or call the Celgene Customer Care Center at **1-888-423-5436**
- Be prepared with your patient identification number
- After completing your survey, your healthcare provider will also complete a survey. Your healthcare provider will then receive authorization to write your prescription
- The prescription will be sent to a REVLIMID REMS[®] certified pharmacy. The REVLIMIDREMS[®] certified pharmacy will contact you to discuss your REVLIMID[®] (lenalidomide) therapy. You will not receive your medication until you speak with the REVLIMID REMS[®] certified pharmacy
- For more information, contact the Celgene Customer Care Center at **1-888-423-5436**

Warning to patients taking REVLIMID® (lenalidomide)

Attention females:

Do **not** take REVLIMID if you are pregnant, if you are breastfeeding, or if you are able to get pregnant and are not using **at the same time** at least 1 highly effective method and at least 1 additional effective method of birth control **every time** you have sex with a male.

Attention males:

You must use a latex or synthetic condom **every time** you have sex with a female who is pregnant or who is able to get pregnant, even if you have had a successful vasectomy (tying of the tubes to prevent the passing of sperm).

You must **not** donate sperm while taking REVLIMID, during breaks (dose interruptions), and for 4 weeks after stopping REVLIMID.

Attention all patients:

You must **not** donate blood while taking REVLIMID® (lenalidomide), during breaks (dose interruptions), and for 4 weeks after stopping REVLIMID.

This medicine is **only** for you. **Do not share it with anyone** even if they have symptoms like yours. It may harm them and can cause birth defects.

REVLIMID must be kept out of the reach of children. Return any unused REVLIMID capsules for disposal to Celgene by calling **1-888-423-5436**, or to your REVLIMID prescriber, or to the pharmacy that dispensed the REVLIMID to you.

You may require regular blood tests during REVLIMID treatment. Consult with your healthcare provider.

[Back Cover]

For more information about REVLIMID[®] (lenalidomide) and the REVLIMID REMS[®] program, please visit **www.CelgeneRiskManagement.com**, or call the Celgene Customer Care Center at **1-888-423-5436**.

Celgene Corporation

86 Morris Ave

Summit, NJ 07901

REVLIMID is only available under a restricted distribution program, REVLIMID REMS[®].

Please see full Prescribing Information, including Boxed WARNINGS, CONTRAINDICATIONS, WARNINGS AND PRECAUTIONS, ADVERSE REACTIONS, and Medication Guide, enclosed.

((Celgene logo))

((REVLIMID REMS[®] logo))

((REVLIMID logo))

REVLIMID[®] is a registered trademark of Celgene Corporation.

REVLIMID REMS[®] is a trademark of Celgene Corporation.

© 2016 Celgene Corporation 6/16

REMS-REV16734