

NADA Number: 108-114

Trade Name	Tomorrow® Infusion Cefa-Dri®
Sponsor	Fort Dodge Animal Health, Division of Wyeth
Ingredients	Cephapirin Benzathine
Species	Cattle, dry cows
Routes of Administration	Intramammary
Dose Form	Gel
Drug Form	Gel
Dispensing Status	OTC
Dosage Amount, Indications & Limitations	<p>526.363 Cephapirin benzathine.</p> <p>Specifications: Each 10 milliliter disposable syringe contains 300 milligrams of cephapirin activity (as cephapirin benzathine) in a peanut-oil gel.</p> <p>Conditions of use:</p> <p>Cattle (dry cows)</p> <p>Amount: Infuse contents of one syringe into each infected quarter.</p> <p>Indications: Use in dry cows for treatment of mastitis caused by susceptible strains of <i>Streptococcus agalactiae</i> and <i>Staphylococcus aureus</i> including penicillin-resistant strains.</p> <p>Limitations: Infuse each infected quarter following last milking or early in the dry period, but no later than 30 days before calving. Milk from treated cows must not be used for food during the first 72 hours after calving. Animals infused with this product must not be slaughtered for food until 42 days after the latest infusion. For use in dry cows only.</p>
Tolerances	A tolerance of 0.02 parts per million (ppm) is established for residues of cephapirin in the milk and 0.1 ppm in the uncooked edible tissues of dairy cattle.