

NADA Number: 038-878

Trade Name	Coban® - 60 Elancoban-100 Coban® - 45 Coban® - 110
Sponsor	Elanco Animal Health, A Division of Eli Lilly & Co.
Ingredients	Monensin Sodium
Species	CHICKENS (Chicken, replacements) CHICKENS (Chicken, broilers) CHICKENS (Chicken, not laying eggs for human consumption) QUAIL (Bobwhite quail, growing) TURKEYS (Turkey, no use class stated or implied)
Routes of Administration	Per Os
Dosage Form	Medicated feed
Dispensing Status	OTC
Patent Number (Expiration Date)	3839557 (No Expiration Date) 3995027 (No Expiration Date) 4061755 (No Expiration Date) 4075323 (No Expiration Date) 4083962 (No Expiration Date) 4218438 (No Expiration Date) 4333919 (No Expiration Date) 4366168 (No Expiration Date) 4405609 (No Expiration Date) 4468380 (No Expiration Date)
Dosage Amount, Indications & Limitations	<p>558.355 Monensin.</p> <p>Specifications: Monensin as the base or the sodium salt, contains a minimum of 90 percent monensin activity derived from monensin A and a minimum of 95 percent derived from monensin A plus B. Approvals for Type A medicated articles containing 45 and 60 grams per pound, as monensin sodium.</p> <p>Amount: 45, or 60 grams per pound</p> <p>Conditions of use:</p> <p>Chickens (broilers)</p> <p>Amount per ton: Monensin, 90 - 110 grams.</p> <p>Indications for use: As an aid in the prevention of coccidiosis caused by <i>E. necatrix</i>, <i>E. tenella</i>, <i>E. acervulina</i>, <i>E. brunetti</i>, <i>E. mivati</i>, and <i>E. maxima</i>.</p> <p>Limitations: Do not feed to laying chickens. Feed continuously as the sole ration. In the absence of coccidiosis, the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain. As monensin or monensin sodium.</p> <p>Chickens (replacements intended for use as cage layers)</p> <p>Amount per ton: Monensin, 90 to 110 grams.</p> <p>Indications for use: As an aid in the prevention of coccidiosis caused by <i>E. necatrix</i>, <i>E. tenella</i>, <i>E. acervulina</i>, <i>E. brunetti</i>, <i>E. mivati</i>, and <i>E. maxima</i>.</p> <p>Limitations. Do not feed to laying chickens. Feed continuously as sole ration. As monensin sodium. Do not feed to chickens over 16 weeks of age.</p> <p>Amount. 110 grams per pound</p> <p>Conditions of use:</p> <p>Chickens (broilers)</p> <p>Amount per ton. Monensin, 90 - 110 grams.</p> <p>Indications for use: As an aid in the prevention of coccidiosis caused by <i>E. necatrix</i>, <i>E. tenella</i>, <i>E. acervulina</i>, <i>E. brunetti</i>, <i>E. mivati</i>, and <i>E. maxima</i>.</p> <p>Limitations: Do not feed to laying chickens. Feed continuously as the sole ration. In the absence of coccidiosis, the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain. As monensin or monensin sodium.</p> <p>Labeling: The labeling of Type A articles containing monensin intended for use in chickens shall bear the following caution statements: Do not allow horses or other equines access to formulations containing monensin. Ingestion of monensin by equines has been fatal. Must be thoroughly mixed in feeds before use. Do not feed undiluted.</p> <p>Conditions of use:</p>

	<p>Turkeys</p> <p>Amount: 45 and 60 grams per pound, as monensin sodium.</p> <p>Amount per ton: Monensin, 54 to 90 grams.</p> <p>Indications for use: For the prevention of coccidiosis in turkeys caused by <i>E. adenoeides</i>, <i>E. meleagrimitis</i>, and <i>E. gallopavonis</i>.</p> <p>Limitations: Feed continuously as the sole ration to growing turkeys from 1 day of age as monensin sodium. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin.</p> <p>Amount: 45 and 60 grams per pound, as monensin sodium.</p> <p>Labeling: The labeling of all formulations containing monensin shall bear the following caution statement: Do not allow horses or other equines access to formulations containing monensin. Ingestion of monensin by equines has been fatal.</p> <p>Quail (bobwhite)</p> <p>Amount per ton: Monensin, 73 grams.</p> <p>Indications for use: For the prevention of coccidiosis in growing bobwhite quail caused by <i>Eimeria dispersa</i> and <i>E. Lettyae</i>.</p> <p>Limitations: Feed continuously as the sole ration. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin.</p> <p>Labeling: The labeling of all formulations containing monensin shall bear the following caution statement: Do not allow horses or other equines access to formulations containing monensin. Ingestion of monensin by equines has been fatal.</p> <p>Amount: 45, and 60 grams per pound, as monensin sodium.</p>
<p>Tolerances</p>	<p>Monensin Sodium: A tolerance of 0.10 part per million (ppm) in cattle liver, 0.05 ppm in muscle, kidney, and fate.</p> <p>A tolerance of 0.05 ppm in goats edible tissue.</p> <p>A tolerance for residue of monensin in chickens, turkeys and quail is not needed.</p> <p>A tolerance for residues of monensin in milk is not needed.</p>