

NADA Number: 092-444

Trade Name	Rumatel® 88
Sponsor	Phibro Animal Health Corp.
Ingredients	Morantel Tartrate
Species	Cattle (Cattle, no use class stated or implied) GOATS (Goat, no use class stated or implied)
Routes of Administration	Per Os
Dosage Form	Medicated feed
Dispensing Status	OTC
Patent Number (Expiration Date)	3644624 (No Expiration Date)
Dosage Amount, Indications & Limitations	<p>558.360 Morantel tartrate.</p> <p>Specifications: Type A medicated articles: 88 grams per pound</p> <p>Conditions of use:</p> <p>Cattle</p> <p>Amount: 0.44 to 4.4 grams of morantel tartrate per pound of feed.</p> <p>Indications: For removal and control of mature gastrointestinal nematode infections of cattle including stomach worms (<i>Haemonchus</i> species, <i>Ostertagia</i> species <i>Trichostrongylus</i> species), worms of the small intestine (<i>Cooperia</i> species, <i>Trichostrongylus</i> species, <i>Nematodirus</i> species), and worms of the large intestine (<i>Oesophagostomum radiatum</i>).</p> <p>Limitations: Feed as a single therapeutic treatment at 0.44 gram of morantel tartrate per 100 pounds of body weight. Fresh water should be available at all times. When medicated feed is consumed, resume normal feeding. Conditions of constant worm exposure may require re-treatment in 2 to 4 weeks. Do not use in Type B or Type C medicated feeds containing bentonite. Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.</p> <p>Goats</p> <p>Amount: 0.44 to 4.4 grams of morantel tartrate per pound of feed.</p> <p>Indications: For removal and control of mature gastrointestinal nematode infections of goats including <i>Haemonchus contortus</i>, <i>Ostertagia</i> (*<i>Teladorsagia</i>) <i>circumcincta</i>, and <i>Trichostrongylus axei</i>.</p> <p>Limitations: Feed as a single therapeutic treatment at 0.44 gram of morantel tartrate per 100 pounds of body weight. Fresh water should be available at all times. When medicated feed is consumed, resume normal feeding. Conditions of constant worm exposure may require re-treatment in 2 to 4 weeks. Do not use in Type B or Type C medicated feeds containing bentonite. Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.</p>
Tolerances	Morantel Tartrate: A tolerance of 0.7 part per million is established for N-methyl-1,3-propanediamine (MAPA, marker residue) in the liver (target tissue) of cattle and goats. A tolerance for residues of morantel tartrate in milk is not required.