

**NADA Number: 125-933**

<b>Trade Name</b>	Romet®-30
<b>Sponsor</b>	Pharmaq AS
<b>Ingredients</b>	Sulfadimethoxine Ormetoprim
<b>Species</b>	<b>FISH</b> (Salmonids, reared) <b>FISH</b> (Catfish, reared)
<b>Routes of Administration</b>	Per Os
<b>Dosage Form</b>	Medicated feed
<b>Dispensing Status</b>	OTC
<b>Patent Number (Expiration Date)</b>	3903275 (No Expiration Date)
<b>Dosage Amount, Indications &amp; Limitations</b>	<p>558.575 Sulfadimethoxine, ormetoprim.</p> <p>Specifications: Type A medicated articles: 25 percent sulfadimethoxine and 5 percent ormetoprim.</p> <p>Conditions of use:</p> <p>Salmonids</p> <p>Amount: 50 milligrams of active ingredients per kilogram of body weight per day.</p> <p>Indications: For the control of furunculosis in salmonids (trout and salmon) caused by <i>Aeromonas salmonicida</i> strains susceptible to sulfadimethoxine and ormetoprim combination.</p> <p>Limitations: Administer for 5 consecutive days. Withdraw 42 days before release as stocker fish or slaughter.</p> <p>Catfish</p> <p>Amount: 50 milligrams of active ingredients per kilogram of body weight per day.</p> <p>Indications: For control of enteric septicemia of catfish caused by <i>Edwardsiella ictaluri</i> strains susceptible to sulfadimethoxine and ormetoprim combination.</p> <p>Limitations: Administer for 5 consecutive days. Withdraw 3 days before slaughter or release as stocker fish.</p>
<b>Tolerances</b>	<p>Sulfadimethoxine: Tolerances are established for residues of sulfadimethoxine in edible products of animals as follows: a. Chickens, turkeys, cattle, ducks, salmonids, catfish and chukar partridges: 0.1 part per million (negligible residue) in uncooked edible tissues b. Milk: 0.01 part per million (negligible residue). Tolerances are established for residues of sulfadimethoxine in edible products of animals as follows: a. Chickens, turkeys, cattle, ducks, salmonids, catfish and chukar partridges: 0.1 part per million (negligible residue) in uncooked edible tissues b. Milk: 0.01 part per million (negligible residue).</p> <p>Ormetoprim: A tolerance of 0.1 part per million is established for negligible residues of ormetoprim in the edible tissues of chickens, turkeys, ducks, salmonids, and catfish.</p>