NADA Number: 128-686	
Trade Name	Bio-Cox® Type A Medicated Article
Sponsor	Zoetis Inc.
Ingredients	Salinomycin Sodium
Species	**CHICKENS** (Chicken, broilers excluding layers) **CHICKENS** (Chicken, roasters) **CHICKENS** (Chicken, replacements) **QUAIL** (Quail, no use class stated or implied)
Routes of Administration	Per Os
Dosage Form	Medicated feed
Dispensing Status	OTC
Patent Number (Expiration Date)	3857948 (No Expiration Date)
Exclusivity	Granted to add two new species roasters and replacements (breeder and layers) chickens.
Dosage Amount, Indications & Limitations	558.550 Salinomycin. Specifications: Type A medicated articles: 30 or 60 grams of activity from salinomycin sodium biomass. Conditions of use: Chickens (broilers) Amount: Salinomycin 40 to 60 grams per ton of feed. Indications: For the prevention of coccidiosis caused by Eimeria tenella, E. necatrix, E. acervulina, E. maxima, E. brunetti, and E. mivati. Limitations: Feed continuously as sole ration. Do not feed to layers. Not approved for use with pellet binders. May be fatal if accidentally fed to adult turkeys or horses. Chickens (roaster and replacement) Amount: Salinomycin 40 to 60 grams per ton of feed. Indications: For prevention of coccidiosis caused by Eimeria tenella, E. necatrix, E. acervulina, E. maxima, E. brunetti, and E. mivati. Limitations: Feed continuously as sole ration. Do not feed to laying hens producing eggs for human consumption. Not approved for use with pellet binders. May be fatal if accidentally fed to horses or adult turkeys. Quail Amount: Salinomycin 50 grams per ton of feed. Indications: For the prevention of coccidiosis caused by E. dispersa and E. lettyae. Limitations: Feed continuously as sole ration. Not approved for use with pellet binders. May be fatal if accidentally fed to adult turkeys or horses.