

NADA Number: 141-087

Trade Name	Quest® Gel Quest® 2% Equine Oral Gel
Sponsor	Zoetis Inc.
Ingredients	Moxidectin
Species	EQUIDS (Horse, not for meat production) EQUIDS (Pony, no use class stated or implied)
Routes of Administration	Per Os
Dosage Form	Gel
Dispensing Status	OTC
Patent Number (Expiration Date)	4916154 (No Expiration Date)
Exclusivity	Granted for the treatment and control of a species of small strongyles (<i>Coronocyclus labratus</i>) in horses and ponies. Granted for treatment and control of <i>Gasterophilus nasalis</i> (3rd instars) in horses and ponies. Granted for the treatment of certain stages of certain gastrointestinal parasites of horses and ponies. Exclusivity is granted based on a supplemental approval showing efficacy in the treatment and control of two additional species of small strongyles (<i>Cylicocyclus radiatus</i> and <i>Petrovinema poculatus</i> .)
Dosage Amount, Indications & Limitations	520.1452 Moxidectin gel. Specifications: The gel contains 2 percent moxidectin (20 milligrams per milliliter). Conditions of use: Horses (and ponies) Amount: Each milligram of gel contains per 20 milligrams (2 percent) of moxidectin. Indications: For the treatment and control of large strongyles: <i>Strongylus vulgaris</i> (adults and L4/L5 arterial stages), <i>S. edentatus</i> (adult and tissue stages), <i>Triodontophorus brevicauda</i> (adults), and <i>T. serratus</i> (adults); small strongyles (adults): <i>Cyathostomum</i> species, including <i>C. catinatum</i> and <i>C. pateratum</i> ; <i>Cylicocyclus</i> species, including <i>C. insigne</i> , <i>C. leptostomum</i> , <i>C. nassatus</i> and <i>C. radiatus</i> ; <i>Cyliocostephanus</i> species, including <i>C. calicatus</i> , <i>C. goldi</i> , <i>C. longibursatus</i> , and <i>C. minutus</i> ; <i>Coronocyclus</i> species, including <i>C. coronatus</i> , <i>C. labiatus</i> , and <i>C. labratus</i> ; and <i>Gyalocephalus capitatus</i> ; and <i>Petrovinema poculatus</i> ; small strongyles: undifferentiated lumenal larvae; encysted cyathostomes (late L3 and L4 mucosal cyathostome larvae); ascarids: <i>Parascaris equorum</i> (adults and L4 larval stages); pinworms: <i>Oxyuris equi</i> (adults and L4 larval stages); hairworms: <i>Trichostrongylus axei</i> (adults); large-mouth stomach worms: <i>Habronema muscae</i> (adults); and horse stomach bots: <i>Gasterophilus intestinalis</i> (2nd and 3rd instars) and <i>G. nasalis</i> (3rd instars). One dose also suppresses strongyle egg production for 84 days. Limitations: For oral use in horses and ponies 6 months of age and older. Do not use in horses intended for human consumption.
Tolerances	Moxidectin: The tolerance for parent moxidectin (the marker residue) in edible tissues of cattle is 200 parts per billion (ppb) in liver (the target tissue) and 50 ppb in muscle. The tolerance for parent moxidectin is 40 ppb in milk.