

ANADA Number: 200-075

Trade Name	Sacox® 60 Sacox® Type A Medicated Article
Sponsor	Huvepharma AD
Ingredients	Salinomycin Sodium
Pioneer	128-686 (Zoetis Inc.)
Species	CHICKENS (Chicken, broilers excluding layers) CHICKENS (Chicken, replacements) CHICKENS (Chicken, roasters) QUAIL (Quail, no use class stated or implied)
Routes of Administration	Per Os
Dosage Form	Powder
Dispensing Status	OTC
Dosage Amount, Indications & Limitations	<p>558.550 Salinomycin.</p> <p>Specifications: Type A medicated articles--30 or 60 grams of activity per pound from salinomycin sodium biomass.</p> <p>Conditions of use:</p> <p>Chickens (broilers)</p> <p>Amount: Salinomycin 40 to 60 grams per ton of feed.</p> <p>Indications: For the prevention of coccidiosis caused by Eimeria tenella, E. necatrix, E. acervulina, E. Maxima, E. Brunetti, and E. mivati.</p> <p>Limitations: Feed continuously as sole ration. Do not feed to layers. Not approved for use with pellet binders. May be fatal if accidentally fed to adult turkeys or horses.</p> <p>Chickens (roaster and replacement)</p> <p>Amount: Salinomycin 40 to 60 grams per ton of feed.</p> <p>Indications: For prevention of coccidiosis caused by Eimeria tenella, E. necatrix, E. acervulina, E. maxima, E. brunetti, and E. mivati.</p> <p>Limitations: Feed continuously as sole ration. Do not feed to laying hens producing eggs for human consumption. Not approved for use with pellet binders. May be fatal if accidentally fed to horses or adult turkeys.</p> <p>Quail</p> <p>Amount: Salinomycin 50 grams per ton of feed.</p> <p>Indications: For the prevention of coccidiosis caused by E. dispersa and E. lettyae.</p> <p>Limitations: Feed continuously as sole ration. Not approved for use with pellet binders. May be fatal if accidentally fed to adult turkeys or horses.</p>