

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

73416

DRAFT FINAL PRINTED LABELING

PRINTED IN U.S.A. CH 2272 (7-98)

DIRECTIONS FOR USE - 6 MINUTE SCRUB
(1) Wet hands and forearms with warm water. (2) Use nail cleaner and then apply scrub solution from sponge side. Work up lather. (3) Scrub nails, cuticles and interdigital spaces thoroughly for 3 minutes with brush side and the hands and forearms with the sponge side. (4) Rinse with warm water. **REPEAT** scrub for 3 more minutes: use sponge side only. (5) Rinse hands and arms thoroughly. (6) Dry thoroughly.

E-Z Scrub® 106

CAT. NO. 371065

NDC 17271-106-05

Antimicrobial Surgical Scrub Brush/Sponge with HIBICLENS® (chlorhexidine gluconate 4% w/v) Filled with 22 mL of HIBICLENS® (4% w/v HIBICLENS® (chlorhexidine gluconate)) an antiseptic with inactive ingredients: D30 Red #40 gluconolactone USP, herbacol 15.393 TB, isopropyl alcohol USP, lauramine oxide 13%, aluminum oxide, poloxamer 237, purified water USP, sodium hydroxide NF. Chlorhexidine gluconate provides rapid bactericidal action and has a potent antimicrobial effect against a wide range of microorganisms. HIBICLENS® is a product of Zeneca Pharmaceuticals, a Business Unit of Zeneca Inc., Wilmington, DE 19850-5437. For quality and technical services call 1-800-333-4813. **ONE BRUSH/SPONGE FOR SURGICAL HAND SCRUB • FOR EXTERNAL USE ONLY DISCARD AFTER USE • Store at controlled room temperature, 15-30 C (59-86 F)**

**BECTON
DICKINSON**

Manufactured For:

Becton Dickinson Surgical Systems
Becton Dickinson and Company
Franklin Lakes, NJ 07417

WARNINGS: FOR EXTERNAL USE ONLY. KEEP OUT OF EYES, EARS AND MOUTH. CHLORHEXIDINE GLUCONATE PRODUCTS SHOULD NOT BE USED AS A PREOPERATIVE SKIN PREPARATION OF THE FACE OR HEAD. MISUSE OF CHLORHEXIDINE GLUCONATE PRODUCTS HAS BEEN REPORTED TO CAUSE SERIOUS AND PERMANENT EYE INJURY WHEN PERMITTED TO ENTER AND REMAIN IN THE EYE DURING SURGICAL PROCEDURES. IF THIS PRODUCT SHOULD CONTACT THE EYE, RINSE OUT PROMPTLY AND THOROUGHLY WITH WATER. Avoid contact with the ear. This product should not be used by persons who have a sensitivity to it. An allergic reaction to chlorhexidine gluconate has been reported to cause deafness when applied to the middle ear through perforated ear drums. Irritation, sensitization and general allergic reactions have been reported with chlorhexidine-containing products, especially in the genital area. Adverse reactions occur; discontinue use immediately and if severe contact a physician. Keep this and all drugs out of the reach of children. In case of accidental ingestion seek professional assistance or contact a Poison Control Center immediately.