



NDA 208025/S-008

SUPPLEMENT APPROVAL

Dexcel Pharma Technologies Ltd
c/o: Icon Clinical Research LLC (Authorized Agent)
Attention: Amy Kneifel, RAC
Director, Regulatory Affairs
79 TW Alexander Drive
4401 Research Commons Bldg, Suite 300
Durham, NC 27709

Dear Ms. Kneifel:

Please refer to your Supplemental New Drug Application (sNDA) dated and received October 10, 2018, and your amendment, submitted under section 505(b) of the Federal Food, Drug, and Cosmetic Act (FDCA) for lansoprazole delayed-release, orally disintegrating tablet, 15 mg.

This “changes being effected” supplemental new drug application provides for a modification of the Drug Facts label to include “**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Acid reducers may interact with certain prescription drugs.” This update to the drug to drug interaction warning is in response to the Agency’s communications of June 29 and August 10, 2018.

We have completed our review of this application, as amended. It is approved, effective on the date of this letter, for use as recommended in the agreed-upon labeling text.

LABELING

Submit final printed labeling, as soon as they are available, but no more than 30 days after they are printed. The final printed labeling must be identical to the labeling listed in the below table, and must be in the “Drug Facts” format (21 CFR 201.66), where applicable.

Submitted Labeling	Date(s) Submitted
14-count inner carton	October 23, 2018
14-count outer carton - <i>Main</i>	October 10, 2018
28-count outer carton – <i>Main</i>	October 10, 2018
42-count outer carton - <i>Main</i>	October 10, 2018

14-count outer carton with 'Orally Disintegrating Tablets' flag	October 10, 2018
28-count outer carton with 'Orally Disintegrating Tablets' flag	October 10, 2018
42-count outer carton with 'Orally Disintegrating Tablets' flag	October 10, 2018
14-count outer carton with 'Child-resistant' flag	October 10, 2018
28-count outer carton with 'Child-resistant' flag	October 10, 2018
42-count outer carton with 'Child-resistant' flag	October 10, 2018
28-count outer carton with 'Value 2 Pack' flag	October 10, 2018
42-count outer carton with 'Value 3 Pack' flag	October 10, 2018
28-count outer carton with 'Buy One 14 Count, Get One 14 Count Free!' flag	October 10, 2018
28-count outer carton with '100% More Free! 14 Free Tablets!' flag	October 10, 2018
28-count outer carton with 'Bonus Size! 14 Tablets Free' flag	October 10, 2018
28-count outer carton with 'Bonus Size! 14 Free' flag	October 10, 2018
28-count outer carton with 'Bonus! + 14 Free' flag	October 10, 2018
28-count outer carton with 'Bonus! 14 Tablets FREE' flag	October 10, 2018
42-count outer carton with 'Bonus Pack: 28 + 14 Free' flag	October 10, 2018
42-count outer carton with 'Bonus Size! 14 Tablets Free' flag	October 10, 2018
42-count outer carton with 'Bonus Size! 14 Free' flag	October 10, 2018
42-count outer carton with 'Bonus! + 14 Free' flag	October 10, 2018
42-count outer carton with 'Bonus! 14 Tablets FREE' flag	October 10, 2018
Consumer Information Leaflet	October 10, 2018

The final printed labeling should be submitted electronically according to the guidance for industry titled *Providing Regulatory Submissions in Electronic Format — Certain Human Pharmaceutical Product Applications and Related Submissions Using the eCTD Specifications (April 2017, Revision 4)*. For administrative purposes, designate this submission “**Final Printed Labeling for approved NDA 208025/S-008.**” Approval of this submission by FDA is not required before the labeling is used.

If you are interested in marketing other package configurations in the future, (e.g., bottles containing greater than 14 tablets, package sizes greater than 42-count), a prior approval labeling

supplement that includes data to adequately demonstrate appropriate consumer comprehension of limitations of use must be submitted.

DRUG REGISTRATION AND LISTING

All drug establishment registration and drug listing information is to be submitted to FDA electronically, via the FDA automated system for processing structured product labeling (SPL) files (eLIST). At the time that you submit your final printed labeling (FPL), the content of labeling (Drug Facts) should be submitted in SPL format as described at <http://www.fda.gov/ForIndustry/DataStandards/StructuredProductLabeling/default.htm>. Information on submitting SPL files using eLIST may be found in the guidance for industry titled "SPL Standard for Content of Labeling Technical Qs and As" at <http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM072392.pdf>. In addition, representative container or carton labeling, whichever includes Drug Facts, (where differences exist only in the quantity of contents statement) should be submitted as a JPG file.

REPORTING REQUIREMENTS

We remind you that you must comply with reporting requirements for an approved NDA (21 CFR 314.80 and 314.81).

If you have any questions, call Janice Adams-King, Safety Regulatory Health Project Manager, at 301-796-3713.

Sincerely,

{See appended electronic signature page}

Valerie Pratt, MD
Deputy Director for Safety
Division of Nonprescription Drug Products
Office of Drug Evaluation IV
Center for Drug Evaluation and Research

ENCLOSURES:
Carton and Container Labeling

Made in Israel
 Manufactured by:
Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by
 GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Drug Facts (continued)

Directions ■ adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity.
 ■ protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, flavon, hypromellose, hypromellose phthalate, maize maltoedextrin, methylcellulose, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium lauryl sulfate, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST

Drug Facts

Active ingredient (in each tablet) Lansoprazole 15 mg; Acid Reducer

Purpose
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ **Allergy alert:** do not use if you are allergic to lansoprazole
 ■ do not use
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools; these may be signs of a serious condition. See your doctor.
 ■ Ask a doctor before use if you have
 ■ liver disease
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with **lightheadedness, sweating or dizziness**
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 ■ frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 ■ In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Location for lot number and expiration date

8U174 FA C6 092818



KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC 64239-181-01

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

■ May take 1 to 4 days for full effect

14 TABLETS

ONE 14-DAY COURSE OF TREATMENT

Melts in your mouth
 Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



Made in Israel
 Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 *This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Drug Facts
 Active ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer
Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
Warnings
 ■ **Do not use** if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ **Allergy alert:** do not use if you are allergic to lansoprazole
Ask a doctor before use if you have
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with **lightheadedness, sweating or dizziness**
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
 ■ frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting ■ stomach pain
Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain
If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)
Directions
 ■ adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol
Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
Other information
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F) ■ Keep product out of high heat and humidity
 ■ protect product from moisture
Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolvidone, flavor, hypromellose, hypromellose phthalate, maize maltoedextrin, maltitol, mannitol, microcrystalline cellulose, polyethylene glycol, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

8U130 FA C6 092818

Location for lot number and expiration date

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
 ■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6 092818

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active ingredient (in each tablet) Acid Reducer
Lansoprazole 15 mg

Use
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Do not use
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have
■ liver disease
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
Frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain
Ask a doctor or pharmacist before use if you are taking a prescription drug.
Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Other information
■ read the directions, warnings and package insert before use
■ keep the carton and package insert. They contain important information.
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
■ protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize polyborate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST


Drug Facts (continued)

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ take 1 tablet before eating in the morning
■ do not crush or chew tablets
■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor.
■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months
unless directed by a doctor
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-03

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

24 Hour
Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

- May take 1 to 4 days for full effect

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Made in Israel
 Manufactured by:
Dexel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by
 GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Lansoprazole 15 mg; Acid Reducer</p> <p>Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ Allergy alert: do not use if you are allergic to lansoprazole</p> <p>Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have ■ liver disease ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness</p> <p>chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p> <p>If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>Other information ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition. ■ read the directions, warnings and package insert before use ■ keep the carton and package insert. They contain important information. ■ store at 20-25°C (68-77°F). ■ Keep product out of high heat and humidity. ■ protect product from moisture</p> <p>Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, HAVON, hypromellose, hypromellose pthalate, maize maibedextrin, malitol, mannitol, meglumine, microcrystalline cellulose, polyacrylate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9266; weekdays 7:30 AM to 5:00 PM EST</p>
---	---

Location for lot number and expiration date

8U174 FA C6-V 100418



KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC 64239-181-01

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
 ■ May take 1 to 4 days for full effect

Orally Disintegrating Tablets



Actual size

14 TABLETS ONE 14-DAY COURSE OF TREATMENT

Melts in your mouth
 Dissolves without water

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 Made in Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Drug Facts
 Active ingredient (in each tablet) Lansoprazole 15 mg
 Purpose Acid Reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 Allergy alert: do not use if you are allergic to lansoprazole
 Do not use
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 Ask a doctor before use if you have
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
 ■ frequent wheezing, particularly with heartburn
 ■ frequent chest pain
 ■ unexplained weight loss
 ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 ■ If pregnant or breast-feeding, ask a health care professional before use.
 ■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Location for lot number and expiration date

8U130 FA C6-V 100418

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

■ May take 1 to 4 days for full effect

Orally Disintegrating Tablets

Melts in your mouth
 Dissolves without water



28 TABLETS

TWO 14-DAY COURSES OF TREATMENT

STANDARD FORMAT

TYPE SIZES:
 TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
 HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
 LEADING: 6.5 pt
 BULLETS: 5 pt Square with 2 M'S spacing between statements
 TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6-V 100418

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active ingredient (in each tablet) Lansoprazole 15 mg, Acid Reducer

Purpose

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)

Directions

- adults 18 years of age and older: this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- take 1 tablet before eating in the morning

14-Day Course of Treatment

- do not crush or chew tablets
- place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not take this medicine with alcohol

Repeated 14-Day Course (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information

- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F). ■ keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize malodextrin, methyl, mannitol, magnesium, microcrystalline cellulose, polyorbate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hekidma Street, Yokneam 209200, Israel
Made in Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.



Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-03

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

24 Hour
Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Acid Reducer

- May take 1 to 4 days for full effect

Orally Disintegrating Tablets

42 TABLETS THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed

HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique

SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed

TEXT: 6 pt Helvetica Neue LT Std 57 Condensed

LEADING: 6.5 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

Hairlines extend to within two spaces of the "Drug Facts" box

Made in Israel
 Manufactured by:
Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by
 GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

<p>Drug Facts (continued)</p> <p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning</p> <p>do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.</p> <p>■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other Information ■ read the directions, warnings and package insert before use ■ keep the carton and package insert. They contain important information. ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity. ■ protect product from moisture</p> <p>Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, hawthorn, hypromellose, hypromellose phthalate, maize maltoedextrin, mannitol, magnesium, microcrystalline cellulose, poly sorbate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST</p>	<p>Drug Facts</p> <p>Active ingredient (in each tablet) Lansoprazole 15 mg;Acid Reducer</p> <p>Purpose ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings ■ Allergy alert: do not use if you are allergic to lansoprazole</p> <p>Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have ■ liver disease ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating, pain spreading to arms, neck or shoulders; or lightheadedness</p> <p>Common side effects ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if ■ your heartburn continues for more than 14 days ■ you need to take this product for more than 14 months ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p> <p>If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>
--	---

Location for lot number and expiration date

8U174 FA C6-J 100418



KEEP THE CARTON AND PACKAGE INSERT.
 THEY CONTAIN IMPORTANT INFORMATION.

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC 64239-181-01

Treats Frequent Heartburn

24 Hour

Lansoprazole

Delayed Release Orally Disintegrating Tablets 15 mg

Acid Reducer

- May take 1 to 4 days for full effect

14 TABLETS ONE 14-DAY COURSE OF TREATMENT

Melts In Your Mouth

Compare to the active ingredient in Prevacid® 24 HR*

Child-resistant

Strawberry Flavor

Actual size

Melts in your mouth
 Dissolves without water

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued); 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Made in Israel
Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2069200, Israel

Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts (continued)

Active ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: do not use if you are allergic to lansoprazole

Do not use
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
 ■ liver disease
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.

Heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

Frequent chest pain
 ■ frequent wheezing, particularly with heartburn

■ unexplained weight loss
 ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST

Directions
 ■ adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning

do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.

do not use if you have
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Course (if needed)
 ■ do not take this medicine with alcohol

unless directed by a doctor
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
 ■ protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize macrodextrin, maltitol, mannitol, meglumine, microcrystalline cellulose, polyborate 80, propylene glycol, silicon dioxide, sodium stearoyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

8U130 FA C6-J 100418

Location for lot number and expiration date




Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.



NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Child-resistant

24 Hour

Lansoprazole

Delayed Release
Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

- May take 1 to 4 days for full effect

28 TABLETS

TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:

- TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- (continued) 8 pt Helvetica Neue LT Std 57 Condensed
- HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
- TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
- LEADING: 6.5 pt
- BULLETS: 5 pt Square with 2 M'S spacing between statements
- TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
- HEAVY LINES: 1.5 pt
- HAIRLINES: 0.5 pt
- Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6-J 100418

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active ingredient (in each tablet) Lansoprazole 15 mg, Acid Reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ **Do not use** if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ **Ask a doctor before use if you have** liver disease, had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ **chest pain or shoulder pain** with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
 ■ **frequent chest pain** frequent wheezing, particularly with heartburn
 ■ **unexplained weight loss** nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST

Other information
 ■ **children under 18 years of age:** ask a doctor before use. Heartburn in children unless directed by a doctor
 ■ **do not take for more than 14 days or more often than every 4 months** unless directed by a doctor
 ■ you may repeat a 14-day course every 4 months
Repeated 14-Day Course (if needed)
 ■ do not take this medicine with alcohol
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take more than 1 tablet a day
 ■ take every day for 14 days
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ **do not crush or chew tablets**
 ■ take 1 tablet before eating in the morning

14-Day Course of Treatment
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ adults 18 years of age and older

Directions
 ■ take 1 tablet before eating in the morning

Drug Facts (continued)

Other information
 ■ keep the carton and package insert. They contain important information.
 ■ protect product from moisture
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hydroxypropyl methylcellulose, milled polyborate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-03

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Child-resistant

24 Hour
Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

- May take 1 to 4 days for full effect

42 TABLETS THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box



Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel
Made in Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Drug Facts

Active ingredient (in each tablet)
Lansoprazole 15 mg.....Acid Reducer

Use
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: do not use if you are allergic to lansoprazole

Do not use
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
■ liver disease
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness

Frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260. weekdays 7:30 AM to 5:00 PM EST

Other information
■ read the directions, warnings and package insert before use
■ keep the carton and package insert. They contain important information.
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
■ protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize starch, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucrose, sugar spheres, talc, titanium dioxide, triethyl citrate

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ take 1 tablet before eating in the morning
■ do not crush or chew tablets
■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
■ take every day for 14 days
■ do not use for more than 14 days unless directed by your doctor
■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor


■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15mg

8U130 FA C6-A 100418

Location for lot number and expiration date

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02


Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Value 2 Pack

24 Hour
Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6-B 100418

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ Ask a doctor before use if you have liver diseases
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
 ■ frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260. weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)

Directions
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
 ■ take 1 tablet before eating in the morning

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
 ■ protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyorbital, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate, poly sorbate 80, propylene glycol, silicon dioxide, sodium stearyl fumarate, naldexidine, methylol, mannitol, magnesium, microcrystalline cellulose, crospovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyorbital, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Manufactured by:
 Decel Pharma Technologies Ltd
 10 Hakdama Street, Yokneam 2069200, Israel
 Made in Israel
 *This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.



Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-03

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Value 3 Pack

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

- May take 1 to 4 days for full effect

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



STANDARD FORMAT

TYPE SIZES:
 TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
 HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
 LEADING: 6.5 pt
 BULLETS: 5 pt Square with 2 M'S spacing between statements
 TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Manufactured by:
Dexcel Pharma Technologies Ltd
10 Hakdima Street, Yokneam 2069200, Israel

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active Ingredient (in each tablet)
Lansoprazole 15 mg
Acid Reducer

Purpose
Acid Reducer

Use
■ Treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: do not use if you are allergic to lansoprazole

Do not use
■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
■ liver disease
■ had heartburn over 3 months. This may be a sign of a more serious condition.

chest pain
■ frequent wheezing, particularly with heartburn
■ frequent weight loss
■ unexplained weight loss

Ask a doctor or pharmacist before use if you are taking a prescription drug.
Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ take 1 tablet before eating in the morning

Warnings
■ do not crush or chew tablets
■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.

Do not use
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Course (if needed)
■ do not take this medicine with alcohol

Other information
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information
■ read the directions, warnings and package insert before use
■ keep the carton and package insert. They contain important information.
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
■ protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, flavor, hypromellose, hypromellose phthalate, maize polyborate 80, propylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

8U130 FA C6-K 100418

Location for lot number and expiration date



Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Buy One 14 Count, Get One 14 Count Free!

24 Hour
Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:

- TITLE:** 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- TITLE (continued):** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued)
- 8 pt Helvetica Neue LT Std 57 Condensed**
- HEADINGS:** 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
- SUBHEADINGS:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- TEXT:** 6 pt Helvetica Neue LT Std 57 Condensed
- LEADING:** 6.5 pt
- BULLETS:** 5 pt Square with 2 M'S spacing between statements
- TELEPHONE #:** 6 pt Helvetica Neue LT Std 77 Bold Condensed
- HEAVY LINES:** 1.5 pt
- HAIRLINES:** 0.5 pt
- Hairlines extend to within two spaces of the "Drug Facts" box**



Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdama Street, Yokneam 2069200, Israel
 *This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Drug Facts
 Active Ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 Allergy alert: do not use if you are allergic to lansoprazole

Do not use
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
 ■ liver disease
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST

Other information
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F). ■ keep product out of high heat and humidity
 ■ protect product from moisture

Inactive ingredients
 ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, flavor, hypromellose, hypromellose phthalate, maize starch, mannitol, microcrystalline cellulose, polyethylene glycol, polypropylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Directions
 (continued)

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

8U130 FA C6-L 100418

Location for lot number and expiration date

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

100% More Free!
 14 Free Tablets!

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
 ■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:
 TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
 HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
 LEADING: 6.5 pt
 BULLETS: 5 pt Square with 2 M'S spacing between statements
 TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdama Street, Yokneam 2069200, Israel
 *This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.


Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15mg

Drug Facts
 Active ingredient (in each tablet) Lansoprazole 15 mg, Acid Reducer
Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
Warnings
 Allergy alert: do not use if you are allergic to lansoprazole
 Do not use
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 Ask a doctor before use if you have
 ■ liver disease
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting ■ stomach pain
 Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain
 If pregnant or breast-feeding, ask a health care professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
Other information
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
 ■ protect product from moisture
Ingredients
 lansoprazole, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize starch, polyethylene glycol, polyethylene glycol, silicon dioxide, sodium stearoyl fumarate, maledextrin, mannitol, magnesium, microcrystalline cellulose, methylcellulose, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate
Directions
 (continued)
 14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 Repeated 14-Day Course (if needed)
 ■ you repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
Other information
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
 ■ protect product from moisture
Ingredients
 lansoprazole, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize starch, polyethylene glycol, polyethylene glycol, silicon dioxide, sodium stearoyl fumarate, maledextrin, mannitol, magnesium, microcrystalline cellulose, methylcellulose, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate
Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST

8U130 FA C6-N 100418

Location for lot number and expiration date

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Bonus Size!
 14 Tablets Free

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
 ■ May take 1 to 4 days for full effect


28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



Actual size

STANDARD FORMAT
TYPE SIZES:
 TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued) 8 pt Helvetica Neue LT Std 57 Condensed
 HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
 LEADING: 6.5 pt
 BULLETS: 5 pt Square with 2 M'S spacing between statements
 TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 Made in Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Drug Facts
 Active ingredient (in each tablet)
 Lansoprazole 15 mg.....Acid Reducer

Use
 ■ Treats frequent heartburn (occurs 2 or more days a week)
 ■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 Allergy alert: do not use if you are allergic to lansoprazole
 Do not use
 ■ If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 Ask a doctor before use if you have
 ■ Liver disease
 ■ Had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ Heartburn with lightheadedness, sweating or dizziness
 ■ Chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
 ■ Frequent wheezing, particularly with heartburn
 ■ Unexplained weight loss
 ■ Nausea or vomiting
 ■ Stomach pain
 Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ Your heartburn continues or worsens
 ■ You need to take this product for more than 14 days
 ■ You need to take more than 1 course of treatment every 4 months
 ■ You get diarrhea
 ■ You develop a rash or joint pain
 If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)
Directions
 ■ Adults 18 years of age and older
 ■ This product is to be used once a day (every 24 hours), every day for 14 days
 ■ It may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ Take 1 tablet before eating in the morning
 ■ Do not crush or chew tablets
 ■ Place the tablet on tongue, tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ Take every day for 14 days
 ■ Do not take for more than 14 days unless directed by your doctor.
 ■ Do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ You may repeat a 14-day course every 4 months
 ■ Do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information
 ■ Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
 ■ Read the directions, warnings and package insert before use
 ■ Keep the carton and package insert. They contain important information.
 ■ Store at 20-25°C (68-77°F) ■ Keep product out of high heat and humidity
 ■ Protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize starch, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, malleodextrin, mannitol, magnesium, microcrystalline cellulose, polyborate 80, propylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

8U130 FA C6-P 100418

Location for lot number and expiration date



Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Bonus Size! 14 Free

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
 ■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



Actual size

STANDARD FORMAT
TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.
 Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel
 Manufactured by:
 Made in Israel

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

<p>Drug Facts Active ingredient (in each tablet) Lansoprazole 15 mg Acid Reducer</p>	<p>Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p>
<p>Warnings ■ Allergy alert: do not use if you are allergic to lansoprazole ■ Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</p>	<p>Ask a doctor before use if you have ■ liver disease ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p>
<p>Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</p>	<p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p>
<p>Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p>	<p>If pregnant or breast-feeding, ask a health care professional before use. ■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>
<p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours</p>	<p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol ■ do not take this medicine with alcohol</p>
<p>14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p>	<p>Other information ■ read the directions, warnings and package insert before use ■ keep the carton and package insert. They contain important information. ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity ■ protect product from moisture</p>
<p>Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopovidone, flavor, hypromellose, hypromellose phthalate, maize maltoдекtrin, methylcellulose, mannitol, magnesium, microcrystalline cellulose, polyorbate 80, propylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p>	<p>Questions or comments? 1-800-719-9260, weekdays 7:30 AM to 5:00 PM EST</p>

8U130 FA C6-R 100418

Location for lot number and expiration date

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Bonus! + 14 Free

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
 ■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:
 TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
 HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
 LEADING: 6.5 pt
 BULLETS: 5 pt Square with 2 M'S spacing between statements
 TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box



Manufactured by:
Cexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 Made in Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Drug Facts
Active ingredient (in each tablet)
 Lansoprazole 15 mg.....Acid Reducer

Use
 ■ Treats frequent heartburn (occurs 2 or more days a week)
 ■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ Ask a doctor before use if you have:
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Location for lot number and expiration date

8U130 FA C6-S 100418

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.



NDC 64239-181-02

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Bonus! 14 Tablets FREE

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer
 ■ May take 1 to 4 days for full effect

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



Actual size

STANDARD FORMAT
TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6-M 100418

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer

Use
 ■ Treats frequent heartburn (occurs 2 or more days a week)
 ■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ Do not use if you are allergic to lansoprazole

Do not use
 ■ If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
 ■ liver disease
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

Frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260. weekdays 7:30 AM to 5:00 PM EST

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyborate 80, propylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Other information
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
 ■ protect product from moisture

Directions
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol

14-Day Course of Treatment
 ■ take every day for 14 days
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ do not crush or chew tablets
 ■ take 1 tablet before eating in the morning

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Manufactured by:
Cexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2069200, Israel
 Made in Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.



Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-03

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Bonus Pack: 28 + 14 Free

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg



Strawberry Flavor

Acid Reducer

- May take 1 to 4 days for full effect

42 TABLETS THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6-N 100418

Made in Israel
 Manufactured by:
 Cexel Pharma Technologies Ltd
 10 Hakdama Street, Yokneam 2069200, Israel
 *This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Drug Facts
Active ingredient (in each tablet) Lansoprazole 15 mg, Acid Reducer
Purpose
Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
Warnings
 ■ Allergy alert: do not use if you are allergic to lansoprazole
 ■ Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ Ask a doctor before use if you have:
 ■ liver disease
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with **lightheadedness, sweating or dizziness**
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
 ■ frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain
 ■ Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain
If pregnant or breast-feeding, ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
 5:00 PM EST
Questions or comments? 1-800-719-9260, weekdays 7:30 AM to

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Drug Facts (continued)
Directions
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
14-Day Treatment
 ■ take 1 tablet before eating in the morning
 ■ do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol
Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
Other information
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
 ■ protect product from moisture
Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize starch, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-03

Treats Frequent Heartburn

MELTech™
 Melts In Your Mouth

Compare to the active ingredient in Prevacid® 24 HR*

Bonus Size!
14 Tablets Free

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15 mg

Acid Reducer
 ■ May take 1 to 4 days for full effect

42 TABLETS THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water

Strawberry Flavor

Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
TITLE (continued): 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6-R 100418

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Drug Facts

Active ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ Ask a doctor before use if you have liver diseases
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
 ■ frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260. weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)

Directions
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
 ■ take 1 tablet before eating in the morning

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
 ■ read the directions, warnings and package insert before use
 ■ keep the carton and package insert. They contain important information.
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
 ■ protect product from moisture

Inactive ingredients ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyorbital, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate, poly sorbate 80, propylene glycol, silicon dioxide, sodium stearyl fumarate, naldexidine, methylol, mannitol, magnesium, microcrystalline cellulose, crospovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyorbital, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

NDC 64239-181-03

Treats Frequent Heartburn

MELTech™
Melts In Your Mouth

Compare to the active ingredient in Prevacid® 24 HR*

Bonus! + 14 Free

24 Hour

Lansoprazole
Delayed Release
Orally Disintegrating Tablets 15 mg

Acid Reducer
 ■ May take 1 to 4 days for full effect

42 TABLETS THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water

Strawberry Flavor

15 Actual size

STANDARD FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued); 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Location for lot number and expiration date

8U155 FA C6-S 100418

Made in Israel
 Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Hakidma Street, Yokneam 2069200, Israel

*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Prevacid® 24 HR.



Drug Facts
 Active ingredient (in each tablet)
 Lansoprazole 15 mg.....Acid Reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ Ask a doctor before use if you have:
 ■ liver disease
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath; swelling; pain spreading to arms, neck or shoulders; or rightheadedness
 ■ frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Drug Facts (continued)
Directions
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning

Do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Inactive ingredients
 ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolvone, flavoc, hypromellose, hypromellose phthalate, maize polyorbate 80, propylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15mg



Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.



NDC 64239-181-03

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR*

Bonus! 14 Tablets FREE

24 Hour
Lansoprazole
 Delayed Release
 Orally Disintegrating Tablets 15mg



Strawberry Flavor

Acid Reducer

- May take 1 to 4 days for full effect

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

Melts in your mouth
 Dissolves without water



Actual size

STANDARD FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued); 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
 (continued) 8 pt Helvetica Neue LT Std 57 Condensed
HEADINGS: 8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
SUBHEADINGS: 6 pt Helvetica Neue LT Std 77 Bold Condensed
TEXT: 6 pt Helvetica Neue LT Std 57 Condensed
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6 pt Helvetica Neue LT Std 77 Bold Condensed
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt
 Hairlines extend to within two spaces of the "Drug Facts" box

Treats Frequent Heartburn

Lansoprazole

Delayed Release

Orally Disintegrating Tablets 15 mg

acid reducer

- May take 1 to 4 days for full effect

Please read the entire package insert before taking
Lansoprazole Delayed Release Orally Disintegrating Tablets.
Save for future reference.

How Lansoprazole delayed release orally disintegrating tablets Treats Your Frequent Heartburn

Lansoprazole delayed release orally disintegrating tablets stops acid production at the source – the **pumps** that release acid into the stomach. Lansoprazole delayed release orally disintegrating tablet is taken once a day (every 24 hours), every day for 14 days.

What You Can Expect When Taking Lansoprazole delayed release orally disintegrating tablets

Frequent heartburn can occur anytime during the 24-hour period (day or night). Take Lansoprazole delayed release orally disintegrating tablets in the morning before eating. Lansoprazole is clinically proven to treat frequent heartburn. Although some people get complete relief of symptoms within 24 hours, it may take 1 to 4 days for full effect. Make sure you take Lansoprazole delayed release orally disintegrating tablets every day for 14 days to treat your frequent heartburn.

Who Should Take Lansoprazole delayed release orally disintegrating tablets

Adults (18 years and older) with **frequent heartburn** – when you have heartburn 2 or more days a week.

Who Should NOT Take Lansoprazole delayed release orally disintegrating tablets

People who have one episode of heartburn a week or less, or who want immediate relief of heartburn.

How to Take Lansoprazole delayed release orally disintegrating tablets

14-DAY Course of Treatment

- Take 1 tablet before eating in the morning.
- **Do not crush or chew tablets.**
- Place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
- Take every day for 14 days.
- Do not take more than 1 tablet a day.
- Do not use for more than 14 days unless directed by your doctor.
- Do not take this medicine with alcohol.

When to Take Lansoprazole delayed release orally disintegrating tablets Again

You may repeat a 14-day course of therapy every 4 months.

When to Talk to Your Doctor

Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.



Warnings and When to Ask Your Doctor

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Clinical studies prove lansoprazole effectively treats frequent heartburn

In three clinical studies, lansoprazole was shown to be significantly better than placebo in treating frequent heartburn.

How Lansoprazole delayed release orally disintegrating tablets is Sold

This medicine is available in 14 tablet, 28 tablet and 42 tablet sizes. These sizes contain one, two and three 14-day courses of treatment, respectively. Do not use for more than 14 days in a row unless directed by your doctor. For the 28 count (two 14-day courses) and the 42 count (three 14-day courses), you may repeat a 14-day course every 4 months.

For Questions or Comments About Lansoprazole delayed release orally disintegrating tablets

Call 1-800-719-9260


weekdays 7:30 AM to 5:00 PM EST

Made in Israel

Manufactured by:

Dexcel Pharma Technologies Ltd
10 Hakidma Street, Yokneam 2069200, Israel

: 80100 00 J3 100418



This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
04/09/2019 02:42:39 PM