

NEUPOGEN® (Filgrastim)

DESCRIPTION

Filgrastim is a human granulocyte colony stimulating factor (G-CSF), produced by recombinant DNA technology. NEUPOGEN® is the Amgen Inc. trademark for Filgrastim, which has been selected as the name for recombinant methionyl human granulocyte colony stimulating factor (r-metHuG-CSF).

NEUPOGEN is a 175 amino acid protein manufactured by recombinant DNA technology.¹ NEUPOGEN is produced by *Escherichia coli* (*E coli*) bacteria into which has been inserted the human granulocyte colony stimulating factor gene. NEUPOGEN has a molecular weight of 18,800 daltons. The protein has an amino acid sequence that is identical to the natural sequence predicted from human DNA sequence analysis, except for the addition of an N-terminal methionine necessary for expression in *E coli*. Because NEUPOGEN is produced in *E coli*, the product is nonglycosylated and thus differs from G-CSF isolated from a human cell.

NEUPOGEN is a sterile, clear, colorless, preservative-free liquid for parenteral administration. Each single-use vial of NEUPOGEN contains 300 mcg/mL of Filgrastim at a specific activity of $1.0 \pm 0.6 \times 10^8$ U/mg, (as measured by a cell mitogenesis assay). The product is formulated in a 10 mM sodium acetate buffer at pH 4.0, containing 5% sorbitol, and 0.004% Tween® 80. The quantitative composition (per mL) of NEUPOGEN is:

Filgrastim	300 mcg
Acetate	0.59 mg
Sorbitol	50.0 mg
Tween® 80	0.004%
Sodium	0.035 mg
Water for Injection USP q.s. ad	1.0 mL

CLINICAL PHARMACOLOGY

Colony Stimulating Factors

Colony stimulating factors are glycoproteins which act on hematopoietic cells by binding to specific cell surface receptors and stimulating proliferation, differentiation commitment, and some end-cell functional activation.

Endogenous G-CSF is a lineage specific colony stimulating factor which is produced by monocytes, fibroblasts, and endothelial cells. G-CSF regulates the production of neutrophils within the bone marrow and affects neutrophil progenitor proliferation,^{2,3} differentiation,^{2,4} and selected end-cell functional activation (including enhanced phagocytic ability,⁵ priming of the cellular metabolism associated with respiratory burst,⁶ antibody dependent killing,⁷ and the increased expression of some functions associated with cell surface antigens⁸). G-CSF is not species specific and has been shown to have minimal direct in vivo or in vitro effects on the production of hematopoietic cell types other than the neutrophil lineage.

Preclinical Experience

Filgrastim was administered to monkeys, dogs, hamsters, rats, and mice as part of a preclinical toxicology program which included single-dose acute, repeated-dose subacute, subchronic, and chronic studies. Single-dose administration of Filgrastim by the oral, intravenous (IV), subcutaneous (SC), or intraperitoneal (IP) routes resulted in no significant toxicity in mice, rats, hamsters, or monkeys. Although no deaths were observed in mice, rats, or monkeys at dose levels up to 3450 mcg/kg or in hamsters using single doses up to approximately 860 mcg/kg, deaths were observed in a subchronic (13-week) study in monkeys. In this study, evidence of neurological symptoms was seen in monkeys treated with doses of Filgrastim greater than 1150 mcg/kg/day for up to 18 days. Deaths were seen in five of the eight treated animals and were associated with 15- to 28-fold increases in peripheral leukocyte counts, and neutrophil-infiltrated hemorrhagic foci were seen in both the cerebrum and cerebellum. In contrast, no monkeys died following 13 weeks of daily IV administration of Filgrastim at a dose level of 115 mcg/kg. In an ensuing 52-week study, one 115 mcg/kg dose female monkey died after 18 weeks of daily IV administration of Filgrastim. Death was attributed to cardiopulmonary insufficiency.

In subacute, repeated-dose studies, changes observed were attributable to the expected pharmacological actions of Filgrastim (ie, dose-dependent increases in white cell counts, increased circulating segmented neutrophils, and increased myeloid:erythroid ratio in bone marrow). In all species, histopathologic examination of the liver and spleen revealed evidence of ongoing extramedullary granulopoiesis; increased spleen weights were seen in all species and appeared to be dose-related. A dose-dependent increase in serum alkaline

phosphatase was observed in rats, and may reflect increased activity of osteoblasts and osteoclasts. Changes in serum chemistry values were reversible following discontinuation of treatment.

In rats treated at doses of 1150 mcg/kg/day for 4 weeks (5 of 32 animals) and for 13 weeks at doses of 100 mcg/kg/day (4 of 32 animals) and 500 mcg/kg/day (6 of 32 animals), articular swelling of the hind legs was observed. Some degree of hind leg dysfunction was also observed; however, symptoms reversed following cessation of dosing. In rats, osteoclasts and osteoanagenesis were found in the femur, humerus, coccyx, and hind legs (where they were accompanied by synovitis) after IV treatment for 4 weeks (115 to 1150 mcg/kg/day), and in the sternum after IV treatment for 13 weeks (115 to 575 mcg/kg/day). These effects reversed to normal within 4 to 5 weeks following cessation of treatment.

In the 52-week chronic, repeated-dose studies performed in rats (IP injection up to 57.5 mcg/kg/day), and cynomolgus monkeys (IV injection of up to 115 mcg/kg/day), changes observed were similar to those noted in the subacute studies. Expected pharmacological actions of Filgrastim included dose-dependent increases in white cell counts, increased circulating segmented neutrophils and alkaline phosphatase levels, and increased myeloid:erythroid ratios in the bone marrow. Decreases in platelet counts were also noted in primates. In no animals tested were hemorrhagic complications observed. Rats displayed dose-related swelling of the hind limb, accompanied by some degree of hind limb dysfunction; osteopathy was noted microscopically. Enlarged spleens (both species) and livers (monkeys), reflective of ongoing extramedullary granulopoiesis, as well as myeloid hyperplasia of the bone marrow, were observed in a dose-dependent manner.

Pharmacologic Effects of NEUPOGEN

In phase 1 studies involving 96 patients with various nonmyeloid malignancies, NEUPOGEN administration resulted in a dose-dependent increase in circulating neutrophil counts over the dose range of 1 to 70 mcg/kg/day.⁹⁻¹¹ This increase in neutrophil counts was observed whether NEUPOGEN was administered IV (1 to 70 mcg/kg twice daily),⁹ SC (1 to 3 mcg/kg once daily),¹¹ or by continuous SC infusion (3 to 11 mcg/kg/day).¹⁰ With discontinuation of NEUPOGEN therapy, neutrophil counts returned to baseline, in most cases within 4 days. Isolated neutrophils displayed normal phagocytic (measured by zymosan-stimulated chemoluminescence) and chemotactic [measured by migration under agarose using N-formyl-methionyl-leucyl-phenylalanine (fMLP) as the chemotaxin] activity in vitro.

The absolute monocyte count was reported to increase in a dose-dependent manner in most patients receiving NEUPOGEN, however, the percentage of monocytes in the differential count remained within the normal range. In all studies to date, absolute counts of both eosinophils and basophils did not change and were within the normal range following administration of NEUPOGEN. Increases in lymphocyte counts following

NEUPOGEN administration have been reported in some normal subjects and cancer patients.

White blood cell (WBC) differentials obtained during clinical trials have demonstrated a shift towards earlier granulocyte progenitor cells (left shift), including the appearance of promyelocytes and myeloblasts, usually during neutrophil recovery following the chemotherapy-induced nadir. In addition, Dohle bodies, increased granulocyte granulation, as well as hypersegmented neutrophils have been observed. Such changes were transient, and were not associated with clinical sequelae nor were they necessarily associated with infection.

Pharmacokinetics

Absorption and clearance of NEUPOGEN follows first-order pharmacokinetic modeling without apparent concentration dependence. A positive linear correlation occurred between the parenteral dose and both the serum concentration and area under the concentration-time curves. Continuous IV infusion of 20 mcg/kg of NEUPOGEN over 24 hours resulted in mean and median serum concentrations of approximately 48 and 56 ng/mL, respectively. Subcutaneous administration of 3.45 mcg/kg and 11.5 mcg/kg resulted in maximum serum concentrations of 4 and 49 ng/mL, respectively, within 2 to 8 hours. The volume of distribution averaged 150 mL/kg in both normal subjects and cancer patients. The elimination half-life, in both normal subjects and cancer patients, was approximately 3.5 hours. Clearance rates of NEUPOGEN were approximately 0.5 to 0.7 mL/minute/kg. Single parenteral doses or daily IV doses, over a 14-day period, resulted in comparable half-lives. The half-lives were similar for IV administration (231 minutes, following doses of 34.5 mcg/kg) and for SC administration (210 minutes, following NEUPOGEN doses of 3.45 mcg/kg). Continuous 24-hour IV infusions of 20 mcg/kg over an 11- to 20-day period produced steady-state serum concentrations of NEUPOGEN with no evidence of drug accumulation over the time period investigated.

CLINICAL EXPERIENCE

Cancer Patients Receiving Myelosuppressive Chemotherapy

NEUPOGEN has been shown to be safe and effective in accelerating the recovery of neutrophil counts following a variety of chemotherapy regimens. In a phase 3 clinical trial in small cell lung cancer, patients received SC administration of NEUPOGEN (4 to 8 mcg/kg/day, days 4 to 17) or placebo. In this study, the benefits of NEUPOGEN therapy were shown to be prevention of infection as manifested by febrile neutropenia, decreased hospitalization, and decreased IV antibiotic usage. No difference in survival or disease progression was demonstrated.

In the phase 3, randomized, double-blind, placebo-controlled trial conducted in patients

with small cell lung cancer, patients were randomized to receive NEUPOGEN (n = 99) or placebo (n = 111) starting on day 4, after receiving standard dose chemotherapy with cyclophosphamide, doxorubicin, and etoposide. A total of 210 patients were evaluated for efficacy and 207 evaluated for safety. Treatment with NEUPOGEN resulted in a clinically and statistically significant reduction in the incidence of infection, as manifested by febrile neutropenia; the incidence of at least one infection over all cycles of chemotherapy was 76% (84/111) for placebo-treated patients, versus 40% (40/99) for NEUPOGEN-treated patients ($p < 0.001$). The following secondary analyses were also performed. The requirements for in-patient hospitalization and antibiotic use were also significantly decreased during the first cycle of chemotherapy; incidence of hospitalization was 69% (77/111) for placebo-treated patients in cycle 1, versus 52% (51/99) for NEUPOGEN-treated patients ($p = 0.032$). The incidence of IV antibiotic usage was 60% (67/111) for placebo-treated patients in cycle 1, versus 38% (38/99) for NEUPOGEN-treated patients ($p = 0.003$). The incidence, severity, and duration of severe neutropenia [absolute neutrophil count (ANC) $< 500/\text{mm}^3$] following chemotherapy were all significantly reduced. The incidence of severe neutropenia in cycle 1 was 84% (83/99) for patients receiving NEUPOGEN versus 96% (106/110) for patients receiving placebo ($p = 0.004$). Over all cycles, patients randomized to NEUPOGEN had a 57% (286/500 cycles) rate of severe neutropenia versus 77% (416/543 cycles) for patients randomized to placebo. The median duration of severe neutropenia in cycle 1 was reduced from 6 days (range 0 to 10 days) for patients receiving placebo to 2 days (range 0 to 9 days) for patients receiving NEUPOGEN ($p < 0.001$). The mean duration of neutropenia in cycle 1 was 5.64 ± 2.27 days for patients receiving placebo versus 2.44 ± 1.90 days for patients receiving NEUPOGEN. Over all cycles, the median duration of neutropenia was 3 days for patients randomized to placebo versus 1 day for patients randomized to NEUPOGEN. The median severity of neutropenia (as measured by ANC nadir) was $72/\text{mm}^3$ (range $0/\text{mm}^3$ to $7912/\text{mm}^3$) in cycle 1 for patients receiving NEUPOGEN versus $38/\text{mm}^3$ (range $0/\text{mm}^3$ to $9520/\text{mm}^3$) for patients receiving placebo ($p = 0.012$). The mean severity of neutropenia in cycle 1 was $496/\text{mm}^3 \pm 1382/\text{mm}^3$ for patients receiving NEUPOGEN versus $204/\text{mm}^3 \pm 953/\text{mm}^3$ for patients receiving placebo. Over all cycles, the ANC nadir for patients randomized to NEUPOGEN was $403/\text{mm}^3$, versus $161/\text{mm}^3$ for patients randomized to placebo. Administration of NEUPOGEN resulted in an earlier ANC nadir following chemotherapy than was experienced by patients receiving placebo (day 10 vs day 12). NEUPOGEN was well tolerated when given SC daily at doses of 4 to 8 mcg/kg for up to 14 consecutive days following each cycle of chemotherapy (see ADVERSE REACTIONS).

Several other phase 1/2 studies, which did not directly measure the incidence of infection, but which did measure increases in neutrophils, support the efficacy of NEUPOGEN. The regimens are presented to provide some background on the clinical experience with NEUPOGEN. No claim regarding the safety or efficacy of the chemotherapy regimens is made. The effects of NEUPOGEN on tumor growth or on the anti-tumor activity of the

chemotherapy were not assessed. The doses of NEUPOGEN used in these studies are considerably greater than those found to be effective in the phase 3 study described above. Such phase 1/2 studies are summarized in the following table.

Type of Malignancy	Regimen	Chemotherapy Dose	No. Pts.	Trial Phase	NEUPOGEN Daily Dosage ^a
Small Cell Lung Cancer	Cyclophosphamide Doxorubicin Etoposide	1 g/m ² /day 50 mg/m ² /day 120 mg/m ² /day x 3 q 21 days	210	3	4-8 mcg/kg SC days 4-17
Small Cell Lung Cancer ¹¹	Ifosfamide Doxorubicin Etoposide Mesna	5 g/m ² /day 50 mg/m ² /day 120 mg/m ² /day x 3 8 g/m ² /day q 21 days	12	1/2	5.75-46 mcg/kg IV days 4-17
Urothelial Cancer ¹²	Methotrexate Vinblastine Doxorubicin Cisplatin	30 mg/m ² /day x 2 3 mg/m ² /day x 2 30 mg/m ² /day 70 mg/m ² /day q 28 days	40	1/2	3.45-69 mcg/kg IV days 4-11
Various Nonmyeloid Malignancies ¹³	Cyclophosphamide Etoposide Cisplatin	2.5 g/m ² /day x 2 500 mg/m ² /day x 3 50 mg/m ² /day x 3 q 28 days	18	1/2	23-69 mcg/kg ^b IV days 8-28
Breast/Ovarian Cancer ¹⁴	Doxorubicin ^c	75 mg/m ² 100 mg/m ² 125 mg/m ² 150 mg/m ² q 14 days	21	2	11.5 mcg/kg IV days 2-9 5.75 mcg/kg IV days 10-12
Neuroblastoma	Cyclophosphamide Doxorubicin Cisplatin	150 mg/m ² x 7 35 mg/m ² 90 mg/m ² q 28 days (cycles 1,3,5) ^d	12	2	5.45-17.25 mcg/kg SC days 6-19
^a NEUPOGEN doses were those that accelerated neutrophil production. Doses which provided no additional acceleration beyond that achieved at the next lower dose are not reported. ^b Lowest dose(s) tested in the study. ^c Patients received doxorubicin at either 75, 100, 125, or 150 mg/m ² . ^d Cycles 2,6 = cyclophosphamide 150 mg/m ² x 7 and etoposide 280 mg/m ² x 3. Cycle 4 = cisplatin 90 mg/m ² x 1 and etoposide 280 mg/m ² x 3.					

Patients With Acute Myeloid Leukemia Receiving Induction or Consolidation Chemotherapy

In a randomized, double-blind, placebo-controlled, multi-center, phase 3 clinical trial, 521 patients (median age 54, range 16 to 89 years) were treated for de novo acute myeloid leukemia (AML). Following a standard induction chemotherapy regimen comprising daunorubicin, cytosine arabinoside, and etoposide¹⁷ (DAV 3+7+5), patients received either NEUPOGEN at 5 mcg/kg/day or placebo, SC, from 24 hours after the last dose of chemotherapy until neutrophil recovery (ANC 1000/mm³ for 3 consecutive days or 10,000/mm³ for 1 day) or for a maximum of 35 days.

Treatment with NEUPOGEN significantly reduced the median time to ANC recovery and the median duration of fever, antibiotic use, and hospitalization following induction chemotherapy. In the NEUPOGEN-treated group, the median time from initiation of chemotherapy to ANC recovery (ANC \geq 500/mm³) was 20 days (vs 25 days in the control group, $p = 0.0001$), the median duration of fever was reduced by 1.5 days ($p = 0.009$), and there were statistically significant reductions in the durations of IV antibiotic use and hospitalization. During consolidation therapy (DAV 2+5+5), patients treated with NEUPOGEN also experienced significant reductions in the incidence of severe neutropenia, time to neutrophil recovery, the incidence and duration of fever, and in the durations of IV antibiotic use and hospitalization. Patients treated with a further course of standard (DAV 2+5+5) or high-dose cytosine arabinoside consolidation also experienced significant reductions in the duration of neutropenia.

There were no statistically significant differences between NEUPOGEN and placebo groups in complete remission rate (69% NEUPOGEN vs 68% placebo, $p = 0.77$), disease-free survival [median 342 days NEUPOGEN ($n = 178$), 322 days placebo ($n = 177$), $p = 0.99$], time to progression of all randomized patients (median 165 days NEUPOGEN, 186 days placebo, $p = 0.87$), or overall survival (median 380 days NEUPOGEN, 425 days placebo, $p = 0.83$).

Cancer Patients Receiving Bone Marrow Transplant

In two separate randomized, controlled trials, patients with Hodgkin's disease (HD) and non-Hodgkin's lymphoma (NHL) were treated with myeloablative chemotherapy and autologous bone marrow transplantation (ABMT). In one study ($n = 54$), NEUPOGEN was administered at doses of 10 or 30 mcg/kg/day; a third treatment group in this study received no NEUPOGEN. A statistically significant reduction in the median number of days of severe neutropenia (ANC $<$ 500/mm³) occurred in the NEUPOGEN-treated group versus the control group [23 days in the control group, 11 days in the 10 mcg/kg/day group, and 14 days in the 30 mcg/kg/day group, (11 days in the combined treatment groups, $p = 0.004$)]. In the second study ($n = 44$, 43 patients evaluable), NEUPOGEN was administered at doses of 10 or 20 mcg/kg/day; a third treatment group in this study received no NEUPOGEN. A statistically significant reduction in the median number of

days of severe neutropenia occurred in the NEUPOGEN-treated group versus the control group (21.5 days in the control group and 10 days in both treatment groups, $p < 0.001$). The number of days of febrile neutropenia was also reduced significantly in this study [13.5 days in the control group, 5 days in the 10 mcg/kg/day group, and 5.5 days in the 20 mcg/kg/day group, (5 days in the combined treatment groups, $p < 0.0001$)]. Reductions in the number of days of hospitalization and antibiotic use were also seen, although these reductions were not statistically significant. There were no effects on red blood cell or platelet levels.

In a randomized, placebo-controlled trial, 70 patients with myeloid and nonmyeloid malignancies were treated with myeloablative therapy and allogeneic bone marrow transplant followed by 300 mcg/m²/day of a Filgrastim product. A statistically significant reduction in the median number of days of severe neutropenia occurred in the treated group versus the control group (19 days in the control group and 15 days in the treatment group, $p < 0.001$) and time to recovery of ANC to $\geq 500/\text{mm}^3$ (21 days in the control group and 16 days in the treatment group, $p < 0.001$).

In three nonrandomized studies ($n = 119$), patients received ABMT and treatment with NEUPOGEN. One study ($n = 45$) involved patients with breast cancer and malignant melanoma. A second study ($n = 39$) involved patients with HD. The third study ($n = 35$) involved patients with NHL, acute lymphoblastic leukemia (ALL), and germ cell tumor. In these studies, the recovery of the ANC to $\geq 500/\text{mm}^3$ ranged from a median of 11.5 to 13 days.

None of the conditioning regimens used in the ABMT studies included radiation therapy.

While these studies were not designed to compare survival, this information was collected and evaluated. The overall survival and disease progression of patients receiving NEUPOGEN in these studies were similar to those observed in the respective control groups and to historical data.

Peripheral Blood Progenitor Cell Collection and Therapy in Cancer Patients

All patients in the Amgen-sponsored trials received a similar mobilization/collection regimen: NEUPOGEN was administered for 6 to 7 days, with an apheresis procedure on days 5, 6, and 7 (except for a limited number of patients receiving apheresis on days 4, 6, and 8). In a non-Amgen-sponsored study, patients underwent mobilization to a target number of mononuclear cells (MNC), with apheresis starting on day 5. There are no data on the mobilization of peripheral blood progenitor cells (PBPC) after days 4 to 5 that are not confounded by leukapheresis.

Mobilization: Mobilization of PBPC was studied in 50 heavily pretreated patients (median number of prior cycles = 9.5) with NHL, HD, or ALL (Amgen study 1). CFU-

GM was used as the marker for engraftable PBPC. The median CFU-GM level on each day of mobilization was determined from the data available (CFU-GM assays were not obtained on all patients on each day of mobilization). These data are presented below.

The data from Amgen study 1 were supported by data from Amgen study 2 in which 22 pretreated breast cancer patients (median number of prior cycles = 3) were studied. Both the CFU-GM and CD34⁺ cells reached a maximum on day 5 at > 10-fold over baseline and then remained elevated with leukapheresis.

Progenitor Cell Levels in Peripheral Blood by Mobilization Day						
	Overall Study 1 CFU-GM/mL		Study 2 CFU-GM/mL		Study 2 CD34⁺ (x 10⁴/mL)	
	No. Samples	Median (25%-75%)	No. Samples	Median (25%-75%)	No. Samples	Median (25%-75%)
Day 1	11	18 (13-62)	20	42 (15-151)	20	0.13 (0.02-0.66)
Day 2	7	22 (3-61)	n/a	n/a	n/a	n/a
Day 3	10	138 (39-364)	n/a	n/a	n/a	n/a
Day 4	18	365 (158-864)	18	576 (108-1819)	17	2.11 (0.58-3.93)
Day 5	36	781 (391-1608)	21	960 (72-1677)	22	3.16 (1.08-6.11)
Day 6	46	505 (199-1397)	22	756 (70-3486)	22	2.67 (1.09-4.40)
Day 7	37	333 (111-938)	22	597 (118-2009)	21	2.64 (0.78-4.22)
Day 8	15	383 (94-815)	12	51 (10-746)	12	1.61 (0.38-4.31)

n/a = not available

In three studies of patients with prior exposure to chemotherapy, the median CFU-GM yield in the leukapheresis product ranged from 20.9 to 32.7 x 10⁴/kg body weight (n = 105). In two of these studies where CD34⁺ yields in the leukapheresis product were also determined, the median CD34⁺ yields were 3.11 and 2.80 x 10⁶/kg, respectively (n = 56). In an additional study of 18 chemotherapy-naive patients, the median CFU-GM yield was 123.4 x 10⁴/kg.

Engraftment: Engraftment following NEUPOGEN-mobilized PBPC is summarized for 101 patients in the table below. In all studies a Cox regression model showed that the total number of CFU-GM and/or CD34⁺ cells collected was a significant predictor of time to platelet recovery.

In a randomized unblinded study of patients with HD or NHL undergoing myeloablative chemotherapy (Amgen study 3), 27 patients received NEUPOGEN-mobilized PBPC followed by NEUPOGEN and 31 patients received ABMT followed by NEUPOGEN. Patients randomized to the NEUPOGEN-mobilized PBPC group compared to the ABMT group had significantly fewer days of platelet transfusions (median 6 vs 10 days), a significantly shorter time to a sustained platelet count > 20,000/mm³ (median 16 vs 23 days), a significantly shorter time to recovery of a sustained ANC ≥ 500/mm³ (median 11 vs 14 days), significantly fewer days of red blood cell transfusions (median 2 vs 3 days) and a significantly shorter duration of posttransplant hospitalization.

		Amgen-sponsored Study 1 N = 13	Amgen-sponsored Study 2 N = 22	Amgen-sponsored Study 3 N = 27	Non-Amgen-sponsored Study N = 39
Median PBPC/kg Collected	MNC	9.5 x 10 ⁸	9.5 x 10 ⁸	8.1 x 10 ⁸	10.3 x 10 ⁸
	CD34 ⁺	n/a	3.1 x 10 ⁶	2.8 x 10 ⁶	6.2 x 10 ⁶
	CFU-GM	63.9 x 10 ⁴	25.3 x 10 ⁴	32.6 x 10 ⁴	n/a
Days to ANC ≥ 500/mm ³	Median	9	10	11	10
	Range	8-10	8-15	9-38	7-40
Days to Plt. ≥ 20,000/mm ³	Median	10	12.5	16	15.5
	Range	7-16	10-30	8-52	7-63

n/a = not available

Three of the 101 patients (3%) did not achieve the criteria for engraftment as defined by a platelet count ≥ 20,000/mm³ by day 28. In clinical trials of NEUPOGEN for the mobilization of PBPC, NEUPOGEN was administered to patients at 5 to 24 mcg/kg/day after reinfusion of the collected cells until a sustainable ANC (≥ 500/mm³) was reached. The rate of engraftment of these cells in the absence of NEUPOGEN posttransplantation has not been studied.

Patients With Severe Chronic Neutropenia

Severe chronic neutropenia (SCN) (idiopathic, cyclic, and congenital) is characterized by a selective decrease in the number of circulating neutrophils and an enhanced susceptibility to bacterial infections.

The daily administration of NEUPOGEN has been shown to be safe and effective in causing a sustained increase in the neutrophil count and a decrease in infectious morbidity

in children and adults with the clinical syndrome of SCN.¹⁵ In the phase 3 trial, summarized in the following table, daily treatment with NEUPOGEN resulted in significant beneficial changes in the incidence and duration of infection, fever, antibiotic use, and oropharyngeal ulcers. In this trial, 120 patients with a median age of 12 years (range 1 to 76 years) were treated.

**Overall Significant Changes in Clinical Endpoints Median Incidence^a
(Events) or Duration (Days) per 28-day Period**

	Control Patients ^b	NEUPOGEN-treated Patients	p-value
Incidence of Infection	0.50	0.20	< 0.001
Incidence of Fever	0.25	0.20	< 0.001
Duration of Fever	0.63	0.20	0.005
Incidence of Oropharyngeal Ulcers	0.26	0.00	< 0.001
Incidence of Antibiotic Use	0.49	0.20	< 0.001

^a Incidence values were calculated for each patient, and are defined as the total number of events experienced divided by the number of 28-day periods of exposure (on-study). Median incidence values were then reported for each patient group.

^b Control patients were observed for a 4-month period.

The incidence for each of these five clinical parameters was lower in the NEUPOGEN arm compared to the control arm for cohorts in each of the three major diagnostic categories. All three diagnostic groups showed favorable trends in favor of treatment. An analysis of variance showed no significant interaction between treatment and diagnosis, suggesting that efficacy did not differ substantially in the different diseases. Although NEUPOGEN substantially reduced neutropenia in all patient groups, in patients with cyclic neutropenia, cycling persisted but the period of neutropenia was shortened to 1 day.

As a result of the lower incidence and duration of infections, there was also a lower number of episodes of hospitalization [28 hospitalizations in 62 patients in the treated group vs 44 hospitalizations in 60 patients in the control group over a 4-month period (p = 0.0034)]. Patients treated with NEUPOGEN also reported a lower number of episodes of diarrhea, nausea, fatigue, and sore throat.

In the phase 3 trial, untreated patients had a median ANC of 210/mm³ (range 0 to

1550/mm³). NEUPOGEN therapy was adjusted to maintain the median ANC between 1500 and 10,000/mm³. Overall, the response to NEUPOGEN was observed in 1 to 2 weeks. The median ANC after 5 months of NEUPOGEN therapy for all patients was 7460/mm³ (range 30 to 30,880/mm³). NEUPOGEN dosing requirements were generally higher for patients with congenital neutropenia (2.3 to 40 mcg/kg/day) than for patients with idiopathic (0.6 to 11.5 mcg/kg/day) or cyclic (0.5 to 6 mcg/kg/day) neutropenia.

INDICATIONS AND USAGE

Cancer Patients Receiving Myelosuppressive Chemotherapy

NEUPOGEN is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with nonmyeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a significant incidence of severe neutropenia with fever (see CLINICAL EXPERIENCE). A complete blood count (CBC) and platelet count should be obtained prior to chemotherapy, and twice per week (see LABORATORY MONITORING) during NEUPOGEN therapy to avoid leukocytosis and to monitor the neutrophil count. In phase 3 clinical studies, NEUPOGEN therapy was discontinued when the ANC was $\geq 10,000/\text{mm}^3$ after the expected chemotherapy-induced nadir.

Patients With Acute Myeloid Leukemia Receiving Induction or Consolidation Chemotherapy

NEUPOGEN is indicated for reducing the time to neutrophil recovery and the duration of fever, following induction or consolidation chemotherapy treatment of adults with AML.

Cancer Patients Receiving Bone Marrow Transplant

NEUPOGEN is indicated to reduce the duration of neutropenia and neutropenia-related clinical sequelae, eg, febrile neutropenia, in patients with nonmyeloid malignancies undergoing myeloablative chemotherapy followed by marrow transplantation (see CLINICAL EXPERIENCE). It is recommended that CBCs and platelet counts be obtained at a minimum of three times per week (see LABORATORY MONITORING) following marrow infusion to monitor the recovery of marrow reconstitution.

Patients Undergoing Peripheral Blood Progenitor Cell Collection and Therapy

NEUPOGEN is indicated for the mobilization of hematopoietic progenitor cells into the peripheral blood for collection by leukapheresis. Mobilization allows for the collection of increased numbers of progenitor cells capable of engraftment compared with collection by leukapheresis without mobilization or bone marrow harvest. After myeloablative chemotherapy, the transplantation of an increased number of progenitor cells can lead to more rapid engraftment, which may result in a decreased need for supportive care (see

CLINICAL EXPERIENCE).

Patients With Severe Chronic Neutropenia

NEUPOGEN is indicated for chronic administration to reduce the incidence and duration of sequelae of neutropenia (eg, fever, infections, oropharyngeal ulcers) in symptomatic patients with congenital neutropenia, cyclic neutropenia, or idiopathic neutropenia (see CLINICAL EXPERIENCE). It is essential that serial CBCs with differential and platelet counts, and an evaluation of bone marrow morphology and karyotype be performed prior to initiation of NEUPOGEN therapy. The use of NEUPOGEN prior to confirmation of SCN may impair diagnostic efforts and may thus impair or delay evaluation and treatment of an underlying condition, other than SCN, causing the neutropenia.

CONTRAINDICATIONS

NEUPOGEN is contraindicated in patients with known hypersensitivity to *E coli*-derived proteins, Filgrastim, or any component of the product.

WARNINGS

Allergic-type reactions occurring on initial or subsequent treatment have been reported in < 1 in 4000 patients treated with NEUPOGEN. These have generally been characterized by systemic symptoms involving at least two body systems, most often skin (rash, urticaria, facial edema), respiratory (wheezing, dyspnea), and cardiovascular (hypotension, tachycardia). Some reactions occurred on initial exposure. Reactions tended to occur within the first 30 minutes after administration and appeared to occur more frequently in patients receiving NEUPOGEN IV. Rapid resolution of symptoms occurred in most cases after administration of antihistamines, steroids, bronchodilators, and/or epinephrine. Symptoms recurred in more than half the patients who were rechallenged.

Patients With Severe Chronic Neutropenia

The safety and efficacy of NEUPOGEN in the treatment of neutropenia due to other hematopoietic disorders (eg, myelodysplastic disorders or myeloid leukemia) have not been established. Care should be taken to confirm the diagnosis of SCN before initiating NEUPOGEN therapy.

While 9 of 325 patients developed myelodysplasia or myeloid leukemia while receiving NEUPOGEN during clinical trials, AML or abnormal cytogenetics have been reported to occur in the natural history of SCN without cytokine therapy.¹⁶ Abnormal cytogenetics have been associated with the eventual development of myeloid leukemia. The effect of NEUPOGEN on the development of abnormal cytogenetics and the effect of continued NEUPOGEN administration in patients with abnormal cytogenetics are unknown. If a

patient with SCN develops abnormal cytogenetics, the risks and benefits of continuing NEUPOGEN should be carefully considered (see ADVERSE REACTIONS).

PRECAUTIONS

General

Simultaneous Use With Chemotherapy and Radiation Therapy

The safety and efficacy of NEUPOGEN given simultaneously with cytotoxic chemotherapy have not been established. Because of the potential sensitivity of rapidly dividing myeloid cells to cytotoxic chemotherapy, do not use NEUPOGEN in the period 24 hours before through 24 hours after the administration of cytotoxic chemotherapy (see DOSAGE AND ADMINISTRATION).

The efficacy of NEUPOGEN has not been evaluated in patients receiving chemotherapy associated with delayed myelosuppression (eg, nitrosoureas) or with mitomycin C or with myelosuppressive doses of anti-metabolites such as 5-fluorouracil.

The safety and efficacy of NEUPOGEN have not been evaluated in patients receiving concurrent radiation therapy. Simultaneous use of NEUPOGEN with chemotherapy and radiation therapy should be avoided.

Potential Effect on Malignant Cells

NEUPOGEN is a growth factor that primarily stimulates neutrophils. However, the possibility that NEUPOGEN can act as a growth factor for any tumor type cannot be excluded. In a randomized study evaluating the effects of NEUPOGEN versus placebo in patients undergoing remission induction for AML, there was no significant difference in remission rate, disease-free or overall survival (see CLINICAL EXPERIENCE).

The safety of NEUPOGEN in chronic myeloid leukemia (CML) and myelodysplasia (MDS) has not been established.

When NEUPOGEN is used to mobilize PBPC, tumor cells may be released from the marrow and subsequently collected in the leukapheresis product. The effect of reinfusion of tumor cells has not been well-studied, and the limited data available are inconclusive.

Leukocytosis

Cancer Patients Receiving Myelosuppressive Chemotherapy

White blood cell counts of 100,000/mm³ or greater were observed in approximately 2% of patients receiving NEUPOGEN at doses above 5 mcg/kg/day. There were no reports of adverse events associated with this degree of leukocytosis. In order to avoid the potential complications of excessive leukocytosis, a CBC is recommended twice per week during NEUPOGEN therapy (see LABORATORY MONITORING).

Premature Discontinuation of NEUPOGEN Therapy

Cancer Patients Receiving Myelosuppressive Chemotherapy

A transient increase in neutrophil counts is typically seen 1 to 2 days after initiation of NEUPOGEN therapy. However, for a sustained therapeutic response, NEUPOGEN therapy should be continued following chemotherapy until the post nadir ANC reaches 10,000/mm³. Therefore, the premature discontinuation of NEUPOGEN therapy, prior to the time of recovery from the expected neutrophil nadir, is generally not recommended (see DOSAGE AND ADMINISTRATION).

Other

In studies of NEUPOGEN administration following chemotherapy, most reported side effects were consistent with those usually seen as a result of cytotoxic chemotherapy (see ADVERSE REACTIONS). Because of the potential of receiving higher doses of chemotherapy (ie, full doses on the prescribed schedule), the patient may be at greater risk of thrombocytopenia, anemia, and nonhematologic consequences of increased chemotherapy doses (please refer to the prescribing information of the specific chemotherapy agents used). Regular monitoring of the hematocrit and platelet count is recommended. Furthermore, care should be exercised in the administration of NEUPOGEN in conjunction with other drugs known to lower the platelet count. In septic patients receiving NEUPOGEN, the physician should be alert to the theoretical possibility of adult respiratory distress syndrome, due to the possible influx of neutrophils at the site of inflammation.

There have been rare reports (< 1 in 7000 patients) of cutaneous vasculitis in patients treated with NEUPOGEN. In most cases, the severity of cutaneous vasculitis was moderate or severe. Most of the reports involved patients with SCN receiving long-term NEUPOGEN therapy. Symptoms of vasculitis generally developed simultaneously with an increase in the ANC and abated when the ANC decreased. Many patients were able to continue NEUPOGEN at a reduced dose.

Information for Patients

In those situations in which the physician determines that the patient can safely and effectively self-administer NEUPOGEN, the patient should be instructed as to the proper dosage and administration. Patients should be referred to the “Information for Patients” labeling included with the package insert in each dispensing carton of NEUPOGEN. This patient information, however, is not intended to be a disclosure of all known or possible effects. If home use is prescribed, patients should be thoroughly instructed in the importance of proper disposal and cautioned against the reuse of needles, syringes, or drug product. A puncture-resistant container for the disposal of used syringes and needles should be available to the patient. The full container should be disposed of according to the directions provided by the physician.

Laboratory Monitoring

Cancer Patients Receiving Myelosuppressive Chemotherapy

A CBC and platelet count should be obtained prior to chemotherapy, and at regular intervals (twice per week) during NEUPOGEN therapy. Following cytotoxic chemotherapy, the neutrophil nadir occurred earlier during cycles when NEUPOGEN was administered, and WBC differentials demonstrated a left shift, including the appearance of promyelocytes and myeloblasts. In addition, the duration of severe neutropenia was reduced, and was followed by an accelerated recovery in the neutrophil counts. Therefore, regular monitoring of WBC counts, particularly at the time of the recovery from the postchemotherapy nadir, is recommended in order to avoid excessive leukocytosis.

Cancer Patients Receiving Bone Marrow Transplant

Frequent CBCs and platelet counts are recommended (at least three times per week) following marrow transplantation.

Patients With Severe Chronic Neutropenia

During the initial 4 weeks of NEUPOGEN therapy and during the 2 weeks following any dose adjustment, a CBC with differential and platelet count should be performed twice weekly. Once a patient is clinically stable, a CBC with differential and platelet count should be performed monthly.

In clinical trials, the following laboratory results were observed:

- Cyclic fluctuations in the neutrophil counts were frequently observed in patients with congenital or idiopathic neutropenia after initiation of NEUPOGEN therapy.
- Platelet counts were generally at the upper limits of normal prior to NEUPOGEN therapy. With NEUPOGEN therapy, platelet counts decreased but usually

remained within normal limits (see ADVERSE REACTIONS).

- Early myeloid forms were noted in peripheral blood in most patients, including the appearance of metamyelocytes and myelocytes. Promyelocytes and myeloblasts were noted in some patients.
- Relative increases were occasionally noted in the number of circulating eosinophils and basophils. No consistent increases were observed with NEUPOGEN therapy.
- As in other trials, increases were observed in serum uric acid, lactic dehydrogenase, and serum alkaline phosphatase.

Drug Interaction

Drug interactions between NEUPOGEN and other drugs have not been fully evaluated. Drugs which may potentiate the release of neutrophils, such as lithium, should be used with caution.

Carcinogenesis, Mutagenesis, Impairment of Fertility

The carcinogenic potential of NEUPOGEN has not been studied. NEUPOGEN failed to induce bacterial gene mutations in either the presence or absence of a drug metabolizing enzyme system. NEUPOGEN had no observed effect on the fertility of male or female rats, or on gestation at doses up to 500 mcg/kg.

Pregnancy Category C

NEUPOGEN has been shown to have adverse effects in pregnant rabbits when given in doses 2 to 10 times the human dose. There are no adequate and well-controlled studies in pregnant women. NEUPOGEN should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

In rabbits, increased abortion and embryoletality were observed in animals treated with NEUPOGEN at 80 mcg/kg/day. NEUPOGEN administered to pregnant rabbits at doses of 80 mcg/kg/day during the period of organogenesis was associated with increased fetal resorption, genitourinary bleeding, developmental abnormalities, decreased body weight, live births, and food consumption. External abnormalities were not observed in the fetuses of dams treated at 80 mcg/kg/day. Reproductive studies in pregnant rats have shown that NEUPOGEN was not associated with lethal, teratogenic, or behavioral effects on fetuses when administered by daily IV injection during the period of organogenesis at dose levels up to 575 mcg/kg/day.

In Segment III studies in rats, offspring of dams treated at > 20 mcg/kg/day exhibited a delay in external differentiation (detachment of auricles and descent of testes) and slight

growth retardation, possibly due to lower body weight of females during rearing and nursing. Offspring of dams treated at 100 mcg/kg/day exhibited decreased body weights at birth, and a slightly reduced 4-day survival rate.

Nursing Mothers

It is not known whether NEUPOGEN is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised if NEUPOGEN is administered to a nursing woman.

Pediatric Use

Serious long-term risks associated with daily administration of NEUPOGEN have not been identified in pediatric patients (ages 4 months to 17 years) with SCN. Limited data from patients who were followed in the phase 3 study for 1.5 years did not suggest alterations in growth and development, sexual maturation, or endocrine function.

The safety and efficacy in neonates and patients with autoimmune neutropenia of infancy have not been established.

In the cancer setting, 12 pediatric patients with neuroblastoma have received up to 6 cycles of cyclophosphamide, cisplatin, doxorubicin, and etoposide chemotherapy concurrently with NEUPOGEN; in this population, NEUPOGEN was well tolerated. There was one report of palpable splenomegaly associated with NEUPOGEN therapy, however, the only consistently reported adverse event was musculoskeletal pain, which is no different from the experience in the adult population.

ADVERSE REACTIONS

Cancer Patients Receiving Myelosuppressive Chemotherapy

In clinical trials involving over 350 patients receiving NEUPOGEN following nonmyeloablative cytotoxic chemotherapy, most adverse experiences were the sequelae of the underlying malignancy or cytotoxic chemotherapy. In all phase 2 and 3 trials, medullary bone pain, reported in 24% of patients, was the only consistently observed adverse reaction attributed to NEUPOGEN therapy. This bone pain was generally reported to be of mild-to-moderate severity, and could be controlled in most patients with non-narcotic analgesics; infrequently, bone pain was severe enough to require narcotic analgesics. Bone pain was reported more frequently in patients treated with higher doses (20 to 100 mcg/kg/day) administered IV, and less frequently in patients treated with lower SC doses of NEUPOGEN (3 to 10 mcg/kg/day).

In the randomized, double-blind, placebo-controlled trial of NEUPOGEN therapy following combination chemotherapy in patients (n = 207) with small cell lung cancer, the

following adverse events were reported during blinded cycles of study medication (placebo or NEUPOGEN at 4 to 8 mcg/kg/day). Events are reported as exposure adjusted since patients remained on double-blind NEUPOGEN a median of 3 cycles versus 1 cycle for placebo.

Event	% of Blinded Cycles With Events	
	NEUPOGEN N = 384 Patient Cycles	Placebo N = 257 Patient Cycles
Nausea/Vomiting	57	64
Skeletal Pain	22	11
Alopecia	18	27
Diarrhea	14	23
Neutropenic Fever	13	35
Mucositis	12	20
Fever	12	11
Fatigue	11	16
Anorexia	9	11
Dyspnea	9	11
Headache	7	9
Cough	6	8
Skin Rash	6	9
Chest Pain	5	6
Generalized Weakness	4	7
Sore Throat	4	9
Stomatitis	5	10
Constipation	5	10
Pain (Unspecified)	2	7

In this study, there were no serious, life-threatening, or fatal adverse reactions attributed to NEUPOGEN therapy. Specifically, there were no reports of flu-like symptoms, pleuritis, pericarditis, or other major systemic reactions to NEUPOGEN.

Spontaneously reversible elevations in uric acid, lactate dehydrogenase, and alkaline phosphatase occurred in 27% to 58% of 98 patients receiving blinded NEUPOGEN therapy following cytotoxic chemotherapy; increases were generally mild-to-moderate. Transient decreases in blood pressure (< 90/60 mmHg), which did not require clinical treatment, were reported in 7 of 176 patients in phase 3 clinical studies following

administration of NEUPOGEN. Cardiac events (myocardial infarctions, arrhythmias) have been reported in 11 of 375 cancer patients receiving NEUPOGEN in clinical studies; the relationship to NEUPOGEN therapy is unknown. No evidence of interaction of NEUPOGEN with other drugs was observed in the course of clinical trials (see PRECAUTIONS).

There has been no evidence for the development of antibodies or of a blunted or diminished response to NEUPOGEN in treated patients, including those receiving NEUPOGEN daily for almost 2 years.

Patients With Acute Myeloid Leukemia

In a randomized phase 3 clinical trial, 259 patients received NEUPOGEN and 262 patients received placebo postchemotherapy. Overall, the frequency of all reported adverse events was similar in both the NEUPOGEN and placebo groups (83% vs 82% in Induction 1, 61% vs 64% in Consolidation 1). Adverse events reported more frequently in the NEUPOGEN-treated group included: petechiae (17% vs 14%), epistaxis (9% vs 5%), and transfusion reactions (10% vs 5%). There were no significant differences in the frequency of these events.

There were a similar number of deaths in each treatment group during induction (25 NEUPOGEN vs 27 placebo). The primary causes of death included infection (9 vs 18), persistent leukemia (7 vs 5), and hemorrhage (6 vs 3). Of the hemorrhagic deaths, five cerebral hemorrhages were reported in the NEUPOGEN group and one in the placebo group. Other serious nonfatal hemorrhagic events were reported in the respiratory tract (4 vs 1), skin (4 vs 4), gastrointestinal tract (2 vs 2), urinary tract (1 vs 1), ocular (1 vs 0), and other nonspecific sites (2 vs 1). While 19 (7%) patients in the NEUPOGEN group and five (2%) patients in the placebo group experienced severe or fatal hemorrhagic events, overall, hemorrhagic adverse events were reported at a similar frequency in both groups (40% vs 38%). The time to transfusion-independent platelet recovery and the number of days of platelet transfusions were similar in both groups.

Cancer Patients Receiving Bone Marrow Transplant

In clinical trials, the reported adverse effects were those typically seen in patients receiving intensive chemotherapy followed by bone marrow transplant (BMT). The most common events reported in both control and treatment groups included stomatitis, nausea, and vomiting, generally of mild-to-moderate severity and were considered unrelated to NEUPOGEN. In the randomized studies of BMT involving 167 patients who received study drug, the following events occurred more frequently in patients treated with Filgrastim than in controls: nausea (10% vs 4%), vomiting (7% vs 3%), hypertension (4% vs 0%), rash (12% vs 10%), and peritonitis (2% vs 0%). None of these events were reported by the Investigator to be related to NEUPOGEN. One event of erythema

nodosum was reported moderate in severity and possibly related to NEUPOGEN.

Generally, adverse events observed in nonrandomized studies were similar to those seen in randomized studies, occurred in a minority of patients, and were of mild-to-moderate severity. In one study (n = 45), three serious adverse events reported by the Investigator were considered possibly related to NEUPOGEN. These included two events of renal insufficiency and one event of capillary leak syndrome. The relationship of these events to NEUPOGEN remains unclear since they occurred in patients with culture-proven infection with clinical sepsis who were receiving potentially nephrotoxic antibacterial and antifungal therapy.

Cancer Patients Undergoing Peripheral Blood Progenitor Cell Collection and Therapy

In clinical trials, 126 patients received NEUPOGEN for PBPC mobilization. In this setting, NEUPOGEN was generally well tolerated. Adverse events related to NEUPOGEN consisted primarily of mild-to-moderate musculoskeletal symptoms, reported in 44% of patients. These symptoms were predominantly events of medullary bone pain (33%). Headache was reported related to NEUPOGEN in 7% of patients. Transient increases in alkaline phosphatase related to NEUPOGEN were reported in 21% of the patients who had serum chemistries measured; most were mild-to-moderate.

All patients had increases in neutrophil counts during mobilization, consistent with the biological effects of NEUPOGEN. Two patients had a WBC count $> 100,000/\text{mm}^3$. No sequelae were associated with any grade of leukocytosis.

Sixty-five percent of patients had mild-to-moderate anemia and 97% of patients had decreases in platelet counts; five patients (out of 126) had decreased platelet counts to $< 50,000/\text{mm}^3$. Anemia and thrombocytopenia have been reported to be related to leukapheresis; however, the possibility that NEUPOGEN mobilization may contribute to anemia or thrombocytopenia has not been ruled out.

Patients With Severe Chronic Neutropenia

Mild-to-moderate bone pain was reported in approximately 33% of patients in clinical trials. This symptom was readily controlled with non-narcotic analgesics. Generalized musculoskeletal pain was also noted in higher frequency in patients treated with NEUPOGEN. Palpable splenomegaly was observed in approximately 30% of patients. Abdominal or flank pain was seen infrequently and thrombocytopenia ($< 50,000/\text{mm}^3$) was noted in 12% of patients with palpable spleens. Fewer than 3% of all patients underwent splenectomy, and most of these had a prestudy history of splenomegaly. Fewer than 6% of patients had thrombocytopenia ($< 50,000/\text{mm}^3$) during NEUPOGEN therapy, most of whom had a pre-existing history of thrombocytopenia. In most cases, thrombocytopenia was managed by NEUPOGEN dose reduction or interruption. An additional 5% of patients had platelet counts between 50,000 to $100,000/\text{mm}^3$. There were no associated

serious hemorrhagic sequelae in these patients. Epistaxis was noted in 15% of patients treated with NEUPOGEN, but was associated with thrombocytopenia in 2% of patients. Anemia was reported in approximately 10% of patients, but in most cases appeared to be related to frequent diagnostic phlebotomy, chronic illness, or concomitant medications. In clinical trials, myelodysplasia or myeloid leukemia was reported to have developed during NEUPOGEN therapy in approximately 3% of patients (9 of 325) (see WARNINGS). Twelve patients from a subset of 102 who had normal cytogenetic evaluations at baseline were subsequently found to have abnormalities, including monosomy 7, on routine repeat evaluation conducted after 18 to 52 months of NEUPOGEN therapy. It is unknown whether the development of these findings is related to chronic daily NEUPOGEN administration or reflects the natural history of SCN. Other adverse events infrequently observed and possibly related to NEUPOGEN therapy were: injection site reaction, rash, hepatomegaly, arthralgia, osteoporosis, cutaneous vasculitis, hematuria/proteinuria, alopecia, and exacerbation of some pre-existing skin disorders (eg, psoriasis).

OVERDOSAGE

In cancer patients receiving NEUPOGEN as an adjunct to myelosuppressive chemotherapy, it is recommended, to avoid the potential risks of excessive leukocytosis, that NEUPOGEN therapy be discontinued if the ANC surpasses $10,000/\text{mm}^3$ after the chemotherapy-induced ANC nadir has occurred. Doses of NEUPOGEN that increase the ANC beyond $10,000/\text{mm}^3$ may not result in any additional clinical benefit.

The maximum tolerated dose of NEUPOGEN has not been determined. Efficacy was demonstrated at doses of 4 to 8 mcg/kg/day in the phase 3 study of nonmyeloablative chemotherapy. Patients in the BMT studies received up to 138 mcg/kg/day without toxic effects, although there was a flattening of the dose response curve above daily doses of greater than 10 mcg/kg/day.

In NEUPOGEN clinical trials of cancer patients receiving myelosuppressive chemotherapy, WBC counts $> 100,000/\text{mm}^3$ have been reported in less than 5% of patients, but were not associated with any reported adverse clinical effects.

In cancer patients receiving myelosuppressive chemotherapy, discontinuation of NEUPOGEN therapy usually results in a 50% decrease in circulating neutrophils within 1 to 2 days, with a return to pretreatment levels in 1 to 7 days.

DOSAGE AND ADMINISTRATION

Cancer Patients Receiving Myelosuppressive Chemotherapy

The recommended starting dose of NEUPOGEN is 5 mcg/kg/day, administered as a single daily injection by SC bolus injection, by short IV infusion (15 to 30 minutes), or by continuous SC or continuous IV infusion. A CBC and platelet count should be obtained before instituting NEUPOGEN therapy, and monitored twice weekly during therapy. Doses may be increased in increments of 5 mcg/kg for each chemotherapy cycle, according to the duration and severity of the ANC nadir.

NEUPOGEN should be administered no earlier than 24 hours after the administration of cytotoxic chemotherapy. NEUPOGEN should not be administered in the period 24 hours before the administration of chemotherapy (see PRECAUTIONS). NEUPOGEN should be administered daily for up to 2 weeks, until the ANC has reached 10,000/mm³ following the expected chemotherapy-induced neutrophil nadir. The duration of NEUPOGEN therapy needed to attenuate chemotherapy-induced neutropenia may be dependent on the myelosuppressive potential of the chemotherapy regimen employed. NEUPOGEN therapy should be discontinued if the ANC surpasses 10,000/mm³ after the expected chemotherapy-induced neutrophil nadir (see PRECAUTIONS). In phase 3 trials, efficacy was observed at doses of 4 to 8 mcg/kg/day.

Cancer Patients Receiving Bone Marrow Transplant

The recommended dose of NEUPOGEN following BMT is 10 mcg/kg/day given as an IV infusion of 4 or 24 hours, or as a continuous 24-hour SC infusion. For patients receiving BMT, the first dose of NEUPOGEN should be administered at least 24 hours after cytotoxic chemotherapy and at least 24 hours after bone marrow infusion.

During the period of neutrophil recovery, the daily dose of NEUPOGEN should be titrated against the neutrophil response as follows:

Absolute Neutrophil Count	NEUPOGEN Dose Adjustment
When ANC > 1000/mm ³ for 3 consecutive days	Reduce to 5 mcg/kg/day ^a
then: If ANC remains > 1000/mm ³ for 3 more consecutive days	Discontinue NEUPOGEN
then: If ANC decreases to < 1000/mm ³	Resume at 5 mcg/kg/day

^a If ANC decreases to < 1000/mm³ at any time during the 5 mcg/kg/day administration, NEUPOGEN should be increased to 10 mcg/kg/day, and the above steps should then be followed.

Peripheral Blood Progenitor Cell Collection and Therapy in Cancer Patients

The recommended dose of NEUPOGEN for the mobilization of PBPC is 10 mcg/kg/day SC, either as a bolus or a continuous infusion. It is recommended that NEUPOGEN be given for at least 4 days before the first leukapheresis procedure and continued until the last leukapheresis. Although the optimal duration of NEUPOGEN administration and leukapheresis schedule have not been established, administration of NEUPOGEN for 6 to 7 days with leukaphereses on days 5, 6, and 7 was found to be safe and effective (see CLINICAL EXPERIENCE for schedules used in clinical trials). Neutrophil counts should be monitored after 4 days of NEUPOGEN, and NEUPOGEN dose modification should be considered for those patients who develop a WBC count > 100,000/mm³.

In all clinical trials of NEUPOGEN for the mobilization of PBPC, NEUPOGEN was also administered after reinfusion of the collected cells (see CLINICAL EXPERIENCE).

Patients With Severe Chronic Neutropenia

NEUPOGEN should be administered to those patients in whom a diagnosis of congenital, cyclic, or idiopathic neutropenia has been definitively confirmed. Other diseases associated with neutropenia should be ruled out.

Starting Dose:

Congenital Neutropenia: The recommended daily starting dose is 6 mcg/kg BID SC every day.

Idiopathic or Cyclic Neutropenia: The recommended daily starting dose is 5 mcg/kg as a single injection SC every day.

Dose Adjustments:

Chronic daily administration is required to maintain clinical benefit. Absolute neutrophil count should not be used as the sole indication of efficacy. The dose should be

individually adjusted based on the patients' clinical course as well as ANC. In the phase 3 study, the target ANC was 1500/mm³ to 10,000/mm³. However, patients may experience clinical benefit with ANCs below this target range. The dose should be reduced if the ANC is persistently greater than 10,000/mm³.

Dilution

If required, NEUPOGEN may be diluted in 5% dextrose. NEUPOGEN diluted to concentrations between 5 and 15 mcg/mL should be protected from adsorption to plastic materials by the addition of Albumin (Human) to a final concentration of 2 mg/mL. When diluted in 5% dextrose or 5% dextrose plus Albumin (Human), NEUPOGEN is compatible with glass bottles, PVC and polyolefin IV bags, and polypropylene syringes.

Dilution of NEUPOGEN to a final concentration of less than 5 mcg/mL is not recommended at any time. **Do not dilute with saline at any time; product may precipitate.**

Storage

NEUPOGEN should be stored in the refrigerator at 2° to 8°C (36° to 46°F). Avoid shaking. Prior to injection, NEUPOGEN may be allowed to reach room temperature for a maximum of 24 hours. Any vial left at room temperature for greater than 24 hours should be discarded. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit; if particulates or discoloration are observed, the container should not be used.

HOW SUPPLIED

NEUPOGEN: Use only one dose per vial; do not re-enter the vial. Discard unused portions. Do not save unused drug for later administration.

Single-dose, preservative-free vials containing 300 mcg (1 mL) of Filgrastim (300 mcg/mL). Dispensing packs of 10 (NDC 55513-530-10).

Single-dose, preservative-free vials containing 480 mcg (1.6 mL) of Filgrastim (300 mcg/mL). Dispensing packs of 10 (NDC 55513-546-10).

NEUPOGEN should be stored at 2° to 8° C (36° to 46° F). Avoid shaking.

REFERENCES

1. Zsebo KM, Cohen AM, Murdock DC, et al. Recombinant human granulocyte colony-stimulating factor: Molecular and biological characterization. *Immunobiol* 1986;172:175-184.
2. Welte K, Bonilla MA, Gillio AP, et al. Recombinant human G-CSF: Effects on hematopoiesis in normal and cyclophosphamide treated primates. *J Exp Med* 1987;165:941-948.
3. Duhresen U, Villeval JL, Boyd J, et al. Effects of recombinant human granulocyte colony-stimulating factor on hematopoietic progenitor cells in cancer patients. *Blood* 1988;72:2074-2081.
4. Souza LM, Boone TC, Gabrilove J, et al. Recombinant human granulocyte colony-stimulating factor: Effects on normal and leukemic myeloid cells. *Science* 1986;232:61-65.
5. Weisbart RH, Kacena A, Schuh A, and Golde DW. GM-CSF induces human neutrophil IgA-mediated phagocytosis by an IgA Fc receptor activation mechanism. *Nature* 1988;332:647-648.
6. Kitagawa S, Yuo A, Souza LM, Saito M, Miura Y, and Takaku F. Recombinant human granulocyte colony-stimulating factor enhances superoxide release in human granulocytes stimulated by chemotactic peptide. *Biochem Biophys Res Commun* 1987;1443:1146.
7. Glaspy JA, Baldwin GC, Robertson PA, et al. Therapy for neutropenia in hairy cell leukemia with recombinant human granulocyte colony-stimulating factor. *Ann Int Med* 1988;109:789-795.
8. Yuo A, Kitagawa S, Ohsaka A, et al. Recombinant human granulocyte colony-stimulating factor as an activator of human granulocytes: Potentiation of responses triggered by receptor-mediated agonists and stimulation of C3bi receptor expression and adherence. *Blood* 1989;74:2144-2149.
9. Gabrilove JL, Jakubowski A, Fain K, et al. Phase I study of granulocyte colony-stimulating factor in patients with transitional cell carcinoma of the urothelium. *J Clin Invest* 1988;82:1454-1461.
10. Morstyn G, Souza L, Keech J, et al. Effect of granulocyte colony-stimulating factor on neutropenia induced by cytotoxic chemotherapy. *Lancet* 1988;1:667-672.
11. Bronchud MH, Scarffe JH, Thatcher N, et al. Phase I/II study of recombinant human granulocyte colony-stimulating factor in patients receiving intensive chemotherapy for small cell lung cancer. *Br J Cancer* 1987;56:809-813.
12. Gabrilove JL, Jakubowski A, Scher H, et al. Effect of granulocyte colony-stimulating factor on neutropenia and associated morbidity due to chemotherapy for transitional cell carcinoma of the urothelium. *N Engl J Med* 1988;318:1414-1422.

13. Neidhart J, Mangalik A, Kohler W, et al. Granulocyte colony-stimulating factor stimulates recovery of granulocytes in patients receiving dose-intensive chemotherapy without bone-marrow transplantation. *J Clin Oncol* 1989;7:1685-1691.
14. Bronchud MH, Howell A, Crowther D, et al. The use of granulocyte colony-stimulating factor to increase the intensity of treatment with doxorubicin in patients with advanced breast and ovarian cancer. *Br J Cancer* 1989;60:121-128.
15. Dale DC, Bonilla MA, Davis MW, et al. A randomized controlled phase III trial of recombinant human granulocyte colony-stimulating factor (Filgrastim) for treatment of severe chronic neutropenia. *Blood* 1993;81:2496-2502.
16. Schroeder TM and Kurth R. Spontaneous chromosomal breakage and high incidence of leukemia in inherited disease. *Blood* 1971;37:96-112.
17. Heil G, Hoelzer D, Sanz MA, et al. A randomized, double-blind, placebo-controlled, phase III study of Filgrastim in remission induction and consolidation therapy for adults with de novo Acute Myeloid Leukemia. *Blood* 1997;90:4710-4718.

This product and its use are covered by the following US Patent Nos.: 4,810,643; 4,999,291; 5,528,823; 5,580,755.

AMGEN LOGO

Manufactured by:

Amgen Inc.

One Amgen Center Drive

Thousand Oaks, California 91320-1789

© 1991-1998 Amgen Inc. All rights reserved.

Issue Date: 04/02/98