

Carton

Front Panel

new

CURES ATHLETE'S FOOT

**FULL PRESCRIPTION STRENGTH
NOW AVAILABLE WITHOUT A PRESCRIPTION
RELIEVES ITCHING & BURNING**

LAMISIL®^{AT}™ Cream
Terbinafine hydrochloride cream 1%
Antifungal

LAMISIL®^{AT}™ Cream
Terbinafine hydrochloride cream 1%
Antifungal

Net Wt. 12 g (.42 oz)

LAMISIL®^{AT}™ Cream
Terbinafine hydrochloride cream 1%

LAMISIL®^{AT}™ Cream
Terbinafine hydrochloride cream 1%

| Active ingredient | Purpose |
|------------------------------------|----------------|
| Terbinafine hydrochloride 1% | Antifungal |

Uses

- cures athlete's foot (tinea pedis)
- cures jock itch (tinea cruris) and ringworm (tinea corporis)
- relieves itching, burning, cracking and scaling which accompany these conditions

Warnings

For external use only

Do not use

- on nails or scalp
- in or near the mouth or the eyes
- for vaginal yeast infections

When using this product do not get into the eyes. If eye contact occurs, rinse thoroughly with water.

Stop use and ask a doctor if too much irritation occurs or gets worse.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control center right away.

Directions

- adults and children 12 years and older

- use the tip of the cap to break the seal and open the tube
- wash the affected skin with soap and water and dry completely before applying
- **for athlete's foot** wear well-fitting, ventilated shoes. Change shoes and socks at least once daily.
 - **between the toes only:** apply twice a day (morning and night) for **1 week** or as directed by a doctor. (Foot Diagram) **1 week**
- **between the toes**
 - **on the bottom or sides of the foot:** apply twice a day (morning and night) for **2 weeks** or as directed by a doctor. (Foot Diagram) **2 weeks on the bottom or sides of the foot**
- **for jock itch and ringworm** apply once a day (morning or night) for **1 week** or as directed by a doctor.
- wash hands after each use
- children under 12 years: ask a doctor

Other information

- do not use if [insert identifying characteristic of] seal on tube is broken or is not visible
- store at 20^o - 25^o C (68^o - 77^o F)

Inactive ingredients benzyl alcohol, cetyl alcohol, cetyl palmitate, isopropyl myristate, polysorbate 60, purified water, sodium hydroxide, sorbitan monostearate, stearyl alcohol

Questions? call 1-800-570-9611 weekdays 9 am - 5 pm eastern time

Novartis Consumer Health, Inc.

Summit, NJ 07901-1312

©1999

Note: Lot and Expiration date will be embossed on the tuck flap.

LAMISIL® AT™ Cream (on both side flaps)