
Drug Facts

| Active ingredient (in each brown oval capsule) | Purpose |
|---|----------------|
| Solubilized ibuprofen equal to 200 mg ibuprofen..... (present as the free acid and potassium salt) | Pain reliever |

Use treats migraine

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:
• hives • facial swelling • asthma (wheezing) • shock

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if you have

- never had migraines diagnosed by a health professional
 - a headache that is different from your usual migraines
 - the worst headache of your life
 - fever and stiff neck
 - headaches beginning after or caused by head injury, exertion, coughing or bending
 - experienced your first headache after the age of 50
 - daily headaches
 - a migraine so severe as to require bed rest
 - problems or serious side effects from taking pain relievers or fever reducers
 - stomach pain
-

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any continuing medical condition
 - taking other drugs on a regular basis
 - taking another product containing ibuprofen, or any other pain reliever/fever reducer
-

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
 - migraine headache pain is not relieved or gets worse after first dose
 - stomach pain occurs with the use of this product
 - new or unexpected symptoms occur
-

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

| | |
|------------------------|--|
| adults: | <ul style="list-style-type: none">• take 2 capsules with a glass of water• if symptoms persist or worsen, ask your doctor• do not take more than 2 capsules in 24 hours, unless directed by a doctor |
| under 18 years of age: | <ul style="list-style-type: none">• ask a doctor |

Other information

- read all directions and warnings before use. Keep carton.
- store at 20 - 25°C (68 - 77°F)
- avoid excessive heat 40°C (above 104°F)

Inactive ingredients black iron oxide, D&C yellow no. 10, FD&C green no. 3, FD&C red no. 40, gelatin, lecithin, light mineral oil, pharmaceutical glaze, polyethylene glycol, potassium hydroxide, purified water, simethicone, sorbitan, sorbitol

Questions? Call 1-800-88-ADVIL

Do not use if seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing

Distributed by WHITEHALL-ROBINS HEALTHCARE
Madison, NJ 07940 Made in USA
1999 WHITEHALL-ROBINS HEALTHCARE