

Drug Facts (continued)

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- fever gets worse or lasts more than 3 days
- pain gets worse or lasts more than 10 days
- stomach pain occurs with the use of this product
- the painful area is red or swollen
- any new or unexpected symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- adults, take 1 gel caplet every 4 to 6 hours while symptoms occur
- if pain or fever does not respond to 1 gel caplet, 2 gel caplets may be used, but do not exceed 6 gel caplets in 24 hours, unless directed by a doctor
- the smallest effective dose should be used
- children, do not give to children under 12, unless directed by a doctor

Other Information

- read all warnings and directions before use. Keep carton
- store at 20-25°C (68-77°F)
- avoid excessive heat 40°C (above 104°F)

ADVANCED MEDICINE FOR PAIN™

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever / Fever Reducer

24 Coated Gel Caplets*
*Capsule-Shaped Gelatin Coated Tablets

Gel Caplets

ADVANCED MEDICINE FOR PAIN™

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever / Fever Reducer

24 Coated Gel Caplets*
*Capsule-Shaped Gelatin Coated Tablets

Gel Caplets

Drug Facts (continued)

Inactive ingredients: croscarmellose sodium, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, hydroxypropyl methylcellulose, iron oxides, medium chain triglycerides, pharmaceutical ink, propylglycol, silicon dioxide, sodium lauryl sulfate, starch, stearic acid, titanium dioxide, triacetin

Do Not Use if seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing.

Whitehall-Robins Healthcare, Madison, NJ 07940
Made in USA U.S. Patent No. 5,087,454

Appearance of the beige and brown Advil gel caplet is a trademark of Whitehall-Robins Healthcare

Drug Facts

Active ingredient (in each gel caplet)	Purposes
Ibuprofen 200 mg	Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
 - temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction which may include

- hives
- facial swelling
- asthma (wheezing)
- shock

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if you have had problems or side effects with any pain reliever/fever reducer

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any ongoing medical condition
- taking other drugs on a regular basis
- taking any other product containing ibuprofen, or any other pain reliever/fever reducer

NDA 18-989S-053
Amendment



Drug Facts (continued)

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- fever gets worse or lasts more than 3 days
- pain gets worse or lasts more than 10 days
- stomach pain occurs with the use of this product
- the painful area is red or swollen
- any new or unexpected symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- adults: take 1 gel caplet every 4 to 6 hours while symptoms occur
- if pain or fever does not respond to 1 gel caplet, 2 gel caplets may be used, but do not exceed 6 gel caplets in 24 hours, unless directed by a doctor
- the smallest effective dose should be used
- children: do not give to children under 12 unless directed by a doctor

Other information

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat 40°C (above 104°F)

ADVANCED MEDICINE FOR PAIN™

Advil

Ibuprofen Tablets, 200 mg
Pain Reliever / Fever Reducer

50 Coated Gel Caplets*  Gel Caplets

*Capsule-Shaped Gelatin Coated Tablets

ADVANCED MEDICINE FOR PAIN™

Advil

Ibuprofen Tablets, 200 mg
Pain Reliever / Fever Reducer

50 Coated Gel Caplets*  Gel Caplets

*Capsule-Shaped Gelatin Coated Tablets

Drug Facts (continued)

Inactive ingredients: croscarmellose sodium, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, hydroxypropyl methylcellulose, iron oxides, medium chain triglycerides, pharmaceutical ink, propyl gallate, silicon dioxide, sodium lauryl sulfate, starch, stearic acid, titanium dioxide, triacetin

Do Not Use if seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing.

Whitehall-Robins Healthcare, Madison, NJ 07940
Made in USA - U.S. Patent No. 5,087,454
Appearance of the beige and brown Advil gel caplet is a trademark of Whitehall-Robins Healthcare

Drug Facts

Active ingredient (in each gel caplet)
Ibuprofen 200 mg

Purposes
Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - the common cold
 - toothache
 - muscular aches
- temporarily reduces fever
- backache
- minor pain of arthritis
- menstrual cramps

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Drug Facts (continued)

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding.


Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer.

Ask a doctor before use if you have had problems or side effects with any pain reliever/fever reducer

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any continuing medical condition
- taking other drugs on a regular basis
- taking any other product containing ibuprofen, or any other pain reliever/fever reducer

NDA 18-989/S-053
Amendment



Drug Facts (continued)
 If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- adults: take 1 gel caplet every 4 to 6 hours while symptoms occur
- if pain or fever does not respond to 1 gel caplet, 2 gel caplets may be used, but do not exceed 6 gel caplets in 24 hours, unless directed by a doctor
- the smallest effective dose should be used
- children: do not give to children under 12 unless directed by a doctor

Drug Facts (continued)
Other information

- read all warnings and directions before use. Keep carton.
- store at 20-25 C (68-77 F)
- avoid excessive heat 40 C (above 104 F)

Inactive ingredients
 croscarmellose sodium, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, hydroxypropyl methylcellulose, iron oxides, medium chain triglycerides, pharmaceutical ink, propyl gallate, silicon dioxide, sodium lauryl sulfate, starch, stearic acid, titanium dioxide, triacetin

ADVANCED MEDICINE FOR PAIN™

Advil®

Ibuprofen Tablets, 200 mg
 Pain Reliever/Fever Reducer

100 Coated Gel Caplets*
*Capsule-Shaped Gelatin Coated Tablets



ADVANCED MEDICINE FOR PAIN™

Advil®

Ibuprofen Tablets, 200 mg
 Pain Reliever/Fever Reducer

100 Coated Gel Caplets*
*Capsule-Shaped Gelatin Coated Tablets



Advil®
 100 Coated Gel Caplets
 Gel Caplets

Do Not Use if seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing.

Whitehall-Robins Healthcare, Madison, NJ 07940
 Made in USA U.S. Patent No. 5,087,454
 Appearance of the beige and brown Advil gel caplet is a trademark of Whitehall-Robins Healthcare

Drug Facts
Active ingredient (in each gel caplet) Purposes
 ibuprofen 200 mg Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - backache
 - the common cold
 - minor pain of arthritis
 - toothache
 - menstrual cramps
 - muscular aches
 - temporarily reduces fever

Warnings
 Allergy alert: ibuprofen may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock


 Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding.

Drug Facts (continued)
 Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if you have had problems or side effects with any pain reliever/fever reducer.
 Ask a doctor or pharmacist before use if you are under a doctor's care for any continuing medical condition.
 taking other drugs on a regular basis
 taking any other product containing ibuprofen, or any other pain reliever/fever reducer
 Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- fever gets worse or lasts more than 3 days
- pain gets worse or lasts more than 10 days
- stomach pain occurs with the use of this product
- the painful area is red or swollen
- any new or unexpected symptoms occur

NDA 18-989/S-053
 Amendment



128 CODE AREA
#222174

Drug Facts (continued)
 If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- adults: take 1 gel caplet every 4 to 6 hours while symptoms occur
- if pain or fever does not respond to 1 gel caplet, 2 gel caplets may be used, but do not exceed 6 gel caplets in 24 hours, unless directed by a doctor
- the smallest effective dose should be used
- children: do not give to children under 12 unless directed by a doctor

Drug Facts (continued)

Other information

- read all warnings and directions before use. Keep carton.
- store at 20-25 C (68-77 F)
- avoid excessive heat 40 C (above 104 F)

Inactive ingredients
 croscarmellose sodium, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, hydroxypropyl methylcellulose, iron oxides, medium chain triglycerides, pharmaceutical ink, propyl gallate, silicon dioxide, sodium lauryl sulfate, starch, stearic acid, titanium dioxide, triacetin

128 CODE AREA
#222174

UNVARNISHED AREA
(PRINTED SIDE OF CARTON)

ADVANCED MEDICINE FOR PAIN™

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer

100 Coated Gel Caplets*
*Capsule-Shaped Gelatin Coated Tablets



TYP (8)

128 CODE AREA
#222174

128 CODE AREA
#222174

ADVANCED MEDICINE FOR PAIN™

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer

100 Coated Gel Caplets*
*Capsule-Shaped Gelatin Coated Tablets



Advil®
100 Coated Gel Caplets
Gel Caplets

Do Not Use if seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing.

Whitehall-Robins Healthcare, Madison, NJ 07940
 Made in USA U.S. Patent No. 5,087,454
 Appearance of the beige and brown Advil gel caplet is a trademark of Whitehall-Robins Healthcare

3 0573-0165-40
2

NDA 18-989/S-053
Amendment

Drug Facts

Active ingredient (in each gel caplet) **Purposes**
 Ibuprofen 200 mg Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - backache
 - the common cold
 - minor pain of arthritis
 - toothache
 - menstrual cramps
 - muscular aches
 - temporarily reduces fever

Warnings
 Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

 Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if you have had problems or side effects with any pain reliever/fever reducer

Ask a doctor or pharmacist before use if you are under a doctor's care for any continuing medical condition

- taking other drugs on a regular basis
- taking any other product containing ibuprofen, or any other pain reliever/fever reducer

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- fever gets worse or lasts more than 3 days
- pain gets worse or lasts more than 10 days
- stomach pain occurs with the use of this product
- the painful area is red or swollen
- any new or unexpected symptoms occur

12

1/8 X
TYP

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the **last 3 months** of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- adults: take 1 gel caplet every 4 to 6 hours while symptoms occur
- if pain or fever does not respond to 1 gel caplet, 2 gel caplets may be used, but do not exceed 6 gel caplets in 24 hours, unless directed by a doctor
- the smallest effective dose should be used
- children: do not give to children under 12 unless directed by a doctor

Other information

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat 40°C (above 104°F)

Inactive ingredients

croscarmellose sodium, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, hydroxypropyl methylcellulose, iron oxides, medium chain triglycerides, pharmaceutical ink, propyl gallate, silicon dioxide, sodium lauryl sulfate, starch, stearic acid, titanium dioxide, triacetin

ADVANCED MEDICINE FOR PAIN™

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer

250 Coated Gel Caplets*

*Capsule-Shaped Gelatin Coated Tablets



Gel Caplets

ADVANCED MEDICINE FOR PAIN™

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer

250 Coated Gel Caplets*

*Capsule-Shaped Gelatin Coated Tablets



Gel Caplets

Advil®

250 Coated Gel Caplets
Gel Caplets

Do Not Use if seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing.

Whitehall-Robins Healthcare, Madison, NJ 07940
Made in USA U.S. Patent No. 5,087,454
Appearance of the beige and brown Advil gel caplet is a trademark of Whitehall-Robins Healthcare

Drug Facts

Active ingredient (in each gel caplet).....
Ibuprofen 200 mg.....

Purposes
Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if you have had problems or side effects with any pain reliever/fever reducer

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any continuing medical condition
- taking other drugs on a regular basis
- taking any other product containing ibuprofen, or any other pain reliever/fever reducer


Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- stomach pain occurs with the use of this product
- fever gets worse or lasts more than 3 days
- the painful area is red or swollen
- pain gets worse or lasts more than 10 days
- any new or unexpected symptoms occur


NDA 18-989S-053
Amendment

1/2

128 CODE AREA
#220084



1/2



128 CODE AREA

128 CODE AREA
#220084



1/2



2516

128 CODE AREA
#220084

3 0573-0165-46 4



1/2

11516
1/8