

**Drug Facts (continued)**

If pregnant or breast feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

do not take more than directed

adults and children 12 years and older

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used, but do not exceed 6 caplets in 24 hours, unless directed by a doctor
- the smallest effective dose should be used

children under 12 years

- ask a doctor

**Other information**

- do not use if container is open or if tear strip is broken or missing
- store between 20 - 25°C (68 - 77°F).
- see container label for lot and expiration date

**Inactive ingredients**

camuba wax, corn starch, FD&C Yellow #6, hydroxypropyl methylcellulose, iron oxide, polydextrose, polyethylene glycol, silicon dioxide, stearic acid, titanium dioxide

**Questions or comments?**


Call toll free 1-800-962-5357



McNeil Consumer Healthcare, DIVISION OF MCNEIL-PPC, INC., FORT WASHINGTON, PA 19034 USA © MCN-PPC, Inc. '01

XXXXXXXX

www.motrin.com



**Drug Facts**

**Active Ingredient (in each caplet)**

Ibuprofen 200 mg.....fever reducer

**Purposes**

Pain reliever/fever reducer

**Uses** temporarily relieves minor aches and pains due to:

- headache
- minor pain of arthritis
- backache
- menstrual cramps
- reduces fever
- muscular aches
- toothache
- the common cold

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include: hives, asthma (wheezing), facial swelling, shock

**Alcohol warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if you have**

- stomach pain
- problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

under a doctor's care for any serious condition

- taking any other drug
- taking any other product that contains ibuprofen, or any other pain reliever/fever reducer

**When using this product give with food or milk if stomach upset occurs**

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear

**Labeling Format Information:**

Fonts: Helvetica regular, bold and bold italic.

<b>Drug Facts:</b>	8.5 pt	<b>Leading:</b>	6.5 pt
<b>Header:</b>	8 pt	<b>Bullets:</b>	5 pt
<b>Subheader:</b>	6 pt	<b>Barlines:</b>	2.5 pt
<b>Body Text:</b>	6 pt	<b>Hairlines:</b>	0.5 pt
<b>Drug Facts (continued):</b>	8 pt		

Horizontal Scale: 94-100%

Average Kerning: 0

**Drug Facts (continued)**

**If pregnant or breast feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

■ do not take more than directed

adults and children 12 years and older	■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used, but do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ the smallest effective dose should be used
children under 12 years	■ ask a doctor

**Other information**

do not use if container is open or if tear strip is broken or missing  
store between 20 - 25 C (68 - 77 F).  
see container label for lot and expiration date

**Inactive ingredients**

camuba wax, corn starch, FD&C Yellow #6, hydroxypropyl methylcellulose, iron oxide, polydextrose, polyethylene glycol, silicon dioxide, stearic acid, titanium dioxide

**Questions or comments?**

Call toll free 1-800-962-5357




McNeil Consumer Healthcare,  
DIVISION OF MCNEIL-PPC, INC.,  
FORT WASHINGTON, PA 19034 USA  
© MCN-PPC, Inc. '01

XXXXXXXX

[www.motrin.com](http://www.motrin.com)

EXP.



**Motrin**  
IB

Pain Reliever/  
Fever Reducer

8

**Drug Facts**

Active Ingredient (in each caplet)	Purposes
Ibuprofen 200 mg.....	Pain reliever/ fever reducer

**Uses** temporarily relieves minor aches and pains due to:

- headache
- minor pain of arthritis
- backache
- menstrual cramps
- reduces fever
- muscular aches
- toothache
- the common cold

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include:

- hives
- asthma (wheezing)
- facial swelling
- shock

**Alcohol warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding.

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use** if you have

- stomach pain
- problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use** if you are

- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains ibuprofen, or any other pain reliever/fever reducer

**When using this product** give with food or milk; if stomach upset occurs

**Stop use and ask a doctor** if

- an allergic reaction occurs. Seek medical help right away.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear



-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/

-----  
Charles Ganley  
9/7/01 06:05:31 PM