

NDA 20-362/S-053

Page 3

CEREZYME® (imiglucerase for injection)

200 Units

400 Units

## DESCRIPTION

**Cerezyme®** (imiglucerase for injection) is an analogue of the human enzyme,  $\beta$ -glucocerebrosidase produced by recombinant DNA technology.  $\beta$ -Glucocerebrosidase ( $\beta$ -D-glucosyl-N-acylsphingosine glucosylhydrolase, E.C. 3.2.1.45) is a lysosomal glycoprotein enzyme which catalyzes the hydrolysis of the glycolipid glucocerebroside to glucose and ceramide.

**Cerezyme®** is produced by recombinant DNA technology using mammalian cell culture (Chinese hamster ovary). Purified imiglucerase is a monomeric glycoprotein of 497 amino acids, containing 4 N-linked glycosylation sites ( $M_r = 60,430$ ). Imiglucerase differs from placental glucocerebrosidase by one amino acid at position 495 where histidine is substituted for arginine. The oligosaccharide chains at the glycosylation sites have been modified to terminate in mannose sugars. The modified carbohydrate structures on imiglucerase are somewhat different from those on placental glucocerebrosidase. These mannose-terminated oligosaccharide chains of imiglucerase are specifically recognized by endocytic carbohydrate receptors on macrophages, the cells that accumulate lipid in Gaucher disease.

**Cerezyme®** is supplied as a sterile, non-pyrogenic, white to off-white lyophilized product. The quantitative composition of the lyophilized drug is provided in the following table:

<b>Ingredient</b>	<b>200 Unit Vial</b>	<b>400 Unit Vial</b>
Imiglucerase (total amount)*	212 units	424 units
Mannitol	170 mg	340 mg
Sodium Citrates (Trisodium Citrate)	70 mg (52 mg)	140 mg (104 mg)
(Disodium Hydrogen Citrate)	(18 mg)	(36 mg)
Polysorbate 80, NF	0.53 mg	1.06 mg
Citric Acid and/or Sodium Hydroxide may have been added at the time of manufacture to adjust pH.		

\*This provides a respective withdrawal dose of 200 and 400 units of imiglucerase.

An enzyme unit (U) is defined as the amount of enzyme that catalyzes the hydrolysis of one micromole of the synthetic substrate para-nitrophenyl- $\beta$ -D-glucopyranoside (pNP-Glc) per minute at 37°C. The product is stored at 2-8°C (36-46°F). After reconstitution with Sterile Water for Injection, USP, the imiglucerase concentration is 40 U/mL (see **DOSAGE AND ADMINISTRATION** for final concentrations and volumes). Reconstituted solutions have a pH of approximately 6.1.

In addition, Haemaccel® (cross-linked gelatin polypeptides), which is used as a stabilizing agent during the manufacturing process, may also be present in very small amounts in the final product.

NDA 20-362/S-053

Page 4

## CLINICAL PHARMACOLOGY

### Mechanism of Action/Pharmacodynamics

Gaucher disease is characterized by a deficiency of  $\beta$ -glucocerebrosidase activity, resulting in accumulation of glucocerebroside in tissue macrophages which become engorged and are typically found in the liver, spleen, and bone marrow and occasionally in lung, kidney, and intestine. Secondary hematologic sequelae include severe anemia and thrombocytopenia in addition to the characteristic progressive hepatosplenomegaly, skeletal complications, including osteonecrosis and osteopenia with secondary pathological fractures. **Cerezyme**<sup>®</sup> (imiglucerase for injection) catalyzes the hydrolysis of glucocerebroside to glucose and ceramide. In clinical trials, **Cerezyme**<sup>®</sup> improved anemia and thrombocytopenia, reduced spleen and liver size, and decreased cachexia to a degree similar to that observed with Ceredase<sup>®</sup> (*alglucerase injection*).

### Pharmacokinetics

During one-hour intravenous infusions of four doses (7.5, 15, 30, 60 U/kg) of **Cerezyme**<sup>®</sup> (imiglucerase for injection) steady-state enzymatic activity was achieved by 30 minutes. Following infusion, plasma enzymatic activity declined rapidly with a half-life ranging from 3.6 to 10.4 minutes. Plasma clearance ranged from 9.8 to 20.3 mL/min/kg (mean  $\pm$  S.D.,  $14.5 \pm 4.0$  mL/min/kg). These variables do not appear to be influenced by dose or duration of infusion. However, only one or two patients were studied at each dose level and infusion rate. The pharmacokinetics of **Cerezyme**<sup>®</sup> do not appear to be different from placental-derived alglucerase (Ceredase<sup>®</sup>).

In patients who developed IgG antibody to **Cerezyme**<sup>®</sup>, an apparent effect on serum enzyme levels resulted in diminished volume of distribution and clearance and increased elimination half-life compared to patients without antibody (see **WARNINGS**).

## INDICATIONS AND USAGE

**Cerezyme**<sup>®</sup> (imiglucerase for injection) is indicated for long-term enzyme replacement therapy for pediatric and adult patients with a confirmed diagnosis of Type 1 Gaucher disease that results in one or more of the following conditions:

- a. anemia
- b. thrombocytopenia
- c. bone disease
- d. hepatomegaly or splenomegaly

## CONTRAINDICATIONS

There are no known contraindications to the use of **Cerezyme**<sup>®</sup> (imiglucerase for injection). Treatment with **Cerezyme**<sup>®</sup> should be carefully re-evaluated if there is significant clinical evidence of hypersensitivity to the product.

NDA 20-362/S-053

Page 5

## WARNINGS

Approximately 15% of patients treated and tested to date have developed IgG antibody to **Cerezyme**<sup>®</sup> (imiglucerase for injection) during the first year of therapy. Patients who developed IgG antibody largely did so within 6 months of treatment and rarely developed antibodies to **Cerezyme**<sup>®</sup> after 12 months of therapy. Approximately 46% of patients with detectable IgG antibodies experienced symptoms of hypersensitivity.

Patients with antibody to **Cerezyme**<sup>®</sup> have a higher risk of hypersensitivity reaction. Conversely, not all patients with symptoms of hypersensitivity have detectable IgG antibody. It is suggested that patients be monitored periodically for IgG antibody formation during the first year of treatment.

Treatment with **Cerezyme**<sup>®</sup> should be approached with caution in patients who have exhibited symptoms of hypersensitivity to the product.

Anaphylactoid reaction has been reported in less than 1% of the patient population. Further treatment with imiglucerase should be conducted with caution. Most patients have successfully continued therapy after a reduction in rate of infusion and pretreatment with antihistamines and/or corticosteroids.

## PRECAUTIONS

### General

In less than 1% of the patient population, pulmonary hypertension and pneumonia have also been observed during treatment with **Cerezyme**<sup>®</sup> (imiglucerase for injection). Pulmonary hypertension and pneumonia are known complications of Gaucher disease, and have been observed both in patients receiving and not receiving **Cerezyme**<sup>®</sup>. No causal relationship with **Cerezyme**<sup>®</sup> has been established. Patients with respiratory symptoms in the absence of fever should be evaluated for the presence of pulmonary hypertension.

Therapy with **Cerezyme**<sup>®</sup> should be directed by physicians knowledgeable in the management of patients with Gaucher disease.

Caution may be advisable in administration of **Cerezyme**<sup>®</sup> to patients previously treated with Ceredase<sup>®</sup> and who have developed antibody to Ceredase<sup>®</sup> (*alglucerase injection*) or who have exhibited symptoms of hypersensitivity to Ceredase<sup>®</sup>.

### Carcinogenesis, Mutagenesis, Impairment of Fertility

Studies have not been conducted in either animals or humans to assess the potential effects of **Cerezyme**<sup>®</sup> (imiglucerase for injection) on carcinogenesis, mutagenesis, or impairment of fertility.

### Teratogenic Effects: Pregnancy Category C

Animal reproduction studies have not been conducted with **Cerezyme**<sup>®</sup> (imiglucerase for injection). It is also not known whether **Cerezyme**<sup>®</sup> can cause fetal harm when administered to a pregnant woman, or can affect reproductive capacity. **Cerezyme**<sup>®</sup> should not be administered during pregnancy except

NDA 20-362/S-053

Page 6

when the indication and need are clear and the potential benefit is judged by the physician to substantially justify the risk.

### **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when **Cerezyme**<sup>®</sup> (imiglucerase for injection) is administered to a nursing woman.

### **Pediatric Use**

The safety and effectiveness of **Cerezyme**<sup>®</sup> (imiglucerase for injection) have been established in patients between 2 and 16 years of age. Use of **Cerezyme**<sup>®</sup> in this age group is supported by evidence from adequate and well-controlled studies of **Cerezyme**<sup>®</sup> and **Ceredase**<sup>®</sup> (αglucuronidase injection) in adults and pediatric patients, with additional data obtained from the medical literature and from long term postmarketing experience. **Cerezyme**<sup>®</sup> has been administered to patients younger than 2 years of age, however the safety and effectiveness in patients younger than 2 have not been established.

### **ADVERSE REACTIONS**

Experience in patients treated with **Cerezyme**<sup>®</sup> (imiglucerase for injection) has revealed that approximately 9.8% of patients experienced adverse events which were judged to be related to **Cerezyme**<sup>®</sup> administration and which occurred with an increase in frequency. Some of the adverse events were related to the route of administration. These include discomfort, pruritus, burning, swelling or sterile abscess at the site of venipuncture. Each of these events were found to occur in < 1% of the total patient population.

Symptoms suggestive of hypersensitivity have been noted in approximately 4.4% of patients. Onset of such symptoms has occurred during or shortly after infusions; these symptoms include pruritus, flushing, urticaria/angioedema, chest discomfort, respiratory symptoms, cyanosis and hypotension. Anaphylactoid reaction has also been reported. (see **WARNINGS**). Each of these events were found to occur in < 1% of the total patient population. Pre-treatment with antihistamines and/or corticosteroids and reduced rate of infusion have allowed continued use of **Cerezyme**<sup>®</sup> in most patients.

Additional adverse reactions that have been reported in approximately 5.4% of patients treated with **Cerezyme**<sup>®</sup> include: nausea, abdominal pain, diarrhea, rash, fatigue, headache, fever, dizziness, chills, backache, and tachycardia. Each of these events were found to occur in < 1% of the total patient population.

Since **Cerezyme**'s approval in May 1994, Genzyme has maintained a database of adverse events occurring in treated patients that were received through spontaneous reporting or discussed in the medical literature. Incidence rates cannot be calculated from the spontaneously reported adverse events. From this data base, the most commonly reported adverse events in children (defined as ages 2-12 years) included dyspnea, fever, nausea, flushing, vomiting, and coughing, whereas in adolescents (>12 – 16 years) and in adults (>16 years) the most commonly reported events included headache pruritus, and rash.

NDA 20-362/S-053

Page 7

In addition to the adverse reactions that have been observed in patients treated with **Cerezyme®**, the following adverse reactions have been reported for this therapeutic class of drug: transient peripheral edema and vomiting.

## OVERDOSE

Experience with doses up to 240 U/kg every two weeks have been reported. At that dose there have been no reports of obvious toxicity.

## DOSAGE AND ADMINISTRATION

**Cerezyme®** (imiglucerase for injection) is administered by intravenous infusion over 1-2 hours. Dosage should be individualized to each patient. Initial dosages range from 2.5 U/kg of body weight 3 times a week to 60 U/kg once every 2 weeks. 60 units/kg every 2 weeks is the dosage for which the most data are available. Disease severity may dictate that treatment be initiated at a relatively high dose or relatively frequent administration. Dosage adjustments should be made on an individual basis, and may increase or decrease, based on achievement of therapeutic goals as assessed by routine comprehensive evaluations of the patient's clinical manifestations.

**Cerezyme®** should be stored at 2-8°C (36-46°F). After reconstitution, **Cerezyme®** should be inspected visually before use. Being a protein solution, slight flocculation (described as thin translucent fibers) occurs occasionally after dilution. The diluted solution may be filtered through an in-line low protein-binding 0.2 µm filter during administration. Any vials exhibiting opaque particles or discoloration should not be used. DO NOT USE **Cerezyme®** after the expiration date on the vial.

On the day of use, after the correct amount of **Cerezyme®** to be administered to the patient has been determined, the appropriate number of vials are each reconstituted with Sterile Water for Injection, USP. The final concentrations and administration volumes are provided in the following table:

	200 Unit Vial	400 Unit Vial
Sterile water for reconstitution	5.1 mL	10.2 mL
Final volume of reconstituted product	5.3 mL	10.6 mL
Concentration after reconstitution	40 U/mL	40 U/mL
Withdrawal volume	5.0 mL	10.0 mL
Units of enzyme within final volume	200 Units	400 units

A nominal 5.0 mL for the 200 unit vial (10.0 mL for the 400 unit vial) is withdrawn from each vial. The appropriate amount of **Cerezyme®** for each patient is diluted with 0.9% Sodium Chloride Injection, USP, to a final volume of 100 to 200 mL. **Cerezyme®** is administered by intravenous infusion over 1 to 2 hours. Aseptic techniques should be used when diluting the dose. Since **Cerezyme®** does not contain any preservative, after reconstitution, vials should be promptly diluted and not stored for subsequent use. **Cerezyme®**, after reconstitution, has been shown to be stable for up to 12 hours when stored at room temperature (25°C) and at 2-8°C. **Cerezyme®**, when diluted, has been shown to be stable for up to 24 hours when stored at 2-8°C.

NDA 20-362/S-053

Page 8

Relatively low toxicity, combined with the extended time course of response, allows small dosage adjustments to be made occasionally to avoid discarding partially used bottles. Thus, the dosage administered in individual infusions may be slightly increased or decreased to utilize fully each vial as long as the monthly administered dosage remains substantially unaltered.

#### **HOW SUPPLIED**

**Cerezyme®** (imiglucerase for injection) is supplied as a sterile, non-pyrogenic, lyophilized product. It is available as follows:

200 Units per Vial NDC 58468-1983-1

400 Units per Vial NDC 58468-4663-1

Store at 2-8°C (36-46°F).

#### **Rx only**

U.S. Patent Numbers: 5,236,838

5,549,892

**Cerezyme®** (imiglucerase for injection) is manufactured by:

**Genzyme Corporation**

**One Kendall Square**

**Cambridge, MA 02139 USA**

Certain manufacturing operations may have been performed by other firms.

---

**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**

---

/s/

-----  
Mary Parks  
6/14/02 10:41:59 AM  
for Dr. Orloff