

## **Benzamycin® Topical Gel**

(erythromycin-benzoyl peroxide gel)

**Topical Gel: erythromycin (3%), benzoyl peroxide (5%)**


**For Dermatological Use Only – Not for Ophthalmic Use**

**Reconstitute Before Dispensing**

### **DESCRIPTION**


BENZAMYCIN® Topical Gel contains erythromycin [(3R\*, 4S\*, 5S\*, 6R\*, 7R\*, 9R\*, 11R\*, 12R\*, 13S\*, 14R\*)-4-[(2,6-Dideoxy-3-C-methyl-3-O-methyl- $\alpha$ -L-ribo-hexopyranosyl)-oxy]-14-ethyl-7,12,13-trihydroxy-3,5,7,9,11,13-hexamethyl-6-[[[3,4,6-trideoxy-3-(dimethylamino)- $\beta$ -D-xyl-hexopyranosyl]oxy]oxacyclotetradecane-2,10-dione]. Erythromycin is a macrolide antibiotic produced from a strain of *Saccharopolyspora erythraea* (formerly *Streptomyces erythraeus*). It is a base and readily forms salts with acids.

Chemically, erythromycin is (C<sub>37</sub>H<sub>67</sub>NO<sub>13</sub>). It has the following structural formula:



Erythromycin has the molecular weight of 733.94. It is a white crystalline powder and has a solubility of approximately 1 mg/mL in water and is soluble in alcohol at 25°C. BENZAMYCIN Topical Gel also contains benzoyl peroxide for topical use. Benzoyl peroxide is an antibacterial and keratolytic agent.

Chemically, benzoyl peroxide is (C<sub>14</sub>H<sub>10</sub>O<sub>4</sub>). It has the following structural formula:



Benzoyl peroxide has the molecular weight of 242.23. It is a white granular powder and is sparingly soluble in water and alcohol and soluble in acetone, chloroform and ether.

Each gram of BENZAMYCIN Topical Gel contains, as dispensed, 30 mg (3%) of erythromycin and 50 mg (5%) of benzoyl peroxide in a base of purified water USP, carbomer, alcohol 20%, sodium hydroxide NF, docusate sodium and fragrance.

## CLINICAL PHARMACOLOGY

The exact mechanism by which erythromycin reduces lesions of acne vulgaris is not fully known; however, the effect appears to be due in part to the antibacterial activity of the drug.

Benzoyl peroxide has a keratolytic and desquamative effect which may also contribute to its efficacy. Benzoyl peroxide has been shown to be absorbed by the skin where it is converted to benzoic acid.

## MICROBIOLOGY

Erythromycin acts by inhibition of protein synthesis in susceptible organisms by reversibly binding to 50 S ribosomal subunits, thereby inhibiting translocation of aminoacyl transfer-RNA and inhibiting polypeptide synthesis. Antagonism has been demonstrated *in vitro* between erythromycin, lincomycin, chloramphenicol and clindamycin.

Benzoyl peroxide is an antibacterial agent which has been shown to be effective against *Propionibacterium acnes*, an anaerobe found in sebaceous follicles and comedones. The antibacterial action of benzoyl peroxide is believed to be due to the release of active oxygen.

## INDICATIONS AND USAGE

BENZAMYCIN Topical Gel is indicated for the topical treatment of acne vulgaris.

## CONTRAINDICATIONS

BENZAMYCIN Topical Gel is contraindicated in those individuals who have shown hypersensitivity to any of its components.

## WARNINGS

**Pseudomembranous colitis has been reported with nearly all antibacterial agents, including erythromycin, and may range in severity from mild to life-threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhea subsequent to the administration of antibacterial agents.**

Treatment with antibacterial agents alters the normal flora of the colon and may permit overgrowth of clostridia. Studies indicate that a toxin produced by *Clostridium difficile* is one primary cause of “antibiotic-associated colitis.”

After the diagnosis of pseudomembranous colitis has been established, therapeutic measures should be initiated. Mild cases of pseudomembranous colitis usually respond to drug discontinuation alone. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation and treatment with an antibacterial drug clinically effective against *C. difficile* colitis.

## PRECAUTIONS

**General:** For topical use only; not for ophthalmic use. Concomitant topical acne therapy should be used with caution because a possible cumulative irritancy effect may occur, especially with the use of peeling, desquamating or abrasive agents. If severe irritation develops, discontinue use and institute appropriate therapy.

The use of antibiotic agents may be associated with the overgrowth of nonsusceptible organisms including fungi. If this occurs, discontinue use and take appropriate measures.

Avoid contact with eyes and all mucous membranes.

**Information for Patients:** Patients using BENZAMYCIN Topical Gel should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes, nose, mouth, and all mucous membranes.
2. This medication should not be used for any disorder other than that for which it was prescribed.
3. Patients should not use any other topical acne preparation unless otherwise directed by physician.
4. Patients should report to their physician any signs of local adverse reactions.
5. BENZAMYCIN® Topical Gel may bleach hair or colored fabric.
6. Keep product refrigerated and discard after 3 months.

### **CARCINOGENESIS, MUTAGENESIS AND IMPAIRMENT OF FERTILITY**

Data from a study using mice known to be highly susceptible to cancer suggests that benzoyl peroxide acts as a tumor promoter. The clinical significance of this is unknown.

No animal studies have been performed to evaluate the carcinogenic and mutagenic potential or effects on fertility of topical erythromycin. However, long-term (2-year) oral studies in rats with erythromycin ethylsuccinate and erythromycin base did not provide evidence of tumorigenicity. There was no apparent effect on male or female fertility in rats fed erythromycin (base) at levels up to 0.25% of diet.

**Pregnancy: Teratogenic Effects: Pregnancy CATEGORY C:** Animal reproduction studies have not been conducted with BENZAMYCIN Topical Gel or benzoyl peroxide.

There was no evidence of teratogenicity or any other adverse effect on reproduction in female rats fed erythromycin base (up to 0.25% diet) prior to and during mating, during gestation and through weaning of two successive litters.

There are no well-controlled trials in pregnant women with BENZAMYCIN Topical Gel. It also is not known whether BENZAMYCIN Topical Gel can cause fetal harm when administered to a pregnant woman or can affect reproductive capacity. BENZAMYCIN Topical Gel should be given to a pregnant woman only if clearly needed.

**Nursing Women:** It is not known whether BENZAMYCIN Topical Gel is excreted in human milk after topical application. However, erythromycin is excreted in human milk following oral and parenteral erythromycin administration. Therefore, caution should be exercised when erythromycin is administered to a nursing woman.

**Pediatric Use:** Safety and effectiveness of this product in pediatric patients below the age of 12 have not been established.

### **ADVERSE REACTIONS**

In controlled clinical trials, the incidence of adverse reactions associated with the use of BENZAMYCIN Topical Gel was approximately 3%. These were dryness and urticarial reaction.

The following additional local adverse reactions have been reported occasionally: irritation of the skin including peeling, itching, burning sensation, erythema, inflammation of the face, eyes and nose, and irritation of the eyes. Skin discoloration, oiliness and tenderness of the skin have also been reported.

#### **DOSAGE AND ADMINISTRATION**

BENZAMYCIN Topical Gel should be applied twice daily, morning and evening, or as directed by a physician, to affected areas after the skin is thoroughly washed, rinsed with warm water and gently patted dry.

#### **How Supplied and Compounding Directions:**

Size (Net Weight)	NDC 0066-	Benzoyl Peroxide Gel	Active Erythromycin Powder (In Plastic Vial)	Ethyl Alcohol (70%) To Be Added
23.3 grams (as dispensed)	0510-23	20 grams	0.8 grams	3 mL
46.6 grams (as dispensed)	0510-46	40 grams	1.6 grams	6 mL

**Prior to dispensing, tap vial until powder flows freely. Add indicated amount of ethyl alcohol (70%) to vial (to the mark) and immediately shake to completely dissolve erythromycin.** Add this solution to gel and stir until homogeneous in appearance (1 to 1½ minutes). BENZAMYCIN Topical Gel should then be stored under refrigeration. Do not freeze. Place a 3-month expiration date on the label.

**NOTE: Prior to reconstitution, store at room temperature between 15° and 30°C (59° – 86°F).**

**After reconstitution, store under refrigeration between 2° and 8°C (36° – 46°F).**

**Do not freeze. Keep tightly closed. Keep out of the reach of children.**

U.S. Patent Nos. 4,387,107 and 4,497,794.

Prescribing Information as of January 2003(a).

#### **Rx only**

Manufactured for

#### **Dermik Laboratories**

A Division of Aventis Pharmaceuticals Inc.

Berwyn, PA 19312 USA

by

Aventis Pharmaceuticals Puerto Rico Inc.

Manati, Puerto Rico 00674

**Benzamycin<sup>®</sup> Topical Gel**  
(erythromycin-benzoyl peroxide gel)






<b>PLEASE READ COMPLETE COMPOUNDING DIRECTIONS</b>
<b>NOTE:</b> TAP VIAL UNTIL ALL POWDER FLOWS FREELY. ADD ETHYL ALCOHOL (70%) TO VIAL (TO THE MARK) AND <b>IMMEDIATELY SHAKE/DISSOLVE COMPLETELY.</b>

ID number:	50067429	
Version:	B	
Country:	USA	
Date:	11-5-02	
Operator:	DM	
Logo version:	Logo not used, just text	
Product Desc.	Label, Benzamycin Gel, 23.3g	
Supplier:	CCL Label – SD	
Artwork created in:	Adobe Illustrator 8.0, Mac format	
Fonts used:	OceanSansAV Light, Light Italic, Bold	
Minimum point size of text:	5 pt.	
Colors Used:		
Process Cyan	Black	No Varn




Proofreading:	Date: _____
	Signature: _____
	Date: _____
	Signature: _____
Regulatory:	Date: _____
	Signature: _____
Functional:	Date: _____
	Signature: _____

Exp	Lot	NDC 0066-0510-23	<b>Benzamycin®</b>	<b>R, ONLY</b> After reconstitution, each gram of BENZAMYCIN® Topical Gel contains 30mg (5%) erythromycin and 50mg (5%) benzoyl peroxide in a base of purified water USP, carbomer, alcohol 20%, sodium hydroxide NF, docusate sodium and fragrance. <b>Usual Dosage:</b> Apply twice daily, morning and evening, or as directed by physician, to affected areas after skin is thoroughly washed, rinsed with warm water and gently patted dry. See package insert for full prescribing information. <b>WARNING:</b> For external use only. Keep away from eyes. Keep out of reach of children. May bleach fabric or hair. Use within 3 months after mixing. Store mixed formulation under refrigeration between 2-8°C (36-46°F). Do not freeze. Keep tightly closed.
		<b>Topical Gel</b>	<b>erythromycin-benzoyl peroxide gel</b>	Mfd for: Dermik Laboratories, A Division of Aventis Pharmaceuticals Inc. Berwyn, PA 19312 USA C2002 50067429
		FOR TOPICAL USE ONLY	DERMIK®	
		One 23.3g Jar (as dispensed)	✶Aventis	

ID number:	50067440	
Version:	B	
Country:	USA	
Date:	11-4-02	
Operator:	DM	
Logo version:	No logo used/text only	
Product Desc.	Carton, Benzamycin Gel, 23.3g Line 1	
Supplier:	PCI Services	
Artwork created in: Adobe Illustrator 8.0, Mac format		
Fonts used: OceanSansAV Bold, Light, Light Italic		
Minimum point size of text: 5 pt.		
Colors Used:		
Reflex Blue  Proc Cyan  PMS 186  No Varn 		

Proofreading:	Date: _____
	Signature: _____
Regulatory:	Date: _____
	Signature: _____
Functional:	Date: _____
	Signature: _____



ID number: 50067253

Version: B

Country: USA

Date: 10-8-02

Operator: DM

Logo version: A1


Product Desc. Label, Benzamycin Gel, 46.6g




Supplier: CCL Label – SD

Artwork created in: Adobe Illustrator 8.0, Mac format

Fonts used: OceanSansAV Light, Light Italic, Bold

Minimum point size of text: 5 pt.



Colors Used:  
Process Cyan  Black  No Varn 

Proofreading:      Date: \_\_\_\_\_

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Signature: \_\_\_\_\_

Regulatory:      Date: \_\_\_\_\_

Signature: \_\_\_\_\_

Functional:      Date: \_\_\_\_\_

Signature: \_\_\_\_\_

Exp

Lot

NDC 0066-0510-46

**Benzamycin®**

**Topical Gel**

*erythromycin-benzoyl peroxide gel*

FOR TOPICAL USE ONLY

One 46.6g Jar (as dispensed)


DERMIK

Aventis

**Benzamycin® Topical Gel** *(erythromycin-benzoyl peroxide gel)*

After reconstitution, each gram of BENZAMYCIN® Topical Gel contains 30mg (3%) erythromycin and 50mg (5%) benzoyl peroxide in a base of purified water USP, carbomer, alcohol 20%, sodium hydroxide NF, docusate sodium and fragrance. **Usual Dosage:** Apply twice daily, morning and evening, or as directed by physician, to affected areas after skin is thoroughly washed, rinsed with warm water and gently patted dry. See package insert for full prescribing information. **WARNING:** For external use only. Keep away from eyes. Keep out of reach of children. May bleach fabric or hair. Use within 3 months after mixing. Store mixed formulation under refrigeration between 2-8°C (36-46°F). **Do not freeze. Keep tightly closed.** U.S. Patents 4,387,107 and 4,497,794.

Mfd for: Dermik Laboratories, A Division of Aventis Pharmaceuticals Inc.  
Berwyn, PA 19312 USA ©2002  
Mfd by: **Aventis Pharmaceuticals Puerto Rico Inc.**, Manati, Puerto Rico 00674



ONLY

50067253



**Proofreading:**      **Date:** \_\_\_\_\_  
                                  **Signature:** \_\_\_\_\_  
                                  **Date:** \_\_\_\_\_  
                                  **Signature:** \_\_\_\_\_

**Regulatory:**      **Date:** \_\_\_\_\_  
                                  **Signature:** \_\_\_\_\_

**Functional:**      **Date:** \_\_\_\_\_  
                                  **Signature:** \_\_\_\_\_



**Colors Used:**

Reflex Blue		PMS 186		Proc Cyan		No Varn	
-------------	---	---------	---	-----------	---	---------	---

