

NDC 0187-0380-34

Benoquin® Cream 20%
(Monobenzone Cream USP)

CAUTION: Federal (U.S.A.) law prohibits dispensing without prescription.
NET WT. 1 1/4 OZ. (35.44 grams)

ICN

Ingredients: Each gram of Benoquin Cream 20% contains 200 mg of monobenzone USP, in a water-washable base consisting of purified water USP, cetyl alcohol NF, propylene glycol USP, sodium lauryl sulfate NF and white wax NF.
Directions: Apply and rub into the pigmented area two or three times daily, or as directed by physician. Prolonged exposure to sunlight should be avoided during treatment with Benoquin Cream 20%, or a sunscreen should be used.
Indications: For final depigmentation in extensive vitiligo, and to permanently depigment normal skin surrounding vitiliginous lesions in patients with disseminated (greater than 50% of body surface area) idiopathic vitiligo.
FOR EXTERNAL USE ONLY: Keep out of the reach of children. See package insert for full information. To open: pierce sealed end with cap.



N 3 01870 38034 7

NDC 0187-0380-34

Benoquin® Cream 20%
(Monobenzone Cream USP)

CAUTION: Federal (U.S.A.) law prohibits dispensing without prescription.
NET WT. 1 1/4 OZ. (35.44 grams)

ICN

Warning: Benoquin is a potent depigmenting agent. Store at controlled room temperature (15°-30°C) 59°-86°F.
Manufactured by ICN Pharmaceuticals, Inc. 3300 Hyland Ave., Costa Mesa, CA 92626 U.S.A.

1197-00EF
Rev. 7-97

PROOF OF PURCHASE

119700



ICN

ICN

BENOQUIN[®] CREAM 20%


(Monobenzone Cream, USP)

R_x Only.

FOR EXTERNAL USE ONLY

DESCRIPTION:

Monobenzone is the monobenzyl ether of hydroquinone. Monobenzone occurs as a white, almost tasteless crystalline powder, soluble in alcohol and practically insoluble in water. Chemically, monobenzone is designated as p-(benzyloxy) phenol; the empirical formula is C₁₃H₁₂O₂; molecular weight 200.24. The structural formula is:



C₁₃H₁₂O₂ 200.24

Each gram of Benoquin Cream contains 200 mg of monobenzone USP, in a water-washable base consisting of purified water USP, cetyl alcohol NF, propylene glycol USP, sodium lauryl sulfate NF and white wax NF.

CLINICAL PHARMACOLOGY:

Benoquin Cream 20% is a depigmenting agent whose mechanism of action is not fully understood.

The topical application of monobenzone in animals, increases the excretion of melanin from the melanocytes. The same action is thought to be responsible for the depigmenting effect of the drug in humans. Monobenzone may cause destruction of melanocytes and permanent depigmentation. This effect is erratic and may take one to four months to occur while existing melanin is lost with normal sloughing of the stratum corneum. Hyperpigmented skin appears to fade more rapidly



than does normal skin, and exposure to sunlight reduces the depigmenting effect of the drug. The histology of the skin after depigmentation with topical monobenzone is the same as that seen in vitiligo; the epidermis is normal except for the absence of identifiable melanocytes.

INDICATIONS AND USAGE:

Benoquin Cream 20% is indicated for final depigmentation in extensive Vitiligo.

Benoquin Cream 20% is applied topically to permanently depigment normal skin surrounding vitiliginous lesions in patients with disseminated (greater than 50 percent of body surface area) idiopathic vitiligo.

Benoquin Cream 20% is not recommended in freckling; hyperpigmentation caused by photosensitization following the use of certain perfumes (berlock dermatitis); melasma (chloasma) of pregnancy; or hyperpigmentation resulting from inflammation of the skin. Benoquin Cream 20% is not effective for the treatment of cafe-au-lait spots, pigmented nevi, malignant melanoma or pigmentation resulting from pigments other than melanin (e.g.: bile, silver, or artificial pigments).

CONTRAINDICATIONS:

Benoquin Cream 20% contains a potent depigmenting agent and is not a cosmetic skin bleach. Use of Benoquin Cream 20% is contraindicated in any conditions other than disseminated vitiligo. Benoquin Cream 20% frequently produces irreversible depigmentation, and it must not be used as a substitute for hydroquinone.

Benoquin Cream 20% is also contraindicated in individuals with a history of sensitivity or allergic reactions to this product, or any of its ingredients.

WARNINGS:

Benoquin Cream 20% is a potent depigmenting agent, not a mild cosmetic bleach. Do not use except for final depigmentation in extensive vitiligo.



Keep this, and all medications out of the reach of children. In case of accidental ingestion, call a physician or a Poison Control Center immediately.

PRECAUTIONS (See Warnings):

General. Benoquin Cream 20% is for External Use Only. Following therapy with Benoquin Cream 20%, the skin will be sensitive for the rest of the patient's life. He/she must use sunscreens during exposure to the sun.

Information for the Patient. Benoquin Cream 20% contains a potent depigmenting agent and is not a cosmetic skin bleach. Use of Benoquin Cream 20% is contraindicated in any conditions other than disseminated vitiligo. Use only for final depigmentation in extensive vitiligo. Areas of normal skin distant to the site of Benoquin Cream 20% application may become depigmented, and irregular, excessive, unsightly, and frequently permanent depigmentation may occur.

Carcinogenesis, mutagenesis, impairment of fertility. No long term studies have been performed to evaluate carcinogenic potential.

Pregnancy: Category C. Animal reproduction studies have not been conducted with Benoquin Cream 20%. It is also not known whether Benoquin Cream 20% can cause fetal harm when administered to a pregnant woman, or can affect reproduction capacity. Benoquin Cream 20% should be given to a pregnant woman only if clearly needed.

Nursing Mothers. It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Benoquin Cream 20% is administered to a nursing woman.

Pediatric Use. The safety and effectiveness of Benoquin Cream 20% in pediatric patients below the age of 12 years have not been established.

ADVERSE REACTIONS:

Mild, transient skin irritation and sensitization, including erythematous and eczematous reactions have occurred

following topical application of Benoquin Cream 20%. Although those reactions are usually transient, treatment with Benoquin Cream 20% should be discontinued if irritation, a burning sensation, or dermatitis occur. Areas of normal skin distant to the site of Benoquin Cream 20% application frequently have become depigmented, and irregular, excessive, unsightly, and frequently permanent depigmentation has occurred.

DOSAGE AND ADMINISTRATION:

A thin layer of Benoquin Cream 20% should be applied and rubbed into the pigmented area two or three times daily, or as directed by physician. Prolonged exposure to sunlight should be avoided during treatment with Benoquin Cream 20%, or a sunscreen should be used.

Depigmentation is usually accomplished after one to four months of Benoquin Cream 20% treatment. If satisfactory results are not obtained after four months of Benoquin Cream 20% treatment, the drug should be discontinued. When the desired degree of depigmentation is obtained, Benoquin Cream 20% should be applied only as often as needed to maintain depigmentation (usually only two times weekly).

HOW SUPPLIED:

Benoquin Cream 20% in 1 1/4 oz. tubes (35.4 g)
(NDC 0187-0380-34).

Benoquin Cream 20% should be stored at 25°C (77°F);
excursion permitted to 15°C - 30°C (59°F - 86°F).



ICN Pharmaceuticals, Inc.
3300 Hyland Ave.
Costa Mesa, CA 92626
(714)545-0100

2393-05 EL
Rev. 8-00