

Bullets – 5 pt

Bar Line – 1 pt

Body Copy – 6 pt/6.5 pt leading
(Helvetica Neu Condensed – 88% Horizontally Scaled)

STANDARD FORMAT

Date: December 11, 2002rev. **File Name:** 50.GT8267-48-00.Proposal
(Helvetica Neu Condensed Bold Oblique – 100% Horizontal Scaled)

Drug Fact Headers – 8.5pts

(Helvetica Neu Condensed Bold Oblique)

Headers – 8 pts

(Helvetica Neu Condensed Bold Oblique)

SubHeaders – 6 pts

(Helvetica Neu Condensed Bold)

Body Copy – 6/6.5pts

(Helvetica Neu Condensed Regular)

Drug Facts (continued)
Ask a doctor before use if you have

- never had migraines diagnosed by a health professional
- a headache that is different from your usual migraines
- the worst headache of your life
- fever and stiff neck
- headaches beginning after or caused by head injury, exertion, coughing or bending
- experienced your first headache after the age of 50
- daily headaches
- bleeding problems
- stomach problems such as heartburn, upset stomach, or stomach pain that do not go away or recur
- migraines so severe as to require bed rest
- problems or serious side effects from taking pain relievers or blood thinners

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for:
 - anticoagulation (thinning of the blood)
 - diabetes
 - gout
 - arthritis
- an allergic reaction occurs. Seek medical help right away.
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely decided to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)
Keep out of reach of children.
Overdose warning: Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)
- adults take 2 tablets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 tablets in 24 hours, unless directed by a doctor
- under 18 years of age, ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- read all product information before using. Keep this box for important information.

Inactive ingredients: benzoic acid, carnuba wax, FD&C Blue #1, hydropropylcellulose, hypromellose, microcrystalline cellulose, mineral oil, polyacrylate 20, polyoxone, propylene glycol, simethicone emulsion, sorbitan monoleurate, stearic acid, titanium dioxide
*may contain these ingredients

Questions or comments? 1-800-468-7146

DISTR. BY: BRISTOL-MYERS SQUIBB CO., NY, NY 10154
© 1995, MADE IN USA FPO USE UNDER U.S. PATENT NO. 5972916

Drug Facts
Active Ingredients (in each tablet)
Acetaminophen 250 mg
Aspirin 250 mg
Caffeine 65 mg

Use • treats migraine

Warnings
Rhys's syndrome: Children and teenagers should not use this drug for chicken pox, or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
• facial swelling • asthma (wheezing) • shock
Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen and aspirin or other pain relievers/fever reducers. Acetaminophen and aspirin may cause liver damage and stomach bleeding.
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer
• with any other products containing acetaminophen.
Taking more than directed may cause liver damage.

Purposes
Pain reliever
Pain reliever
Pain reliever and

Drug Fact Header – 8.5 pts
(Helvetica Neu Condensed Bold Oblique – 100% Horizontally Scaled)

Headers – 8 pts
(Helvetica Neu Condensed Bold Oblique – 88% Horizontally Scaled)

Hair Line – .5 pt

