

Front Panel - Labeling Format (100%) for ChloroPrep® One-Step
Chlorhexidine Gluconate 2% w/v and Isopropyl Alcohol 70% v/v
Patient Preoperative Skin Preparation • 10.5-mL Applicator

CREATED 5/01/02 REVISED 6/10/02 REVISED 8/20/02 REVISED 1/30/03
REVISED 2/3/03 (annotations) REVISED 2/11/03 REVISED 2/13/03 REVISED 2/17/03

REDUCED 65% TO FIT

Number:.....
14 pt antique
olive compact
Title:
40 pt antique
olive nord
Formulation:.....
15 pt antique
olive compact
Statements:.....
18 pt antique
olive compact
Applicator:.....
15 pt antique
olive bold
Single use:.....
15 pt antique
olive roman
Applicator, etc.:.....
15 pt antique
olive bold
"Sterile"
15 pt antique
olive nord
Storage, etc.:.....
15 pt antique
olive roman

Cat. No. 260700 NDC #054365-400-04 DIN#02160757

ChloroPrep® One-Step

Chlorhexidine Gluconate 2% (w/v) and Isopropyl Alcohol 70% (v/v)
Antiseptic Patient Preoperative Skin Preparation

**ChloroPrep® One-Step is a fast-acting, broad spectrum,
persistent antiseptic that significantly reduces the
number of microorganisms on intact skin.**

10.5-mL Applicator

Single Use


Applicator is sterile if package is intact

Contents: 1 Each Sponge, Latex Free

Quantity: 1 Box, 25 Each/Box

Store at room temperature 20-25 °C (68-77 °F)

Avoid freezing and excessive heat above 40 °C (104 °F)



Back Panel - Labeling Format (100%) for ChlorPrep® One-Step
 Chlorhexidine Gluconate 2% w/v and Isopropyl Alcohol 70% v/v
 Patient Preoperative Skin Preparation • 10.5-mL Applicator

CREATED 5/01/02 REVISED 6/10/02 REVISED 8/20/02 REVISED 1/30/03 REVISED 2/3/03 (annotations)
 REVISED 2/11/03 REVISED 2/13/03 REVISED 2/17/03

REDUCED 65% TO FIT

ChlorPrep® One-Step

Chlorhexidine Gluconate 2% w/v and Isopropyl Alcohol 70% v/v

Patient Preoperative Skin Preparation • 10.5-mL Applicator

WARNING: FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.
DO NOT USE WITH ELECTROCAUTERY PROCEDURES.

Title:
 14 pt Helvetica
 bold italic
 Headings:
 8 pt Helvetica
 bold italic
 left justified
 Subheadings:
 6 pt Helvetica
 bold,
 left justified
 0.5 point headline
 2.5 point
 headline
 2.5 point box
 headline
 Bullets:
 solid square
 Body Text:
 6 pt Helvetica
 regular,
 6.5 leading,
 left justified


Drug Facts	Purpose Antiseptic
Active Ingredients Chlorhexidine gluconate 2% w/v Isopropyl Alcohol 70% v/v	
Use For the preparation of the patient's skin prior to surgery	
Warnings For external use only Keep away from fire or flame. Do not use with electrocautery procedures. Do not use: <ul style="list-style-type: none"> on children less than 2 months of age because of the potential for excessive skin irritation and increased drug absorption on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol on open skin wounds or as a preservative for ointments When using this product: <ul style="list-style-type: none"> Keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cool water right away and contact a physician Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. 	
Directions Rub on the application to break the ampoule and release the antiseptic. Do not touch the sponge. Hold the sponge over the application site for approximately 20 seconds. Dry surgical sites (such as abdomen or arm). Use repeated back-and-forth motions of the sponge for approximately 20 seconds. Do not blot or wipe away. Completely wet the treatment area with antiseptic. Allow the area to air dry for approximately 2 minutes. * Natural treatment area for one applicator is approximately 457 cm ² (approximately 8.4 x 8.4 inches). Discard the applicator after a single use.	
Other Information Store between 20-25 °C (68-77 °F). Avoid freezing and excessive heat above 40 °C (104 °F). * USP purified water	
Questions? Call 1-800-333-0022 (8 a.m.-5 p.m. CST)	



Side Panels - Labeling Format (100%) for Chloraprep® One-Step
Chlorhexidine Gluconate 2% w/v and Isopropyl Alcohol 70% v/v
Patient Preoperative Skin Preparation • 10.5-mL Applicator

CREATED 5/01/02 REVISED 6/10/02 REVISED 8/20/02 REVISED 1/30/03 REVISED 2/3/03 (annotations)
REVISED 2/11/03 REVISED 2/13/03 REVISED 2/17/03


REDUCED 65% TO FIT



MEDI-FLEX®

Chloraprep® One-Step
10.5-ml Applicators • 25 qty.

Store at room temperature 20-25 °C (68-77 °F)
Avoid freezing and excessive heat above 40 °C (104 °F)



MEDI-FLEX®

Chloraprep® One-Step
10.5-ml Applicators • 25 qty.

Store at room temperature 20-25 °C (68-77 °F)
Avoid freezing and excessive heat above 40 °C (104 °F)

Lot No: Exp. Date:


•H9852607002B•

REDUCED 65% TO FIT

Top Panel - Labeling Format (100%) for Chloraprep® One-Step
Chlorhexidine Gluconate 2% w/v and Isopropyl Alcohol 70% v/v
Patient Preoperative Skin Preparation • 10.5-mL Applicator

CREATED 5/01/02 REVISÉD 6/10/02 REVISÉD 8/20/02 REVISÉD 1/30/03 REVISÉD 2/3/03 (annotations)
REVISÉD 2/11/03 REVISÉD 2/13/03 REVISÉD 2/17/03

P.O. Box 42825
Overland Park, KS 66282-2825
1-800-523-0502



Cat. No. 260700 NDC #054365-400-04 DIN#02160757

Chloraprep® One-Step

Chlorhexidine Gluconate 2% (w/v) and Isopropyl Alcohol 70% (v/v)
Antiseptic Patient Preoperative Skin Preparation

Chloraprep® One-Step is a fast-acting, broad spectrum,
persistent antiseptic that significantly reduces the
number of microorganisms on intact skin.

immediate container labeling



Cat. No. 260700 NDC #054365-400-04 DIN#02160757
ChloroPrep® One-Step
 Chlorhexidine Gluconate 2% w/v and Isopropyl Alcohol 70% v/v
 Patient Preoperative Skin Preparation • 10.5mL Applicator
WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.
DO NOT USE WITH ELECTROCAUTERY PROCEDURES.

CREATED 5/01/02 REVISED 6/10/02 REVISED 8/20/02 REVISED 1/30/03 REVISED 2/3/03
 (annotations) REVISED 2/11/03 REVISED 2/13/03 REVISED 2/17/03 REVISED 2/3/03

Applicator barrel artwork - Labeling Format (100%) for ChloroPrep® One-Step
 Chlorhexidine Gluconate 2% w/v and Isopropyl Alcohol 70% v/v
 Patient Preoperative Skin Preparation • 10.5-mL Applicator
 CREATED 5/01/02 REVISED 6/10/02 REVISED 8/20/02 REVISED 1/30/03
 REVISED 2/3/03 (annotations) REVISED 2/11/03 REVISED 2/13/03 REVISED 2/17/03

Drug Facts	
Active Ingredients Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	Purpose Antiseptic Antiseptic
Use for the preparation of the patient's skin prior to surgery	
Warnings For external use only Flammable. Keep away from fire or flame. Do not use with electrocautery procedures. Do not use <ul style="list-style-type: none"> in children less than 2 months of age because of the potential for excessive skin irritation and increased drug absorption on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol for lumbar puncture or in contact with the meninges on open skin wounds or as a general skin cleanser When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a physician. Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions <ul style="list-style-type: none"> pinch the wings on the applicator to break the ampule and release the antiseptic. Do not touch the sponge. Wet the sponge by repeatedly pressing and releasing the sponge against the treatment area until liquid is visible on the skin. dry surgical sites (such as abdomen or arm): Use repeated back-and-forth strokes of the sponge for approximately 30 seconds. Completely wet the treatment area with antiseptic. Allow the area to air dry for approximately 30 seconds. Do not blot or wipe away. moist surgical sites (such as the inguinal fold): Use repeated back-and-forth strokes of the sponge for approximately 2 minutes. Completely wet the treatment area with antiseptic. Allow the area to air dry for approximately one (1) minute. Do not blot or wipe away. maximal treatment area for one applicator is approximately 457 cm² (approximately 8.4 x 8.4 inches). Discard the applicator after a single use. Other information • store between 20-25 °C (68-77 °F) • avoid freezing and excessive heat above 40 °C (104 °F) Inactive ingredients • USP purified water Questions? Call 1-800-823-0902 (8 a.m.-5 p.m. CST)	

Title: 14 pt helvetica bold italic
 Headings: 8 pt helvetica bold italic, left justified
 0.5 point hairline
 Subheadings: 6 pt helvetica bold, left justified
 Bullet: 5 pt zapf dingbats, solid square

2.5 point barline
 2.5 point box barline
 Body Text: 6 pt helvetica regular, 6.5 leading, left justified

Single Use
 Applicator is **sterile** if package is intact

Lot No: _____ Exp. Date: _____



ChloroPrep® 10.5-mL Applicator
 2% w/v Chlorhexidine gluconate
 and 70% v/v Isopropyl alcohol
 Medi-Flex Hospital Products, Overland Park KS
 Lot No: _____ Exp. Date: _____

Title: 9 pt ontrique olive nord
 Font: 8 pt ontrique olive bold

REDUCED 65% TO FIT