

150%

<p><b>Advil</b><sup>®</sup> 1 Tablet MULTI-SYMPPTOM COLD IBUPROFEN 200 mg/PSEUDOEPHEDRINE HCl 30 mg/ CHLORPHENIRAMINE MALEATE 2 mg ALLERGY, BLEEDING AND ALCOHOL WARNING. READ CARTON BEFORE USE. Wyeth Consumer Healthcare, Madison, NJ 07940</p> <p>LOT EXP SEPARATE FIRST THEN PEEL &amp; PUSH</p>	<p><b>Advil</b><sup>®</sup> 1 Tablet MULTI-SYMPPTOM COLD IBUPROFEN 200 mg/PSEUDOEPHEDRINE HCl 30 mg/ CHLORPHENIRAMINE MALEATE 2 mg ALLERGY, BLEEDING AND ALCOHOL WARNING. READ CARTON BEFORE USE. Wyeth Consumer Healthcare, Madison, NJ 07940</p> <p>LOT EXP SEPARATE FIRST THEN PEEL &amp; PUSH</p>
<p>SEPARATE FIRST THEN PEEL &amp; PUSH</p> <p><b>Advil</b><sup>®</sup> 1 Tablet MULTI-SYMPPTOM COLD IBUPROFEN 200 mg/PSEUDOEPHEDRINE HCl 30 mg/ CHLORPHENIRAMINE MALEATE 2 mg ALLERGY, BLEEDING AND ALCOHOL WARNING. READ CARTON BEFORE USE. Wyeth Consumer Healthcare, Madison, NJ 07940</p> <p>LOT EXP SEPARATE FIRST THEN PEEL &amp; PUSH</p>	<p>SEPARATE FIRST THEN PEEL &amp; PUSH</p> <p><b>Advil</b><sup>®</sup> 1 Tablet MULTI-SYMPPTOM COLD IBUPROFEN 200 mg/PSEUDOEPHEDRINE HCl 30 mg/ CHLORPHENIRAMINE MALEATE 2 mg ALLERGY, BLEEDING AND ALCOHOL WARNING. READ CARTON BEFORE USE. Wyeth Consumer Healthcare, Madison, NJ 07940</p> <p>LOT EXP SEPARATE FIRST THEN PEEL &amp; PUSH</p>
<p>SEPARATE FIRST THEN PEEL &amp; PUSH</p> <p><b>Advil</b><sup>®</sup> 1 Tablet MULTI-SYMPPTOM COLD IBUPROFEN 200 mg/PSEUDOEPHEDRINE HCl 30 mg/ CHLORPHENIRAMINE MALEATE 2 mg ALLERGY, BLEEDING AND ALCOHOL WARNING. READ CARTON BEFORE USE. Wyeth Consumer Healthcare, Madison, NJ 07940</p> <p>LOT EXP</p>	<p>SEPARATE FIRST THEN PEEL &amp; PUSH</p> <p><b>Advil</b><sup>®</sup> 1 Tablet MULTI-SYMPPTOM COLD IBUPROFEN 200 mg/PSEUDOEPHEDRINE HCl 30 mg/ CHLORPHENIRAMINE MALEATE 2 mg ALLERGY, BLEEDING AND ALCOHOL WARNING. READ CARTON BEFORE USE. Wyeth Consumer Healthcare, Madison, NJ 07940</p> <p>LOT EXP</p>
<p>SEPARATE FIRST THEN PEEL &amp; PUSH</p> <p><b>Advil</b><sup>®</sup> 1 Tablet MULTI-SYMPPTOM COLD IBUPROFEN 200 mg/PSEUDOEPHEDRINE HCl 30 mg/ CHLORPHENIRAMINE MALEATE 2 mg ALLERGY, BLEEDING AND ALCOHOL WARNING. READ CARTON BEFORE USE. Wyeth Consumer Healthcare, Madison, NJ 07940</p> <p>LOT EXP</p>	<p>SEPARATE FIRST THEN PEEL &amp; PUSH</p> <p><b>Advil</b><sup>®</sup> 1 Tablet MULTI-SYMPPTOM COLD IBUPROFEN 200 mg/PSEUDOEPHEDRINE HCl 30 mg/ CHLORPHENIRAMINE MALEATE 2 mg ALLERGY, BLEEDING AND ALCOHOL WARNING. READ CARTON BEFORE USE. Wyeth Consumer Healthcare, Madison, NJ 07940</p> <p>LOT EXP</p>
<p>SEPARATE FIRST THEN PEEL &amp; PUSH</p> <p><b>Advil</b><sup>®</sup> 1 Tablet MULTI-SYMPPTOM COLD IBUPROFEN 200 mg/PSEUDOEPHEDRINE HCl 30 mg/ CHLORPHENIRAMINE MALEATE 2 mg ALLERGY, BLEEDING AND ALCOHOL WARNING. READ CARTON BEFORE USE. Wyeth Consumer Healthcare, Madison, NJ 07940</p> <p>LOT EXP</p>	<p>SEPARATE FIRST THEN PEEL &amp; PUSH</p> <p><b>Advil</b><sup>®</sup> 1 Tablet MULTI-SYMPPTOM COLD IBUPROFEN 200 mg/PSEUDOEPHEDRINE HCl 30 mg/ CHLORPHENIRAMINE MALEATE 2 mg ALLERGY, BLEEDING AND ALCOHOL WARNING. READ CARTON BEFORE USE. Wyeth Consumer Healthcare, Madison, NJ 07940</p> <p>LOT EXP</p>

SP1082-1.FDA01



128 HD CODE AREASUBSET  
C-10 MIL DENSITY WITH 4  
OR 6 DIGITS (CODE TO  
EXTEND TO EDGE OF  
FLAP - UNVARNISHED)

UNPRINTED AREA FOR LOT & EXP  
CODING (UNVARNISHED)

Job Information	Color Legend
<b>Date</b> 3/31/04 PASS #4 <b>Component #</b> UK24501-1.FDA02 <b>Drawing #</b> 5181	
Identification	<b>NOTE:</b> <b>RUBINE RED INK PRINTS IN PLACE OF MAGENTA</b>
<b>Wyeth</b> Consumer Healthcare Production Packaging Dept. Five / Farms Madison, NJ 07940 Phone: (973) 660-5595/5153 Fax: (973) 660-6048	
Vendor Identification	Printer Notes
Production Packaging Dept. Cheng Lin SC Linda Hill 11/25/3	PRINTER TO REPLACE FPO UPC'S WITH HI-RES DIGITAL ART ALSO PRINTER TO SUPPLY COLOR MATCHPRINT FOR COLOR APPROVAL PRIOR TO PRESS RUN. <b>RUBINE RED INK PRINTS IN PLACE OF MAGENTA</b>

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS Title	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE AFTER BULLETED SECTION	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

DEPARTMENT	REVIEWED BY		DATE	with Changes
	Print Name	Signature		
Graphic Artist				<input type="checkbox"/>
Packaging Engineer				<input type="checkbox"/>
Labeling Compliance				<input type="checkbox"/>
Medical				<input type="checkbox"/>
Regulatory Affairs				<input type="checkbox"/>
Marketing				<input type="checkbox"/>
Legal				<input type="checkbox"/>
Trademark				<input type="checkbox"/>

RELEASED	
(1)	_____ Date _____
(2)	_____ Date _____

**DIGITAL MECHANICAL**  
 COLOR REPRESENTATION MAY BE OFF  
 FOLLOW COLOR LEGEND FOR ACTUAL COLOR.

## ATTACHMENT 1

### ADVIL<sup>®</sup> Multisymptom Cold Caplets

**Ibuprofen 200 mg >> Pain reliever / Fever reducer**  
**Pseudoephedrine HCl 30 mg >> Nasal Decongestant**  
**Chlorpheniramine Maleate 2 mg >> Antihistamine**

#### **Active ingredients (in each caplet)**

#### **Purposes**

Chlorpheniramine maleate 2mg.....Antihistamine  
Ibuprofen 200 mg.....Pain reliever/fever reducer  
Pseudoephedrine HCl 30 mg.....Nasal decongestant

#### **Uses**

temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:

- runny nose
- sneezing
- headache
- itchy, watery eyes
- nasal congestion
- minor aches and pains
- itching of the nose or throat
- sinus pressure
- fever

#### **Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** Taking more than recommended may cause stomach bleeding.

**Alcohol warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take ibuprofen or other pain relievers/fever reducers. Ibuprofen may cause stomach bleeding.

#### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### **Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- heart disease
- kidney disease
- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or pain
- trouble urinating due to an enlarged prostate gland
- high blood pressure
- ulcers
- thyroid disease
- bleeding problems
- diabetes
- glaucoma

#### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking sedatives or tranquilizers
- over 65 years of age
- taking any other product that contains ibuprofen, or any other pain reliever/fever reducer
- taking any other product that contains pseudoephedrine, chlorpheniramine or any other nasal decongestant or antihistamine
- taking a prescription drug for anticoagulation (blood thinning), or a diuretic

- taking any other drug
- taking aspirin for cardioprotection

**When using this product**

- **do not use more than directed**
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- drowsiness may occur
- take with food or milk if stomach upset occurs
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- nasal congestion lasts for more than 7 days
- fever lasts for more than 3 days
- you get nervous, dizzy, or sleepless
- symptoms continue or get worse
- stomach pain occurs with the use of this product even if mild pain persists
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use this product during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults: take 1 caplet every 4-6 hours while symptoms persist.
- do not take more than 6 caplets in any 24-hour period, unless directed by a doctor
- children under 12 years of age: consult a doctor

**Other information**

- read all warnings and directions before use. Keep carton.
- store in a dry place 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**

carnauba wax, croscarmellose sodium, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, glyceryl behenate, hypromellose, iron oxide black, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, silicon dioxide, starch, titanium dioxide