

NDC 63824-057-82

Mucinex^D
600 mg guaifenesin and
60 mg pseudoephedrine HCl per tablet

**EXPECTORANT AND
NASAL DECONGESTANT**

2 Bi-layer Tablets

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

Contains FD&C Yellow Dye #6

PHYSICIAN SAMPLE-
NOT TO BE SOLD

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155
US Pat. 6,372,252 B1
3105702 102003

3-6382405782-8

4 1/4"

15/16"

NDC 63824-057-20

Mucinex^D
600 mg guaifenesin and
60 mg pseudoephedrine HCl per tablet

**EXPECTORANT AND
NASAL DECONGESTANT**



20 Bi-layer Tablets

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

See carton for complete Drug Facts.

US Pat. 6,372,252 B1

102003
3005720

Contains FD&C Yellow Dye #6

*Please visit our Web site:
www.mucinex.com*

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155
© 2003 Adams Laboratories, Inc.

5 1/4"

2"

NDC 63824-057-40

Mucinex^D
600 mg guaifenesin and
60 mg pseudoephedrine HCl per tablet

EXPECTORANT AND
NASAL DECONGESTANT



40 Bi-layer Tablets

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

See carton for complete Drug Facts.

US Pat. 6,372,252 B1

102003
3005740

Contains FD&C Yellow Dye #6

*Please visit our Web site:
www.mucinex.com*

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155
© 2003 Adams Laboratories, Inc.

5 1/4"

2"

NDC 63824-034-82

Mucinex^D
1200 mg guaifenesin and
120 mg pseudoephedrine HCl per tablet

**EXPECTORANT AND
NASAL DECONGESTANT**

2 Bi-layer Tablets

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

PHYSICIAN SAMPLE-
NOT TO BE SOLD

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155
US Pat. 6,372,252 B1
3103402 102003

3-63824-03482-9

4 1/4"

15/16"

NDC 63824-034-01

Mucinex^D
1200 mg guaifenesin and
120 mg pseudoephedrine HCl per tablet

**EXPECTORANT AND
NASAL DECONGESTANT**



10 Bi-layer Tablets

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

See carton for complete Drug Facts.

US Pat. 6,372,252 B1

102003
3003401

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155
© 2003 Adams Laboratories, Inc.

*Please visit our Web site:
www.mucinex.com*

5 1/4"

2"

NDC 63824-034-20

Mucinex[®]D
1200 mg guaifenesin and
120 mg pseudoephedrine HCl per tablet

**EXPECTORANT AND
NASAL DECONGESTANT**



20 Bi-layer Tablets

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

See carton for complete Drug Facts.

US Pat. 6,372,252 B1

102003
3003420

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155
© 2003 Adams Laboratories, Inc.

*Please visit our Web site:
www.mucinex.com*

5 1/4"

2"

MUCINEX D 600/60 mg SAMPLE TRAY

Title:
14 pt. Helvetica Bold
Italic, left justified

Headings:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Subheadings:
6 pt. Helvetica Bold
left justified

Graphic leading
to next panel

Drug Facts		Drug Facts (continued)	
Active ingredients (in each extended-release tablet)		When using this product	
Guaifenesin 600 mg	Expectorant	<ul style="list-style-type: none"> ■ do not use more than directed 	
Pseudoephedrine HCl 60 mg	Nasal Decongestant	Stop use and ask a doctor if <ul style="list-style-type: none"> ■ you get nervous, dizzy, or sleepless ■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness. 	
Uses		If pregnant or breast-feeding , ask a health professional before use.	
<ul style="list-style-type: none"> ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily relieves nasal congestion due to: <ul style="list-style-type: none"> ■ common cold ■ hay fever ■ upper respiratory allergies ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage ■ temporarily relieves sinus congestion and pressure 		Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Warnings		Directions	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.		<ul style="list-style-type: none"> ■ do not crush, chew, or break tablet ■ take with a full glass of water ■ this product can be administered without regard for timing of meals ■ adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours ■ children under 12 years of age: do not use 	
Ask a doctor before use if you have		Other information	
<ul style="list-style-type: none"> ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough accompanied by too much phlegm (mucus) 		<ul style="list-style-type: none"> ■ tamper evident: do not use if seal on bottle printed "SEALED FOR YOUR PROTECTION" is broken or missing ■ store at 20-25°C (68-77°F) ■ see bottom of bottle for lot code and expiration date 	
Inactive ingredients		Inactive ingredients	
carbomer 934P, NF; FD&C yellow #6 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF		carbomer 934P, NF; FD&C yellow #6 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF	

Right justified

Title for continued panel:
8 pt. Helvetica Bold Italic

8 pt. Helvetica Regular

0.5 pt. hairline

2.5 pt. barline

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline

1 3/4"
3 1/4"

20 BI-LAYER TABLETS

Mucinex^D
600 mg guaifenesin and
60 mg pseudoephedrine HCl per tablet
EXPECTORANT AND
NASAL DECONGESTANT

NEW

Drug Facts (continued)

When using this product
 ■ do not use more than directed

Stop use and ask a doctor if
 ■ you get nervous, dizzy, or sleepless
 ■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ do not crush, chew, or break tablet
 ■ take with a full glass of water
 ■ this product can be administered without regard for timing of meals
 ■ adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours
 ■ children under 12 years of age: do not use

Other information
 ■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
 ■ store at 20-25°C (68-77°F)
 ■ see bottom of bottle for lot code and expiration date

Drug Facts (continued)

Inactive ingredients carborer 934P, NF; FD&C yellow #6 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF


Please visit our Web site:
www.mucinex.com

Contains FD&C Yellow Dye #6

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155
© 2003 Adams Laboratories, Inc.
US Pat. 6,372,252 B1

102003
2005720

NEW NDC 63824-057-20
Mucinex^D
 600 mg guaifenesin and
 60 mg pseudoephedrine HCl per tablet
 EXPECTORANT AND
 NASAL DECONGESTANT



20 BI-LAYER TABLETS

3-63824-057-20-0

1 3/4"

Drug Facts

Active ingredients (in each extended-release tablet)

Guaifenesin 600 mg	Expectorant
Pseudoephedrine HCl 60 mg	Nasal Decongestant

Uses
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
 ■ temporarily relieves nasal congestion due to:
 ■ common cold ■ hay fever
 ■ upper respiratory allergies
 ■ temporarily restores freer breathing through the nose
 ■ promotes nasal and/or sinus drainage
 ■ temporarily relieves sinus congestion and pressure

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
 ■ heart disease ■ high blood pressure
 ■ thyroid disease ■ diabetes
 ■ trouble urinating due to an enlarged prostate gland
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
 ■ cough accompanied by too much phlegm (mucus)

MUCINEX D 600/60 mg (20 count) CARTON

Title:
14 pt. Helvetica Bold
Italic, left justified

Headings:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Subheadings:
6 pt. Helvetica Bold
left justified

Graphic leading
to next panel

Right justified

Title for continued panel:
8 pt. Helvetica Bold Italic

8 pt. Helvetica Regular

-2.5 pt. barline

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

-0.5 pt. hairline

-2.5 pt. box barline

Drug Facts

Active ingredients (in each extended-release tablet)

Guafenesin 600 mg	Expectorant
Pseudoephedrine HCl 60 mg	Nasal Decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves nasal congestion due to:
 - common cold
 - hay fever
 - upper respiratory allergies
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- temporarily relieves sinus congestion and pressure

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Drug Facts (continued)

When using this product

- do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions


- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
- store at 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

Drug Facts (continued)

Inactive ingredients carbomer 934P, NF; FD&C yellow #6 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF



40 BI-LAYER TABLETS

Mucinex^D
 600 mg guaifenesin and
 60 mg pseudoephedrine HCl per tablet
 EXPECTORANT AND
 NASAL DECONGESTANT

NEW

Drug Facts (continued)

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
- store at 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

Drug Facts (continued)

Inactive ingredients carboxer 934P, NF; FD&C yellow #6 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

Please visit our Web site:
www.mucinex.com

Contains FD&C Yellow Dye #6




Fort Worth, Texas 76155
 © 2003 Adams Laboratories, Inc.
 US Pat. 6,372,252 B1

102003
 2005740

NEW

NDC 63824-057-40

Mucinex^D
 600 mg guaifenesin and
 60 mg pseudoephedrine HCl per tablet
 EXPECTORANT AND
 NASAL DECONGESTANT



40 BI-LAYER TABLETS

Drug Facts

Active ingredients (in each extended-release tablet)

Guaifenesin 600 mg Expectorant
 Pseudoephedrine HCl 60 mg Nasal Decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves nasal congestion due to:
 - common cold
 - upper respiratory allergies
 - hay fever
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- temporarily relieves sinus congestion and pressure

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

3-63824-057-40-8

MUCINEX D 600/60 mg (40 count) CARTON

Title:
14 pt. Helvetica Bold
Italic, left justified

Headings:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Subheadings:
6 pt. Helvetica Bold
left justified

Graphic leading
to next panel

Right justified

Title for continued panel:
8 pt. Helvetica Bold Italic

8 pt. Helvetica Regular

-2.5 pt. barline

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

-0.5 pt. hairline

-2.5 pt. box barline

Drug Facts

Active ingredients (in each extended-release tablet)

Guafenesin 600 mg	Expectorant
Pseudoephedrine HCl 60 mg	Nasal Decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves nasal congestion due to:
 - common cold
 - hay fever
 - upper respiratory allergies
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- temporarily relieves sinus congestion and pressure

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Drug Facts (continued)

When using this product

- do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
- store at 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

Drug Facts (continued)

Inactive ingredients carbomer 934P, NF; FD&C yellow #6 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

MUCINEX D 1200/120 mg SAMPLE TRAY

Title:
14 pt. Helvetica Bold
Italic, left justified

Headings:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Subheadings:
6 pt. Helvetica Bold
left justified

Graphic leading
to next panel

Drug Facts		Drug Facts (continued)	
Active ingredients (in each extended-release tablet)		When using this product	
Guaifenesin 1200 mg	Expectorant	<ul style="list-style-type: none"> ■ do not use more than directed 	
Pseudoephedrine HCl 120 mg	Nasal Decongestant	Stop use and ask a doctor if <ul style="list-style-type: none"> ■ you get nervous, dizzy, or sleepless ■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness. 	
Uses		If pregnant or breast-feeding , ask a health professional before use.	
<ul style="list-style-type: none"> ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily relieves nasal congestion due to: <ul style="list-style-type: none"> ■ common cold ■ hay fever ■ upper respiratory allergies ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage ■ temporarily relieves sinus congestion and pressure 		Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Warnings		Directions	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.		<ul style="list-style-type: none"> ■ do not crush, chew, or break tablet ■ take with a full glass of water ■ this product can be administered without regard for timing of meals ■ adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours ■ children under 12 years of age: do not use 	
Ask a doctor before use if you have		Other information	
<ul style="list-style-type: none"> ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough accompanied by too much phlegm (mucus) 		<ul style="list-style-type: none"> ■ tamper evident: do not use if seal on bottle printed "SEALED FOR YOUR PROTECTION" is broken or missing ■ store at 20-25°C (68-77°F) ■ see bottom of bottle for lot code and expiration date 	
Inactive ingredients		Inactive ingredients	
carbomer 934P, NF; FD&C red #40 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF		carbomer 934P, NF; FD&C red #40 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF	

Title for continued panel:
8 pt. Helvetica Bold Italic


8 pt. Helvetica Regular

0.5 pt. hairline

2.5 pt. barline

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline



10 BI-LAYER TABLETS

NEW
Mucinex^D
 1200 mg guaifenesin and
 120 mg pseudoephedrine HCl per tablet
 EXPECTORANT AND
 NASAL DECONGESTANT

NEW

1 3/4"

Drug Facts (continued)

When using this product
 ■ do not use more than directed

Stop use and ask a doctor if
 ■ you get nervous, dizzy, or sleepless
 ■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.


Directions
 ■ do not crush, chew, or break tablet
 ■ take with a full glass of water
 ■ this product can be administered without regard for timing of meals
 ■ adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
 ■ children under 12 years of age: do not use

Other information
 ■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
 ■ store at 20-25°C (68-77°F)
 ■ see bottom of bottle for lot code and expiration date

Drug Facts (continued)

Inactive ingredients carbomer 934P, NF; FD&C red #40 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

NEW NDC 63824-034-01
Mucinex^D
 1200 mg guaifenesin and
 120 mg pseudoephedrine HCl per tablet
 EXPECTORANT AND
 NASAL DECONGESTANT



Please visit our Web site:
www.mucinex.com

Adams[®]
 LABORATORIES, INC.
 Fort Worth, Texas 76155
 © 2003 Adams Laboratories, Inc.
 US Pat. 6,372,252 B1

102003
 2003401

10 BI-LAYER TABLETS

Drug Facts

Active ingredients (in each extended-release tablet)
 Guaifenesin 1200 mg.....Expectorant
 Pseudoephedrine HCl 120 mg.....Nasal Decongestant

Uses
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
 ■ temporarily relieves nasal congestion due to:
 ■ common cold ■ hay fever
 ■ upper respiratory allergies
 ■ temporarily restores freer breathing through the nose
 ■ promotes nasal and/or sinus drainage
 ■ temporarily relieves sinus congestion and pressure

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
 ■ heart disease ■ high blood pressure
 ■ thyroid disease ■ diabetes
 ■ trouble urinating due to an enlarged prostate gland
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
 ■ cough accompanied by too much phlegm (mucus) ▶

3-63824-034-01-0

MUCINEX D 1200/120 mg (10 count) CARTON

Title:
14 pt. Helvetica Bold
Italic, left justified

Headings:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Subheadings:
6 pt. Helvetica Bold
left justified

Graphic leading
to next panel

Right justified

Title for continued panel:
8 pt. Helvetica Bold Italic

8 pt. Helvetica Regular

-2.5 pt. barline

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

-0.5 pt. hairline

-2.5 pt. box barline

Drug Facts

Active ingredients (in each extended-release tablet)

Guafenesin 1200 mg.....Expectorant
Pseudoephedrine HCl 120 mg.....Nasal Decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves nasal congestion due to:
 - common cold
 - hay fever
 - upper respiratory allergies
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- temporarily relieves sinus congestion and pressure

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Drug Facts (continued)

When using this product

- do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions


- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
- store at 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

Drug Facts (continued)

Inactive ingredients carbomer 934P, NF; FD&C red #40 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF



20 BI-LAYER TABLETS

Mucinex^D
 1200 mg guaifenesin and
 120 mg pseudoephedrine HCl per tablet
 EXPECTORANT AND
 NASAL DECONGESTANT

NEW

Drug Facts (continued)

When using this product
 ■ do not use more than directed

Stop use and ask a doctor if
 ■ you get nervous, dizzy, or sleepless
 ■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ do not crush, chew, or break tablet
 ■ take with a full glass of water
 ■ this product can be administered without regard for timing of meals
 ■ adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
 ■ children under 12 years of age: do not use

Other information
 ■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
 ■ store at 20-25°C (68-77°F)
 ■ see bottom of bottle for lot code and expiration date

Drug Facts (continued)


Inactive ingredients carbomer 934P, NF; FD&C red #40 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

Please visit our Web site:
www.mucinex.com

Adams[®]
 LABORATORIES, INC.
 Fort Worth, Texas 76155
 © 2003 Adams Laboratories, Inc.
 US Pat. 6,372,252 B1

102003
 2003420

NEW NDC 63824-034-20
Mucinex^D
 1200 mg guaifenesin and
 120 mg pseudoephedrine HCl per tablet
 EXPECTORANT AND
 NASAL DECONGESTANT



20 BI-LAYER TABLETS

3-63824-034-20-1

Drug Facts

Active ingredients (in each extended-release tablet)

Guaifenesin 1200 mg.....	Expectorant
Pseudoephedrine HCl 120 mg.....	Nasal Decongestant

Uses
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
 ■ temporarily relieves nasal congestion due to:
 ■ common cold ■ hay fever
 ■ upper respiratory allergies
 ■ temporarily restores freer breathing through the nose
 ■ promotes nasal and/or sinus drainage
 ■ temporarily relieves sinus congestion and pressure

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

■ heart disease	■ high blood pressure
■ thyroid disease	■ diabetes
■ trouble urinating due to an enlarged prostate gland	
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema	
■ cough accompanied by too much phlegm (mucus)	▶