

NDA 19-452/S-016/S-019 & S-020

Page 4

Derma-Smoothe/FS[®]
fluocinolone acetonide
Topical Oil, 0.01%


(Body Oil)

For Topical Use Only-
Not for Not for Oral, Ophthalmic, or Intravaginal Use

NDC 28105-150-04

DESCRIPTION

Derma-Smoothe/FS[®] Topical Oil contains fluocinolone acetonide {(6 α , 11 β , 16 α)-6,9-difluoro-11,21-dihydroxy-16,17[(1-methylethylidene)bis(oxy)]-pregna-1,4-diene-3,20-dione, cyclic 16,17 acetal with acetone}, a synthetic corticosteroid for topical dermatologic use. This formulation is also marketed as Derma-Smoothe/FS[®] 0.01% fluocinolone acetonide for use as scalp oil with “shower caps” for scalp psoriasis in adults and as fluocinolone acetonide oil, 0.01% for chronic eczematous external otitis. Chemically, fluocinolone acetonide is C₂₄ H₃₀ F₂ O₆. It has the following structural formula:



Fluocinolone acetonide in Derma-Smoothe/FS[®] has a molecular weight of 452.50. It is a white crystalline powder that is odorless, stable in light, and melts at 270°C with decomposition; soluble in alcohol, acetone and methanol; slightly soluble in chloroform; insoluble in water.

Each gram of Derma-Smoothe/FS[®] contains approximately 0.11 mg of fluocinolone acetonide in a blend of oils, which contains isopropyl alcohol, isopropyl myristate, light mineral oil, oleth-2, refined peanut oil NF and fragrances.

CLINICAL PHARMACOLOGY

Like other topical corticosteroids, fluocinolone acetonide has anti-inflammatory, antipruritic, and vasoconstrictive properties. The mechanism of the anti-inflammatory activity of the topical steroids, in general, is unclear. However, corticosteroids are thought to act by the induction of phospholipase A₂ inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A₂.

Pharmacokinetics: The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the vehicle and the integrity of the epidermal barrier. Occlusion of topical corticosteroids can enhance penetration. Topical corticosteroids can be absorbed from normal intact skin. Also, inflammation and/or other disease processes in the skin can increase percutaneous absorption.

NDA 19-452/S-016/S-019 & S-020

Page 5

Derma-Smoothe/FS[®] is in the low to medium range of potency as compared with other topical corticosteroids.

CLINICAL STUDIES

In a vehicle-controlled study for the treatment of psoriasis of the scalp in adults, after 21 days of treatment, 60% of patients on active treatment and 21% of patients on the drug vehicle had excellent to cleared clinical response.

Open-label safety studies on 33 children (20 subjects ages 2 to 6 years, 13 subjects ages 7 to 12 years) with moderate to severe stable atopic dermatitis, and baseline body surface area involvement greater than 75% in 18 patients, and 50% to 75% in 15 patients, were treated with Derma-Smoothe/FS[®] twice daily for 4 weeks. Morning pre-stimulation cortisol level and post-Cortrosyn stimulation cortisol level were obtained in each subject at the beginning of the trial and at the end of 4 weeks of treatment. At the end of treatment, 4 out of 18 subjects aged 2 to 5 years showed low pre-stimulation cortisol levels (3.2 to 6.6 µg/dL; normal: cortisol > 7µg/dL) but all had normal responses to 0.25 mg of Cortrosyn stimulation (cortisol > 18 µg/dL).

A clinical study was conducted to assess the safety of Derma-Smoothe/FS, which contains refined peanut oil, on subjects with known peanut allergies. The study enrolled 13 patients with atopic dermatitis, 6 to 17 years of age. Of the 13 patients, 9 were Radioallergosorbent Test (RAST) positive to peanuts and 4 had no peanut sensitivity (controls). The study evaluated the responses to both prick test and patch test utilizing peanut oil NF, Derma-Smoothe/FS[®] and histamine/saline controls, on the 13 individuals. These subjects were also treated with Derma-Smoothe/FS[®] twice daily for 7 days. Prick test and patch test results for all 13 patients were negative to Derma-Smoothe/FS[®] and the refined peanut oil. One of the 9 peanut-sensitive patients experienced an exacerbation of atopic dermatitis after 5 days of Derma-Smoothe/FS[®]. Importantly, the bulk peanut oil NF, used in Derma-Smoothe/FS[®] is heated at 475° F for at least 15 minutes, which should provide for adequate decomposition of allergenic proteins.

INDICATION AND USAGE

Derma-Smoothe/FS[®] is a low to medium potency corticosteroid indicated:

In adult patients for the treatment of atopic dermatitis. (Body Oil)

In pediatric patients 2 years and older with moderate to severe atopic dermatitis (Body Oil). It may be used for up to 4 weeks.

CONTRAINDICATIONS

Derma-Smoothe/FS[®] is contraindicated in those patients with a history of hypersensitivity to any of the components of the preparation.

This product contains refined peanut oil NF (see PRECAUTIONS section).

PRECAUTIONS

General: Systemic absorption of topical corticosteroids can produce reversible hypothalamic-pituitary-adrenal (HPA) axis suppression with the potential for glucocorticosteroid insufficiency after

NDA 19-452/S-016/S-019 & S-020

Page 6

withdrawal of treatment. Manifestations of Cushing's syndrome, hyperglycemia, and glucosuria can also be produced in some patients by systemic absorption of topical corticosteroids while on treatment.

Patients applying a topical steroid to a large surface area or to areas under occlusion should be evaluated periodically for evidence of HPA axis suppression. This may be done by using the ACTH stimulation, A.M. plasma cortisol, and urinary free cortisol tests.

If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent corticosteroid. Infrequently, signs and symptoms of glucocorticoid insufficiency may occur requiring supplemental systemic corticosteroids. For information on systemic supplementation, see prescribing information for those products.

Children may be more susceptible to systemic toxicity from equivalent doses due to their larger skin surface to body mass ratios. (See PRECAUTIONS-Pediatric use)

Allergic contact dermatitis to any component of topical corticosteroids is usually diagnosed by a *failure to heal* rather than noting a clinical exacerbation, which may occur with most topical products not containing corticosteroids. Such an observation should be corroborated with appropriate diagnostic testing. One peanut sensitive child experienced a flare of his atopic dermatitis after 5 days of twice daily treatment with Derma-Smoothe/FS[®] (see CLINICAL STUDIES section).

If wheal and flare type reactions (which may be limited to pruritus) or other manifestations of hypersensitivity develop, Derma-Smoothe/FS[®] should be discontinued immediately and appropriate therapy instituted.

If concomitant skin infections are present or develop, an appropriate antifungal or antibacterial agent should be used. If a favorable response does not occur promptly, use of Derma-Smoothe/FS[®] should be discontinued until the infection has been adequately controlled.

Derma-Smoothe/FS[®] is formulated with 48% refined peanut oil NF. Peanut oil used in this product is routinely tested for peanut proteins using a sandwich enzyme-linked immunosorbent assay test (S-ELISA) kit, which can detect peanut proteins to as low as 2.5 parts per million (ppm).

Physicians should use caution in prescribing Derma-Smoothe/FS[®] for peanut-sensitive individuals.

Information for Patients: Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes. In case of contact, wash eyes liberally with water.
2. This medication should not be used for any disorder other than that for which it was prescribed.
3. Patients should promptly report to their physician any worsening of their skin condition.
4. Parents of pediatric patients should be advised not to use Derma-Smoothe/FS[®] in the treatment of diaper dermatitis. Derma-Smoothe/FS[®] should not be applied to the diaper area as diapers or plastic pants may constitute occlusive dressing.

NDA 19-452/S-016/S-019 & S-020

Page 7

5. This medication should not be used on the face, underarm, or groin unless directed by the physician.
6. As with other corticosteroids, therapy should be discontinued when control is achieved. If no improvement is seen within 2 weeks, contact the physician.

Laboratory Tests: The following tests may be helpful in evaluating patients for HPA axis suppression:

ACTH stimulation test
A.M. plasma cortisol test
Urinary free cortisol test

Carcinogenesis, mutagenesis, and impairment of fertility: Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of Derma-Smoothe/FS[®]. Studies have not been performed to evaluate the mutagenic potential of fluocinolone acetonide, the active ingredient in Derma-Smoothe/FS[®]. Some corticosteroids have been found to be genotoxic in various genotoxicity tests (i.e. the *in vitro* human peripheral blood lymphocyte chromosome aberration assay with metabolic activation, the *in vivo* mouse bone marrow micronucleus assay, the Chinese hamster micronucleus test and the *in vitro* mouse lymphoma gene mutation assay).

Pregnancy: Teratogenic effects: Pregnancy category C: Corticosteroids have been shown to be teratogenic in laboratory animals when administered systemically at relatively low dosage levels. Some corticosteroids have been shown to be teratogenic after dermal application in laboratory animals.

There are no adequate and well-controlled studies in pregnant women on teratogenic effects from Derma-Smoothe/FS[®]. Therefore, Derma-Smoothe/FS[®] should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers: Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk. Because many drugs are excreted in human milk, caution should be exercised when Derma-Smoothe/FS[®] is administered to a nursing woman.

Pediatric Use: Derma-Smoothe/FS[®] may be used twice daily for up to 4 weeks in pediatric patients 2 years and older with moderate to severe atopic dermatitis. Derma-Smoothe/FS[®] should not be applied to the diaper area.

Application to intertriginous areas should be avoided due to the increased possibility of local adverse events such as striae, atrophy, and telangiectasia, which may be irreversible. The smallest amount of drug needed to cover the affected areas should be applied. Long term safety in the pediatric population has not been established.

Derma-Smoothe/FS[®] is not recommended for use on the face (see ADVERSE REACTIONS section).

Because of a higher ratio of skin surface area to body mass, children are at a greater risk than adults of HPA-axis-suppression when they are treated with topical corticosteroids. They are therefore also at greater risk of glucocorticosteroid insufficiency after withdrawal of treatment and of Cushing's

NDA 19-452/S-016/S-019 & S-020

Page 8

syndrome while on treatment. Adverse effects including striae have been reported with inappropriate use of topical corticosteroids in infants and children. (SEE PRECAUTIONS).

HPA axis suppression, Cushing's syndrome, and intracranial hypertension have been reported in children receiving topical corticosteroids. Children may be more susceptible to systemic toxicity from equivalent doses due to their larger skin surface to body mass ratios. Manifestations of adrenal suppression in children include linear growth retardation, delayed weight gain, low plasma cortisol levels, and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Derma-Smoothe/FS[®] is formulated with 48% refined peanut oil NF, in which peanut protein is not detectable at 2.5 ppm. Physicians should use caution in prescribing Derma-Smoothe/FS[®] for peanut sensitive individuals.

ADVERSE REACTIONS

The following local adverse reactions have been reported infrequently with topical corticosteroids. They may occur more frequently with the use of occlusive dressings, especially with higher potency corticosteroids. These reactions are listed in an approximate decreasing order of occurrence: burning, itching, irritation, dryness, folliculitis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, secondary infection, skin atrophy, striae, and miliaria. One peanut sensitive child experienced a flare of his atopic dermatitis after 5 days of twice daily treatment with Derma-Smoothe/FS[®].

NDA 19-452/S-016/S-019 & S-020

Page 9

A post marketing (open-label) safety study was conducted in 58 children to evaluate the local safety of Derma-Smoothe/FS when applied twice daily for 4 weeks to the face in children (2 to 12 years) with moderate to severe atopic dermatitis (see table of Incidence of Adverse Events).

Incidence of Adverse Events (%)

N=58

Adverse Event (AE)*	# of patients (%)	Day 14	Day 28**	Day 56***
Any AE	15 (25.9)	6 (10.3)	7 (12.1)	7 (12.1)
Telangiectasia	5 (8.6)	3 (5.2)	4 (6.9)	2 (3.5)
Erythema	3 (5.2)			3 (5.2)
Itching	3 (5.2)			3 (5.2)
Irritation	3 (5.2)			3 (5.2)
Burning	3 (5.2)			3 (5.2)
Hypopigmentation	2 (3.5)	2 (3.5)		
Shiny skin	1 (1.7)		1 (1.7)	
Secondary atopic dermatitis	1 (1.7)			1 (1.7)
Papules and pustules	1 (1.7)			1 (1.7)
Keratosis pilaris	1 (1.7)			1 (1.7)
Folliculitis	1 (1.7)		1 (1.7)	
Facial herpes simplex	1 (1.7)	1 (1.7)		
Acneiform eruption	1 (1.7)		1 (1.7)	
Ear infection	1 (1.7)		1 (1.7)	

*The number of individual adverse events reported does not necessarily reflect the number of individual subjects, since one subject could have multiple reportings of an adverse event.

**End of Treatment

***Four Weeks Post Treatment

OVERDOSAGE

Topically applied Derma-Smoothe/FS[®] can be absorbed in sufficient amounts to produce systemic effects (see PRECAUTIONS).

DOSAGE AND ADMINISTRATION

Derma-Smoothe/FS[®] for atopic dermatitis in adults (Body Oil):

For the treatment of atopic dermatitis, Derma-Smoothe/FS[®] should be applied as a thin film to the affected area three times daily.

Derma-Smoothe/FS[®] for atopic dermatitis in pediatric patients 2 years and older (Body Oil):

Moisten skin. Apply Derma-Smoothe/FS[®] as a thin film to the affected areas twice daily for no longer than four weeks.

NDA 19-452/S-016/S-019 & S-020
Page 10

HOW SUPPLIED

Derma-Smoothe/FS[®] is supplied in bottles containing 4 fluid ounces. It is labeled as Body Oil (**NDC # 28105-150-04**).

Keep tightly closed. Store at 25° C (68° to 77° F); excursions permitted to 15°-30° C (59°-86°F.)
[see USP Controlled Room Temperature]

CAUTION: Rx only

MANUFACTURED AND DISTRIBUTED BY:

Hill Laboratories, Inc.
Sanford, Florida 32773

Rev. CODE xxxxx
Date: 1/04

NDA 19-452/S-016/S-019 & S-020
Page 11

CONTAINER LABEL

Left side of Container Label

NDC 28105-150-04

R_x Only

DERMA-SMOOTHIE/FS[®]

Fluocinolone Acetonide

0.01% Topical Oil

BODY OIL

FOR TOPICAL USE ONLY

NOT FOR ORAL, OPHTHALMIC or INTRAVAGINAL USE

SHAKE WELL BEFORE USE

Net Contents 118.28 mL
(4 fl. oz.)

SINCE 1943

**HILL DERMACEUTICALS, INC.
SANFORD, FLORIDA 32773**

NDA 19-452/S-016/S-019 & S-020
Page 12

Right Side of Container Label

Keep Out the Reach of Children
FOR TOPICAL USE ONLY
NOT FOR ORAL, OPHTHALMIC, or INTRAVAGINAL USE

DOSAGE AND ADMINISTRATION: Atopic dermatitis in pediatric patients 2 years and older:
Moisten skin. Apply Derma-Smoothe/FS[®] as a thin film to the affected areas twice daily for up to 4 weeks.

Atopic eczema / dermatitis in adults: Moisten skin. Apply Derma-Smoothe/FS[®] as a thin film to the affected areas three times daily.

DO NOT USE WITH OCCLUSION

SEE PACKAGE INSERT FOR FULL PRESCRIBING INFORMATION

Contains: Fluocinolone Acetonide (0.01%), Isopropyl Alcohol (1.6%), in a base containing Refined Peanut Oil NF, Mineral Oil, Isopropyl Myristate, Oleth-2, Cream Fragrance and Balsam Pine.

Storage: Keep tightly closed. Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F)
[see USP Controlled Room Temperature]

Lot No.
Exp. Date

Manufactured and Distributed by:
HILL DERMACEUTICALS, INC.
Sanford, Florida 32773

Manufactured by:
HILL LABORATORIES, INC.
Sanford, Florida 32773

CODE No. XXXXXXXX
Rev. XX/XX

NDA 19-452/S-016/S-019 & S-020

Page 13

CARTON LABEL

Front and Back of Box

NDC 28105-150-04

R_x only

DERMA-SMOOTHIE/FS[®]

Fluocinolone Acetonide
0.01% Topical Oil

BODY OIL

FOR TOPICAL USE ONLY
NOT FOR ORAL, OPHTHALMIC, or INTRAVAGINAL USE

SHAKE WELL BEFORE USE

Net Contents 118.28 mL
(4 fl. oz.)

SINCE 1949

HILL DERMACEUTICALS, INC.
SANFORD, FLORIDA 32773

NDA 19-452/S-016/S-019 & S-020
Page 14

SIDE PANEL 1

Contains: Fluocinolone Acetonide (0.01%), Isopropyl Alcohol (1.6%), in a base containing Refined Peanut Oil NF, Mineral Oil, Isopropyl Myristate, Oleth-2, Cream Fragrance and Balsam Pine.

Storage: Keep tightly closed. Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F)
[see USP Controlled Room Temperature]

Keep Out of Reach of Children
FOR TOPICAL USE ONLY
NOT FOR ORAL, OPHTHALMIC, or INTRAVAGINAL USE

CODE NO. XXXXXX

NDA 19-452/S-016/S-019 & S-020
Page 15

SIDE PANEL 2

DOSAGE AND ADMINISTRATION:

Atopic dermatitis in pediatric patients 2 years and older: Moisten skin. Apply Derma-Smoothe/FS[®] as a thin film to the affected areas twice daily for up to 4 weeks.

Atopic eczema / dermatitis in adults: Moisten skin. Apply Derma-Smoothe/FS[®] as a thin film to the affected areas three times daily.

DO NOT USE WITH OCCLUSION.

SEE PACKAGE INSERT FOR FULL PRESCRIBING INFORMATION.

Manufactured and Distributed by:
HILL DERMACEUTICALS, INC.
Sanford, Florida 32773

Manufactured by:
HILL LABORATORIES, INC.
Sanford, Florida 32773

NDA 19-452/S-016/S-019 & S-020
Page 16

Bottom Panel

Lot No:

Exp. Date:

Rev.

NDA 19-452/S-016/S-019 & S-020
Page 17

Derma-Smoothe/FS[®]
fluocinolone acetonide
Topical Oil, 0.01%


(Scalp Oil)

For Topical Use Only-
Not for Oral, Ophthalmic, or Intravaginal Use

NDC 28105-149-04

DESCRIPTION

Derma-Smoothe/FS[®] Topical Oil contains fluocinolone acetonide {(6 α , 11 β , 16 α)-6,9-difluoro-11,21-dihydroxy-16,17[(1-methylethylidene)bis(oxy)]-pregna-1,4-diene-3,20-dione, cyclic 16,17 acetal with acetone}, a synthetic corticosteroid for topical dermatologic use. This formulation is also marketed as Derma-Smoothe/FS[®] 0.01%, fluocinolone acetonide for use as body oil for atopic dermatitis in adults and for moderate to severe atopic dermatitis in pediatric patients 2 years and older and as fluocinolone acetonide oil, 0.01% for chronic eczematous external otitis. Chemically, fluocinolone acetonide is C₂₄H₃₀F₂O₆. It has the following structural formula:



Fluocinolone acetonide in Derma-Smoothe/FS[®] has a molecular weight of 452.50. It is a white crystalline powder that is odorless, stable in light, and melts at 270°C with decomposition; soluble in alcohol, acetone and methanol; slightly soluble in chloroform; insoluble in water.

Each gram of Derma-Smoothe/FS[®] contains approximately 0.11 mg of fluocinolone acetonide in a blend of oils, which contains isopropyl alcohol, isopropyl myristate, light mineral oil, oleth-2, refined peanut oil NF and fragrances.

Each packaged product contains 2 shower caps. The shower cap is made of low density polyethylene material with rubber elastic.

CLINICAL PHARMACOLOGY

Like other topical corticosteroids, fluocinolone acetonide has anti-inflammatory, antipruritic, and vasoconstrictive properties. The mechanism of the anti-inflammatory activity of the topical steroids, in general, is unclear. However, corticosteroids are thought to act by the induction of phospholipase A₂ inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A₂.

NDA 19-452/S-016/S-019 & S-020

Page 18

Pharmacokinetics: The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the vehicle and the integrity of the epidermal barrier. Occlusion of topical corticosteroids can enhance penetration. Topical corticosteroids can be absorbed from normal intact skin. Also, inflammation and/or other disease processes in the skin can increase percutaneous absorption.

Derma-Smoothe/FS[®] is in the low to medium range of potency as compared with other topical corticosteroids.

CLINICAL STUDIES

In a vehicle-controlled study for the treatment of psoriasis of the scalp in adults, after 21 days of treatment, 60% of patients on active treatment and 21% of patients on the drug vehicle had excellent to cleared clinical response.

Open-label safety studies on 33 children (20 subjects ages 2 to 6 years, 13 subjects ages 7 to 12 years) with moderate to severe stable atopic dermatitis, and baseline body surface area involvement greater than 75% in 18 patients, and 50% to 75% in 15 patients, were treated with Derma-Smoothe/FS[®] twice daily for 4 weeks. Morning pre-stimulation cortisol level and post-Cortrosyn stimulation cortisol level were obtained in each subject at the beginning of the trial and at the end of 4 weeks of treatment. At the end of treatment, 4 out of 18 subjects aged 2 to 5 years showed low pre-stimulation cortisol levels (3.2 to 6.6 µg/dL; normal: cortisol > 7µg/dL) but all had normal responses to 0.25 mg of Cortrosyn stimulation (cortisol > 18 µg/dL).

A clinical study was conducted to assess the safety of Derma-Smoothe/FS, which contains refined peanut oil, on subjects with known peanut allergies. The study enrolled 13 patients with atopic dermatitis, 6 to 17 years of age. Of the 13 patients, 9 were Radioallergosorbent Test (RAST) positive to peanuts and 4 had no peanut sensitivity (controls). The study evaluated the responses to both prick test and patch test utilizing peanut oil NF, Derma-Smoothe/FS[®] and histamine/saline controls, on the 13 individuals. These subjects were also treated with Derma-Smoothe/FS[®] twice daily for 7 days. Prick test and patch test results for all 13 patients were negative to Derma-Smoothe/FS[®] and the refined peanut oil. One of the 9 peanut-sensitive patients experienced an exacerbation of atopic dermatitis after 5 days of Derma-Smoothe/FS[®]. Importantly, the bulk peanut oil NF, used in Derma-Smoothe/FS[®] is heated at 475° F for at least 15 minutes, which should provide for adequate decomposition of allergenic proteins.

INDICATION AND USAGE

Derma-Smoothe/FS[®] is a low to medium potency corticosteroid indicated:

In adult patients for the treatment of psoriasis of the scalp (Scalp Oil).

~~In pediatric patients 2 years and older with moderate to severe atopic dermatitis. It may be used for up to 4 weeks.~~

NDA 19-452/S-016/S-019 & S-020

Page 19

CONTRAINDICATIONS

Derma-Smoothe/FS[®] is contraindicated in those patients with a history of hypersensitivity to any of the components of the preparation.

This product contains refined peanut oil NF (see PRECAUTIONS section).

PRECAUTIONS

General: Systemic absorption of topical corticosteroids can produce reversible hypothalamic-pituitary-adrenal (HPA) axis suppression with the potential for glucocorticosteroid insufficiency after withdrawal of treatment. Manifestations of Cushing's syndrome, hyperglycemia, and glucosuria can also be produced in some patients by systemic absorption of topical corticosteroids while on treatment.

Patients applying a topical steroid to a large surface area or to areas under occlusion should be evaluated periodically for evidence of HPA axis suppression. This may be done by using the ACTH stimulation, A.M. plasma cortisol, and urinary free cortisol tests.

If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent corticosteroid. Infrequently, signs and symptoms of glucocorticoid insufficiency may occur requiring supplemental systemic corticosteroids. For information on systemic supplementation, see prescribing information for those products.

Children may be more susceptible to systemic toxicity from equivalent doses due to their larger skin surface to body mass ratios.

Allergic contact dermatitis to any component of topical corticosteroids is usually diagnosed by a *failure to heal* rather than noting a clinical exacerbation, which may occur with most topical products not containing corticosteroids. Such an observation should be corroborated with appropriate diagnostic testing. One peanut-sensitive child experienced a flare of his atopic dermatitis after 5 days of twice daily treatment with Derma-Smoothe/FS[®] (see CLINICAL STUDIES section).

If wheal and flare type reactions (which may be limited to pruritus) or other manifestations of hypersensitivity develop, Derma-Smoothe/FS[®] should be discontinued immediately and appropriate therapy instituted.

If concomitant skin infections are present or develop, an appropriate antifungal or antibacterial agent should be used. If a favorable response does not occur promptly, use of Derma-Smoothe/FS[®] should be discontinued until the infection has been adequately controlled.

Derma-Smoothe/FS[®] is formulated with 48% refined peanut oil NF. Peanut oil used in this product is routinely tested for peanut proteins using a sandwich enzyme-linked immunosorbent assay test (S-ELISA) kit, which can detect peanut proteins to as low as 2.5 parts per million (ppm).

Physicians should use caution in prescribing Derma-Smoothe/FS[®] for peanut-sensitive individuals.

NDA 19-452/S-016/S-019 & S-020
Page 20

Information for Patients: Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes. In case of contact, wash eyes liberally with water.
2. This medication should not be used for any disorder other than that for which it was prescribed.
3. Patients should promptly report to their physician any worsening of their skin condition.
4. Parents of pediatric patients should be advised not to use Derma-Smoothe/FS[®] in the treatment of diaper dermatitis. Derma-Smoothe/FS[®] should not be applied to the diaper area as diapers or plastic pants may constitute occlusive dressing.
5. This medication should not be used on the face, underarm, or groin unless directed by the physician.
6. As with other corticosteroids, therapy should be discontinued when control is achieved. If no improvement is seen within 2 weeks, contact the physician.

Laboratory Tests: The following tests may be helpful in evaluating patients for HPA axis suppression:

ACTH stimulation test
A.M. plasma cortisol test
Urinary free cortisol test

Carcinogenesis, mutagenesis, and impairment of fertility: Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of Derma-Smoothe/FS[®]. Studies have not been performed to evaluate the mutagenic potential of fluocinolone acetonide, the active ingredient in Derma-Smoothe/FS[®]. Some corticosteroids have been found to be genotoxic in various genotoxicity tests (i.e. the *in vitro* human peripheral blood lymphocyte chromosome aberration assay with metabolic activation, the *in vivo* mouse bone marrow micronucleus assay, the Chinese hamster micronucleus test and the *in vitro* mouse lymphoma gene mutation assay).

Pregnancy: Teratogenic effects: Pregnancy category C: Corticosteroids have been shown to be teratogenic in laboratory animals when administered systemically at relatively low dosage levels. Some corticosteroids have been shown to be teratogenic after dermal application in laboratory animals.

There are no adequate and well-controlled studies in pregnant women on teratogenic effects from Derma-Smoothe/FS[®]. Therefore, Derma-Smoothe/FS[®] should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers: Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk. Because many drugs are excreted in human milk, caution should be exercised when Derma-Smoothe/FS[®] is administered to a nursing woman.

NDA 19-452/S-016/S-019 & S-020

Page 21

Pediatric Use: Derma-Smoothe/FS[®] may be used twice daily for up to 4 weeks in pediatric patients 2 years and older with moderate to severe atopic dermatitis. Derma-Smoothe/FS[®] should not be applied to the diaper area.

Application to intertriginous areas should be avoided due to the increased possibility of local adverse events such as striae, atrophy, and telangiectasia, which may be irreversible. The smallest amount of drug needed to cover the affected areas should be applied. Long term safety in the pediatric population has not been established.

Derma-Smoothe/FS[®] is not recommended for use on the face (see ADVERSE REACTIONS section).

Because of a higher ratio of skin surface area to body mass, children are at a greater risk than adults of HPA-axis-suppression when they are treated with topical corticosteroids. They are therefore also at greater risk of glucocorticosteroid insufficiency after withdrawal of treatment and of Cushing's syndrome while on treatment. Adverse effects including striae have been reported with inappropriate use of topical corticosteroids in infants and children. (SEE PRECAUTIONS).

HPA axis suppression, Cushing's syndrome, and intracranial hypertension have been reported in children receiving topical corticosteroids. Children may be more susceptible to systemic toxicity from equivalent doses due to their larger skin surface to body mass ratios. Manifestations of adrenal suppression in children include linear growth retardation, delayed weight gain, low plasma cortisol levels, and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Derma-Smoothe/FS[®] is formulated with 48% refined peanut oil NF, in which peanut protein is not detectable at 2.5 ppm. Physicians should use caution in prescribing Derma-Smoothe/FS[®] for peanut sensitive individuals.

ADVERSE REACTIONS

The following local adverse reactions have been reported infrequently with topical corticosteroids. They may occur more frequently with the use of occlusive dressings, especially with higher potency corticosteroids. These reactions are listed in an approximate decreasing order of occurrence: burning, itching, irritation, dryness, folliculitis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, secondary infection, skin atrophy, striae, and miliaria. One peanut sensitive child experienced a flare of his atopic dermatitis after 5 days of twice daily treatment with Derma-Smoothe/FS[®].

NDA 19-452/S-016/S-019 & S-020
Page 22

A post marketing (open-label) safety study was conducted in 58 children to evaluate the local safety of Derma-Smoothe/FS when applied twice daily for 4 weeks to the face in children (2 to 12 years) with moderate to severe atopic dermatitis (see table of Incidence of Adverse Events).

Incidence of Adverse Events (%)
N=58

Adverse Event (AE)*	# of patients (%)	Day 14	Day 28**	Day 56***
Any AE	15 (25.9)	6 (10.3)	7 (12.1)	7 (12.1)
Telangiectasia	5 (8.6)	3 (5.2)	4 (6.9)	2 (3.5)
Erythema	3 (5.2)			3 (5.2)
Itching	3 (5.2)			3 (5.2)
Irritation	3 (5.2)			3 (5.2)
Burning	3 (5.2)			3 (5.2)
Hypopigmentation	2 (3.5)	2 (3.5)		
Shiny skin	1 (1.7)		1 (1.7)	
Secondary atopic dermatitis	1 (1.7)			1 (1.7)
Papules and pustules	1 (1.7)			1 (1.7)
Keratosis pilaris	1 (1.7)			1 (1.7)
Folliculitis	1 (1.7)		1 (1.7)	
Facial herpes simplex	1 (1.7)	1 (1.7)		
Acneiform eruption	1 (1.7)		1 (1.7)	
Ear infection	1 (1.7)		1 (1.7)	

*The number of individual adverse events reported does not necessarily reflect the number of individual subjects, since one subject could have multiple reportings of an adverse event.

**End of Treatment

***Four Weeks Post Treatment

OVERDOSAGE

Topically applied Derma-Smoothe/FS[®] can be absorbed in sufficient amounts to produce systemic effects (see PRECAUTIONS).

DOSAGE AND ADMINISTRATION

Derma-Smoothe/FS[®] for scalp psoriasis in adults (Scalp Oil):

For the treatment of scalp psoriasis, wet or dampen hair and scalp thoroughly. Apply a thin film of Derma-Smoothe/FS[®] on the scalp, massage well and cover scalp with the supplied shower cap. Leave on overnight or for a minimum of 4 hours before washing off. Wash hair with regular shampoo and rinse thoroughly.

NDA 19-452/S-016/S-019 & S-020
Page 23

HOW SUPPLIED

Derma-Smoothe/FS[®] is supplied in bottles containing 4 fluid ounces. It is labeled as Scalp Oil (**NDC # 28105-149-04**).

Scalp Oil is supplied with 2 shower caps.

Keep tightly closed. Store at 25° C (68° to 77° F); excursions permitted to 15°-30° C (59°-86°F.)
[see USP Controlled Room Temperature]

CAUTION: Rx only

MANUFACTURED BY AND DISTRIBUTED BY:
Hill Laboratories, Inc.
Sanford, Florida 32773

Rev. CODE xxxxx
Date: 1/04

NDA 19-452/S-016/S-019 & S-020
Page 24

CARTON LABEL

~~DERMA-SMOOTHIE/FS SCALP OIL~~

Front and Back of Carton

NDC 28105-149-04

R_x only

DERMA-SMOOTHIE/FS[®]

Fluocinolone Acetonide
0.01% Topical Oil

(SCALP OIL)

FOR TOPICAL USE ONLY
NOT FOR ORAL, OPHTHALMIC, or INTRAVAGINAL USE
SHAKE WELL BEFORE USE

Net Wt. 4 fl. oz. (118.28 mL)

Since 1943

HILL
Dermaceuticals, Inc
Sanford, Florida 32773

NDA 19-452/S-016/S-019 & S-020
Page 25

Left Side of Carton

Contains:

Fluocinolone Acetonide (0.01%), Isopropyl Alcohol (1.6%), in a base containing **Refined** Peanut Oil NF, Mineral Oil, Isopropyl Myristate, Oleth-2, Cream Fragrance and Balsam Pine.

Contents of Package:

4 fl. oz. bottle
2 shower caps

Storage:

Keep tightly closed.

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature]

Keep Out of Reach of Children

FOR TOPICAL USE ONLY

NOT FOR ORAL, OPHTHALMIC, or INTRAVAGINAL USE

CODE NO XXXXXXXX

NDA 19-452/S-016/S-019 & S-020
Page 26

Right Side of Carton

DOSAGE AND ADMINISTRATION:

For the treatment of **scalp psoriasis**, wet or dampen hair and scalp thoroughly. Apply a thin film of Derma-Smoothe/FS[®] on the scalp, massage well and cover scalp with one of the supplied shower caps. Leave on overnight or for a minimum of 4 hours before washing off. Wash hair with regular shampoo and rinse thoroughly.

**SEE PACKAGE
INSERT FOR FULL
PRESCRIBING
INFORMATION.**

Manufactured and Distributed By:
HILL DERMACEUTICALS, INC.
Sanford, Florida 32773

Manufactured By:
HILL LABORATORIES, INC.
Sanford, Florida 32773

Product Bar Code Here

NDA 19-452/S-016/S-019 & S-020
Page 27

Bottom Flap of Carton

Lot No.

Exp. Date:

REV. XX/XX

NDA 19-452/S-016/S-019 & S-020
Page 28

CONTAINER LABEL

DERMA-SMOOTHIE/FS SCALP OIL

LEFT SIDE

NDC 28105-149-04

R_x only

DERMA-SMOOTHIE/FS[®]

Fluocinolone Acetonide
0.01% Topical Oil

(SCALP OIL)

FOR TOPICAL USE ONLY
NOT FOR ORAL, OPHTHALMIC, or INTRAVAGINAL USE
SHAKE WELL BEFORE USE

Net Wt. 4 fl. oz. (118.28 mL)

Since 1943

Hill
Dermaceuticals, Inc.
Sanford, Florida 32773

NDA 19-452/S-016/S-019 & S-020
Page 29

RIGHT SIDE

Keep Out of Reach of Children
FOR TOPICAL USE ONLY
Not For Oral, Ophthalmic, or Intravaginal Use

Dosage and Administration:

For the treatment of **scalp psoriasis**, wet or dampen hair and scalp thoroughly. Apply a thin film of Derma-Smoothe/FS[®] on the scalp, massage well and cover scalp with one of the supplied shower caps. Leave on overnight or for a minimum of 4 hours before washing off. Wash hair with regular shampoo and rinse thoroughly.
See package Insert for Full Prescribing Information.

Contains:

Fluocinolone Acetonide (0.01%), Isopropyl Alcohol (1.6%), in a base containing **Refined** Peanut Oil NF, Mineral Oil, Isopropyl Myristate, Oleth-2, Cream Fragrance and Balsam Pine.

Contents of Package: 4 fl. oz. bottle / 2 shower caps

Storage:

Keep tightly closed.

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature]

Lot No.:
Exp. Date:

Manufactured and Distributed By:
HILL DERMACEUTICALS, INC.
Sanford, Florida 32773

Manufactured By:
HILL LABORATORIES, INC.
Sanford, Florida 32773

Code: XXXXXXXX
Rev. XX/XX

NDA 19-452/S-016/S-019 & S-020
Page 30

2

NDC 28105-149-04

Rx only

Derma-Smoothie/FS®

Fluocinolone Acetonide
0.01% Topical Oil

(SCALP OIL)

- FOR TOPICAL USE ONLY
- NOT FOR OPHTHALMIC USE
- SHAKE WELL BEFORE USE

Net Wt. 4 fl. oz. (118.28 mL)

Since 1943

HILL
Dermaceuticals, Inc.
Sanford, Florida 32773

Contains:

Fluocinolone Acetonide (0.01%), Isopropyl Alcohol (1.6%), in a base containing Peanut Oil, Mineral Oil, Isopropyl Myristate, Oleth-2, Cream Fragrance and Balsam Pine.

Contents of Package:

4 fl. oz. bottle
2 shower caps

Storage:

Keep tightly closed.
Store at 25°C (77°F); excursions permitted to 15-30°C (59-86° F) [see USP Controlled Room Temperature]

Keep Out of Reach of
Children

FOR EXTERNAL
USE ONLY

NOT FOR
OPHTHALMIC USE

CODE NO 171A106

NDC 28105-149-04

Rx only

Derma-Smoothie/FS®

Fluocinolone Acetonide
0.01% Topical Oil

(SCALP OIL)

- FOR TOPICAL USE ONLY
- NOT FOR OPHTHALMIC USE
- SHAKE WELL BEFORE USE

Net Wt. 4 fl. oz. (118.28 mL)

Since 1943

HILL
Dermaceuticals, Inc.
Sanford, Florida 32773

**DOSAGE AND
ADMINISTRATION:**

For the treatment of **scalp psoriasis**, wet or dampen hair and scalp thoroughly. Apply a thin film of Derma-Smoothie/FS® on the scalp, massage well and cover scalp with one of the supplied shower caps. Leave on overnight or for a minimum of 4 hours before washing off. Wash hair with regular shampoo and rinse thoroughly.

SEE PACKAGE
INSERT FOR FULL
PRESCRIBING
INFORMATION.

Distributed By:
HILL DERMACEUTICALS,
INC.
Sanford, Florida 32773

Manufactured By:
HILL LABORATORIES, INC.
Sanford, Florida 32773



1	2	3
1	1	1
2	2	2
3	3	3

Lot No.

Exp. Date:

REV. 7/03

NDA 19-452/S-016/S-019 & S-020
Page 31

2

NDC 28105-149-04

Rx only

Derma-Smoothie/FS®

Fluocinolone Acetonide
0.01% Topical Oil

(SCALP OIL)

- FOR TOPICAL USE ONLY
- NOT FOR OPHTHALMIC USE
- SHAKE WELL BEFORE USE

Net Wt. 4 fl. oz. (118.28 mL)

Since 1943

HILL
Dermaceuticals, Inc.
Sanford, Florida 32773

Contains:

Fluocinolone Acetonide (0.01%), Isopropyl Alcohol (1.6%), in a base containing Peanut Oil, Mineral Oil, Isopropyl Myristate, Oleth-2, Cream Fragrance and Balsam Pine.

Contents of Package:

4 fl. oz. bottle
2 shower caps

Storage:

Keep tightly closed.
Store at 25°C (77°F);
excursions permitted to 15-30°C (59-86° F)
[see USP Controlled Room Temperature]

Keep Out of Reach of
Children

FOR EXTERNAL
USE ONLY

NOT FOR
OPHTHALMIC USE

CODE NO 171A106

NDC 28105-149-04

Rx only

Derma-Smoothie/FS®

Fluocinolone Acetonide
0.01% Topical Oil

(SCALP OIL)

- FOR TOPICAL USE ONLY
- NOT FOR OPHTHALMIC USE
- SHAKE WELL BEFORE USE

Net Wt. 4 fl. oz. (118.28 mL)

Since 1943

HILL
Dermaceuticals, Inc.
Sanford, Florida 32773

**DOSAGE AND
ADMINISTRATION:**

For the treatment of **scalp psoriasis**, wet or dampen hair and scalp thoroughly. Apply a thin film of Derma-Smoothie/FS® on the scalp, massage well and cover scalp with one of the supplied shower caps. Leave on overnight or for a minimum of 4 hours before washing off. Wash hair with regular shampoo and rinse thoroughly.

SEE PACKAGE
INSERT FOR FULL
PRESCRIBING
INFORMATION.

Distributed By:
HILL DERMACEUTICALS,
INC.
Sanford, Florida 32773

Manufactured By:
HILL LABORATORIES, INC.
Sanford, Florida 32773



1	2	3
1	1	1
2	2	2
3	3	3

Lot No.

Exp. Date:

REV. 7/03