

Drug Facts (continued)

Other information ● each caplet contains: sodium 20 mg ● store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-800-331-4536 or www.midol.com

Midol
F.P.O. MAXIMUM STRENGTH

Extended Relief
for up to 12 HOURS

NAPROXEN SODIUM TABLETS, 220 mg PAIN RELIEVER

24 CAPLETS* capsule-shaped tablets

Cramps • Backache • Headache

EXP: 4/07 UIC: 287624D

MAXIMUM STRENGTH
Midol® Extended Relief
Dist. by: Bayer HealthCare LLC

READ CARTON BEFORE USE. Each caplet contains: Naproxen sodium 220 mg 8875440

SEPARATE PILL PAPER, PUSH THROUGH PILL

EXP: 4/07 UIC: 287624D

MAXIMUM STRENGTH
Midol® Extended Relief
Dist. by: Bayer HealthCare LLC

READ CARTON BEFORE USE. Each caplet contains: Naproxen sodium 220 mg 8875440

SEPARATE PILL PAPER, PUSH THROUGH PILL

EXP: 4/07 UIC: 287624D

MAXIMUM STRENGTH
Midol® Extended Relief
Dist. by: Bayer HealthCare LLC

READ CARTON BEFORE USE. Each caplet contains: Naproxen sodium 220 mg 8875440

SEPARATE PILL PAPER, PUSH THROUGH PILL

EXP: 4/07 UIC: 287624D

MAXIMUM STRENGTH
Midol® Extended Relief
Dist. by: Bayer HealthCare LLC

READ CARTON BEFORE USE. Each caplet contains: Naproxen sodium 220 mg 8875440

SEPARATE PILL PAPER, PUSH THROUGH PILL

Drug Facts

Active ingredient
(in each caplet)

Naproxen sodium 220 mg
(naproxen 200 mg).....

Purpose

Pain reliever

Uses temporarily relieves minor aches and pains due to ● menstrual cramps ● muscle aches ● headache ● backache

Drug Facts (continued)

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction which may include: ● hives ● facial swelling ● asthma (wheezing) ● shock

Drug Facts (continued)

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take naproxen sodium or other pain relievers/fever reducers. Naproxen sodium may cause stomach bleeding.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if you have ever had serious side effects from any pain reliever

Ask a doctor or pharmacist before use if you are ● under a doctor's care for any serious condition ● taking any other drug ● taking any other product that contains naproxen sodium, or any other pain reliever/fever reducer

Stop use and ask a doctor if ● an allergic reaction occurs. Seek medical help right away. ● pain gets worse or lasts more than 10 days ● you have difficulty swallowing ● it feels like the pill is stuck in your throat ● you develop heartburn ● stomach pain occurs or lasts, even if symptoms are mild ● redness or swelling is present in the painful area ● any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ● do not take more than directed ● drink a full glass of water with each dose

| | |
|----------------|---|
| under 12 years | ● ask a doctor |
| 12 to 65 years | ● take 1 caplet every 8 to 12 hours while symptoms last. For the first dose you may take 2 caplets within the first hour. The smallest effective dose should be used. Do not exceed 2 caplets in any 8- to 12-hour period. Do not exceed 3 caplets in a 24-hour period. |
| over age 65 | ● do not take more than 1 caplet every 12 hours unless directed by a doctor |