

FPO  
DOES NOT PRINT  
DASHED BOX ONLY REPRESENTS  
LIVE AREA

Cat. No. 260800 NDC #054365-400-05 DIN #02160757

# ChloroPrep<sup>®</sup> With Tint

Chlorhexidine Gluconate 2% w/v  
and Isopropyl Alcohol 70% v/v  
Patient Preoperative Skin Preparation  
26-mL Applicator

**WARNING: FLAMMABLE.**  
KEEP AWAY FROM FIRE OR FLAME.  
DO NOT USE WITH ELECTROCAUTERY PROCEDURES

Ingredients:.....  
6 pt AntiqueOlive Bold

Title: .....  
14 pt helvetica black oblique

Headings: .....  
8 pt helvetica black oblique,  
left justified

0.5 point hairline

Subheadings: .....  
6 pt helvetica  
black,  
left justified

Bullet: .....  
5 pt zapf  
dingbats,  
solid square

## Drug Facts

**Active Ingredients Purpose**  
Chlorhexidine gluconate 2% w/v ..... Antiseptic  
Isopropyl alcohol 70% v/v ..... Antiseptic

**Use** for the preparation of the patient's skin prior to surgery

### Warnings

**For external use only**

**Flammable, keep away from fire or flame**

- Solution contains alcohol and gives off flammable vapors while drying - allow to dry 3 minutes on skin.
- Do not use with electrocautery procedures.
- Do not allow to solution to pool
- Remove solution-soaked materials from prep area

### Do not use

- in children less than 2 months of age because of the potential for excessive skin irritation and increased drug absorption
- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a physician.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

### Directions

■ **Maximal treatment area for one applicator is approximately 1126 cm<sup>2</sup> (approx. 13.2 in. x 13.2 in.). Discard the applicator after a single use.**

■ pinch the wings of the applicator to break the ampule and release the antiseptic. Do not touch the sponge. Wet the sponge by repeatedly pressing and releasing the sponge against the treatment area until liquid is visible on the skin.

■ **dry surgical sites** (such as abdomen or arm)  
Use repeated back-and-forth strokes of the sponge for approximately **30 seconds**. Completely wet the treatment area with antiseptic. Allow the area to air dry for approximately **three (3) minutes**. Do not blot or wipe away.

■ **moist surgical sites** (such as the inguinal fold)  
Use repeated back-and-forth strokes of the sponge for approximately **2 minutes**. Completely wet the treatment area with antiseptic. Allow the area to air dry for approximately **three (3) minutes**. Do not blot or wipe away.

### Other Information

- Store between 20-25 °C (68-77 °F)
- Avoid freezing and excessive heat above 40 °C (104 °F)

**Inactive ingredients** ■ USP purified water  
■ Pledget contains FD&C green #3 dye

### Questions?

Call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)  
www.chloroprep.com

Single Use  
Applicator is STERILE if package is intact

Medi-Flex, Inc.  
Leawood, KS 66211

2.35" wide

9.5" high

2.5 point  
box barline

Body Text:  
6 pt helvetica  
regular,  
6.5 leading,  
left justified

heading:  
8 pt helvetica  
black oblique

text:  
6 pt helvetica  
black

8-100800  
26ml Chloragreg With Tint Applicator (White Film)



0 50mm

312mm



**WARNING - FLAMMABLE**

- Keep away from fire or flame.
- To reduce the risk of fire:
  - Solution contains alcohol and gives off flammable vapour while drying - allow to dry 3 minutes on skin.
  - Do not use with hydrocarbon products.
  - Do not touch solution in pool.
  - Remove solution-coated material from prep area.

63.94mm


**WARNING - FLAMMABLE**

- Keep away from fire or flame.
- To reduce the risk of fire:
  - Solution contains alcohol and gives off flammable vapour while drying - allow to dry 3 minutes on skin.
  - Do not use with hydrocarbon products.
  - Do not touch solution in pool.
  - Remove solution-coated material from prep area.



24.308mm

Arial Black 8pt  
Arial Black 8pt  
Arial Bold 7pt

These bars in this area are for use with top edge and center marks and are not for use with side marks.  
NOTE: Text box center points both vertically and horizontally are used to place the text bar in correct position.



Part # 6-775800 Insert  
 Insert Sheet - Labeling Format (100%) for ChlorPrep® One-Step  
 Chlorhexidine Gluconate 2% w/v and Isopropyl Alcohol 70% v/v  
 Patient Preoperative Skin Preparation • 26-ml Applicator



6" wide

Numbers:  
 9 pt antique olive roman Title  
 18 pt antique olive bold  
 9 pt antique olive roman  
 8 pt CI Helvetica Condensed

Cat. No. 260800 NDC 05436S-400-05 DIN 02160757

## ChlorPrep® With Tint

### Patient Preoperative Skin Preparation

Chlorhexidine Gluconate 2% (w/v) • Isopropyl Alcohol 70% (v/v) • 26-ml Applicator

STERILE

Eti

External Use Only  
Professional Use Only

Do Not Reuse

Startle Contents\*  
Applicator (1)  
Cotton Swabs (2)

Lot No.: \_\_\_\_\_ Exp. Date: \_\_\_\_\_

\*Sterility of contents guaranteed unless package is damaged or open.

See Drug Facts for full drug labeling information

**WARNING**

- ChlorPrep solution contains alcohol and is flammable when drying.
- To prevent fires, follow instructions on this insert.

**Directions for Use:**  
Prepare Patient for ChlorPrep Solution.

- When hair removal is necessary, the preferred method is to use a surgical clipper on the morning of the surgery.
- Preparation should be applied to clean, dry, residue-free, intact skin.
- When applying the solution, start at the incision site and apply the solution in a back and forth or up and down motion.

**After ChlorPrep is Applied:**

- Allow solution to completely dry prior to draping.

**While Waiting for ChlorPrep to Dry:**

- Do not drape.
- Check for pooled preparation. Use sterile gauze to absorb pooled solution. Do not blot. Allow solution to dry.
- Remove all solution-soaked materials.

**When ChlorPrep Solution is Dry:**

- Begin draping only after solution is dry and all solution-soaked materials are removed.
- If incise drapes are used, apply directly to dry prepped area.

**Questions or Comments?**


- (800) 523-0602 (M-F 8 a.m. - 5 p.m. CST) • [www.chlorprep.com](http://www.chlorprep.com)

Statement  
9 pt CI Helvetica  
Condensed

6" high

Sub-Heading  
9 pt antique olive  
roman

Bullet  
8 pt Round CI  
Helvetica  
Condensed



-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/

-----  
Curtis Rosebraugh  
5/3/05 02:21:50 PM