



## Patient Information

### TAMIFLU®

(oseltamivir phosphate)

#### **R<sub>x</sub> only**

This leaflet contains important information about TAMIFLU (TAM-ih-flew). Read it well before you begin treatment. This information does not take the place of talking with your health care professional about your medical condition or your treatment. This leaflet does not list all the benefits and risks of TAMIFLU. If you have any questions about TAMIFLU, ask your health care professional. Only your health care professional can determine if TAMIFLU is right for you.

#### **What is TAMIFLU?**

TAMIFLU attacks the influenza virus and stops it from spreading inside your body. TAMIFLU treats flu at its source, by attacking the virus that causes the flu, rather than simply masking symptoms.

TAMIFLU is for treating adults and children age 1 and older with the flu whose flu symptoms started within the last day or two. TAMIFLU can also reduce the chance of getting the flu in people age 1 and older who have a higher chance of getting the flu because they spend time with someone who has the flu. TAMIFLU can also reduce the chance of getting the flu if there is a flu outbreak in the community.

#### **What is “Flu”?**

“The flu” is an infection caused by the influenza virus. Flu symptoms include fever (usually 100°F to 103°F in adults, and sometimes higher in children) and problems such as cough, sore throat, runny or stuffy nose, headaches, muscle aches, fever, and extreme tiredness. Many people use the term “flu” to mean any combination of these symptoms, such as the common cold, but true influenza infection is often worse and may last longer than a cold.

Flu outbreaks happen about once a year, usually in the winter, when the influenza virus spreads widely in the community. Outside of those outbreaks, only a very tiny number of respiratory infections are caused by the influenza virus.

#### **Should I get a flu shot?**

TAMIFLU is not a substitute for a flu vaccination. You should continue to get a flu vaccination every year, according to your health care professional’s advice.

## TAMIFLU® (oseltamivir phosphate)

### 34 **Who should not take TAMIFLU?**

35 Do not take TAMIFLU if you are allergic to the main ingredient, oseltamivir phosphate,  
36 or to any other ingredients of TAMIFLU. Before starting treatment, make sure your  
37 health care professional knows if you take any other medicines, or are pregnant, planning  
38 to become pregnant, or breastfeeding. TAMIFLU is normally not recommended for use  
39 during pregnancy or nursing, as the effects on the unborn child or nursing infant are  
40 unknown. TAMIFLU is not recommended for use in children younger than 1 year of age.

41 Tell your health care professional if you have any type of kidney disease, heart disease,  
42 respiratory disease, or any serious health condition.

### 43 **How should I take TAMIFLU?**

44 It is important that you begin your treatment with TAMIFLU as soon as possible from the  
45 first appearance of your flu symptoms or soon after you are exposed to the flu. If you feel  
46 worse or develop new symptoms during treatment with TAMIFLU, or if your flu  
47 symptoms do not start to get better, you should contact your health care professional.

48 If you have the flu: Take TAMIFLU twice a day for 5 days, once in the morning and  
49 once in the evening. You should complete the entire treatment of 10 doses (capsules or  
50 liquid), even if you feel better.

51 To prevent the flu: If someone in your home has the flu, take TAMIFLU once a day for  
52 10 days or for as long as prescribed. You can take TAMIFLU for up to 6 weeks if you are  
53 exposed to the flu because of an outbreak in your community. Follow your health care  
54 professional's advice on how long to take TAMIFLU.

55 TAMIFLU has not been studied in children 1 to 12 years of age for preventing flu during  
56 an outbreak in your community or for use for more than 10 days.

57 You can take TAMIFLU with food or without food. There is less chance of stomach  
58 upset if you take it with a light snack, milk, or a meal.

59 If you are taking TAMIFLU liquid, your pharmacist will give you a dosing dispenser  
60 marked with three possible doses. Follow your health care professional's instructions on  
61 which dose to take or how to combine them for the proper dose for you. In order to be  
62 sure you receive the proper dose, it is important that you use the dispenser provided.  
63 Review the instructions below on how to use the dispenser and ask your pharmacist if  
64 you have any questions. If you lose or damage the dispenser and cannot use it, contact  
65 your health care professional or pharmacist for advice on the proper dose.

66 If you forget to take your medicine, take the missed dose as soon as you remember,  
67 except if it is 2 hours or less before your next dose. Then continue to take TAMIFLU at  
68 the usual times. Do not take 2 doses at a time to make up for a missed dose. If you miss  
69 several doses, tell your health care professional and follow the advice given to you.

## TAMIFLU® (oseltamivir phosphate)

### 70 **What are the possible side effects of TAMIFLU?**

71 The most common side effects of TAMIFLU are nausea and vomiting. These are usually  
72 mild to moderate. They usually happen in the first 2 days of treatment. Taking TAMIFLU  
73 with food may reduce the chance of getting these side effects.

74 If you develop an allergic reaction or severe rash, stop taking TAMIFLU and contact  
75 your health care professional.

76 If you notice any side effects not mentioned in this leaflet, or if you have any concerns  
77 about the side effects you get, tell your health care professional.

### 78 **How and where should I store TAMIFLU?**

79 TAMIFLU capsules should be stored at room temperature below 77°F (25°C) and kept in  
80 a dry place. Keep this medication out of reach of children.

81 TAMIFLU suspension should be stored under refrigeration at 36° to 46°F (2° to 8°C). Do  
82 not freeze.

### 83 **General advice about prescription medicines:**

84 Medicines are sometimes prescribed for conditions that are not mentioned in patient  
85 information leaflets. Do not use TAMIFLU for a condition for which it was not  
86 prescribed. Do not give TAMIFLU to other people, even if they have the same symptoms  
87 you have. It may not be right for them.

88 This leaflet summarizes the most important information about TAMIFLU. If you would  
89 like more information, talk with your health care professional. You can ask your  
90 pharmacist or health care professional for information about TAMIFLU that is written for  
91 health professionals.

92

### 93 **DOSING INSTRUCTIONS FOR PATIENTS:**

94 **Please follow instructions carefully to ensure proper dosing of the oral suspension.**




95

- 96 • Shake closed bottle well for about 5 seconds before each use.
- 97 • Remove child-resistant cap.

## TAMIFLU® (oseltamivir phosphate)

- 98 • Before inserting the tip of the oral dispenser into bottle adapter, push the plunger  
99 completely down toward the tip of the oral dispenser. Insert tip firmly into opening of  
100 the bottle adapter.
- 101 • Turn the entire unit (bottle and oral dispenser) upside down.
- 102 • Pull the plunger out slowly until the desired amount of medication is withdrawn into  
103 the oral dispenser (see figure). The 75 mg dose is obtained by filling the dispenser  
104 twice, once to the 30 mg graduation, and a second fill to the 45 mg graduation.



- 105
- 106 • Turn the entire unit right side up and remove the oral dispenser slowly from the  
107 bottle.
- 108 • Dispense directly into mouth. Do not mix with any liquid prior to dispensing.
- 109 • Close bottle with child-resistant cap after each use.
- 110 • Disassemble oral dispenser, rinse under running tap water and air dry prior to next  
111 use.

112

113 Manufactured by:

114 F. Hoffmann-La Roche Ltd.

115 Basel, Switzerland

116 Distributed by:



**Pharmaceuticals**

Roche Laboratories Inc.  
340 Kingsland Street  
Nutley, New Jersey 07110-1199

117

118 Licensor:

119 Gilead Sciences, Inc.

120 Foster City, California 94404

121 Revised: Month Year

122 Copyright © 1999-xxxx by Roche Laboratories Inc. All rights reserved.

Filename: clean ppi 12.19.05.doc  
Directory: C:\Documents and Settings\jefferye\Local  
Settings\Temporary Internet Files\OLK6CA  
Template: U:\Application  
Data\Microsoft\Templates\US\_Labeling\_XP.dot  
Title: tamiflu  
Subject: oseltamivir phosphate  
Author:  
Keywords: 21087,21246,proposed,xxdec2005  
Comments: hoffmann-la roche 0004  
Creation Date: 12/19/2005 11:24 AM  
Change Number: 3  
Last Saved On: 12/19/2005 11:24 AM  
Last Saved By:  
Total Editing Time: 0 Minutes  
Last Printed On: 1/10/2006 10:12 AM  
As of Last Complete Printing  
Number of Pages: 4  
Number of Words: 1,124 (approx.)  
Number of Characters: 6,413 (approx.)