



## Patient Information About INVIRASE (in-ver-ase)

### INVIRASE<sup>®</sup>

(saquinavir mesylate)

CAPSULES and TABLETS

**Generic Name: Saquinavir mesylate (sa-KWIN-a-veer mes-il-late)**

#### **Rx only**

**ALERT: Find out about medicines that should NOT be taken with INVIRASE.** Please also read the section MEDICINES YOU SHOULD NOT TAKE WITH INVIRASE.

Please read this product information carefully before you start taking INVIRASE and each time you renew your prescription. There may be new information. Reading this information can help you take this medicine correctly. However, it is not a substitute for your doctor's advice about the safety and benefits of INVIRASE. You should talk to your doctor about INVIRASE as part of your long-term treatment plan for HIV before you start taking your medication and ask any questions you may have at regular checkups. Remember, you should remain under a doctor's care when using INVIRASE and should not change or stop your therapy without talking to your doctor first.

#### ***What is INVIRASE?***

INVIRASE belongs to a class of anti-HIV medicines called protease (PRO-tee-ase) inhibitors. INVIRASE Capsules and Tablets in combination with other anti-HIV drugs are used for the treatment of HIV, the virus that causes AIDS (acquired immunodeficiency syndrome).

#### ***How does INVIRASE work?***

INVIRASE fights HIV as it grows inside cells by blocking an enzyme (protease) that HIV needs to reproduce.

#### ***How is INVIRASE different from FORTOVASE<sup>®</sup> (saquinavir)?***

Both INVIRASE and FORTOVASE contain the same active ingredient—saquinavir. When INVIRASE or FORTOVASE are taken two times a day with ritonavir, a similar amount of saquinavir gets into the blood to fight HIV. However, FORTOVASE can be taken three times a day without ritonavir and the correct amount of saquinavir can get into the blood.

INVIRASE should never be taken without ritonavir. FORTOVASE may be taken without ritonavir if you are not able to tolerate even a small amount of ritonavir.

You should not substitute one for the other. If the medicine you receive does not look like the light brown and green capsule (INVIRASE) it is not the right drug. Talk to your doctor, nurse or pharmacist if you are not sure that you have the right medication.

### ***Who should not take INVIRASE?***

Anyone who has had a severe allergic reaction to INVIRASE or any of the ingredients in the capsule or tablet should not take it. The use of INVIRASE in patients under 16 years of age, over 65 years of age, or patients with severe liver problems has not been fully investigated.

### ***How should INVIRASE/Norvir® (ritonavir) be taken?***

- The recommended dosage of INVIRASE in combination with Norvir® (ritonavir) is INVIRASE 5 capsules or 2 tablets twice a day taken with 1 capsule of Norvir twice a day. In some combinations, your dose may change.
- INVIRASE must be taken along with Norvir (ritonavir).
- INVIRASE must be taken with meals or up to 2 hours after a meal—but it is easiest to remember if you take it with your meals. When INVIRASE is taken without food, the amount of INVIRASE in the blood is lower and may not fight HIV as well.
- When taking INVIRASE and other anti-HIV medicines, it is very important to follow the directions exactly and take your medication every day. If you skip doses—or take less than the prescribed dose—the medicine will not work as well, and your disease could get worse.
  - If you miss a dose, you should take the next dose as soon as possible. However, do not double the dose.

### ***What results have been seen with INVIRASE?***

INVIRASE with ritonavir has been shown to reduce the amount of virus in the blood (“viral load”) and increase CD<sub>4</sub> (T) cells when taken with other HIV therapy.

### ***What are the side effects of INVIRASE?***

People treated with INVIRASE in combination with Norvir may have side effects. The majority of these have been described as mild. In clinical studies of patients who received saquinavir in combination with Norvir and other HIV drugs the side effects seen most often were: body fat change (5.4%), nausea (10.8%), vomiting (7.4%), diarrhea (8.1%), stomach pain (6.1%), tiredness (6.1%), and pneumonia (5.4%).

Diabetes (new onset or worsening) and increased blood sugar levels have been reported with the use of protease inhibitors. In addition, increased bleeding in patients with hemophilia has also been associated with these drugs.

When saquinavir is taken with ritonavir, some patients may experience large increases in triglyceride and lipid levels. The long-term chance of getting complications such as heart attack and stroke due to increases in triglyceride and cholesterol levels caused by protease inhibitors is not known at this time.

Changes in body fat have been seen in some patients taking anti-HIV medications. These changes may include increased amount of fat in the upper back and neck (“buffalo hump”), breasts, and around the trunk. Loss of fat from the legs and arms may also happen. The cause and long-term health effects of these conditions are not known at this time.

These are not the only side effects that can occur with INVIRASE. Your doctor can discuss with you a more complete list of side effects and laboratory abnormalities that may accompany this medication.

If any side effects or unusual symptoms do occur, contact your doctor immediately. Do not stop or decrease your dose on your own. Lowering the dose may make INVIRASE less effective in fighting HIV.

***Are there other medications that I should not take with INVIRASE/Norvir (ritonavir)?***

There are some drugs that should not be taken with INVIRASE. Before starting therapy with INVIRASE, be sure to tell your doctor all of the medicines—prescription medications, as well as over-the-counter drugs and nutritional supplements—that you are now taking or plan to take.

**Medicines you should not take with INVIRASE**

Drug Class	Drugs Within Class Not to Be Taken with INVIRASE/Norvir (ritonavir)
Antiarrhythmics	Pacerone <sup>®</sup> (amiodarone), Tambacor <sup>®</sup> (flecainide), Rhythmol <sup>®</sup> (propafenone), bepridil, quinidine
Antihistamines	Seldane <sup>®</sup> (terfenadine)*, Hismanal <sup>®</sup> (astemizole)*
Antimigraines	Ergot medications (eg, Wigraine <sup>®</sup> and Cafergot <sup>®</sup> )
GI motility agents	Propulsid <sup>®</sup> (cisapride)*
Sedatives, hypnotics	Versed <sup>®</sup> (midazolam), Halcion <sup>®</sup> (triazolam)
Antimycobacterial agents	Rifampin
Neuroleptics	Pimozide

\* No longer sold in the US.

INVIRASE causes increased blood levels of these compounds. This can lead to serious or life-threatening reactions such as irregular heartbeat or prolonged sedation.

Taking INVIRASE with St. John's wort (*hypericum perforatum*), an herbal product sold as a dietary supplement, or products containing St. John's wort is not recommended. Talk with your doctor if you are taking or are planning to take St. John's wort. Taking St. John's wort may decrease INVIRASE levels and lead to increased viral load and possible resistance to INVIRASE or cross-resistance to other antiretroviral drugs.

Garlic capsules should not be used while taking FORTOVASE as the sole protease inhibitor due to the risk of decreased saquinavir in the blood. No data are available for the coadministration of FORTOVASE and Norvir with garlic capsules or INVIRASE and Norvir with garlic capsules.

Your doctor may want to change your medicine if you are taking rifampin (known as Rifadin<sup>®</sup>, Rifamate<sup>®</sup>, Rifater<sup>®</sup> or Rimactane<sup>®</sup>) or Mycobutin<sup>®</sup> (rifabutin); these drugs substantially reduce the level of INVIRASE in the blood. Rifampin, in combination with INVIRASE and ritonavir, may also cause severe liver problems.

The following drugs increase blood levels of INVIRASE: Norvir<sup>®</sup> (ritonavir)<sup>‡</sup>, Viracept<sup>®</sup> (nelfinavir)<sup>§</sup>, Rescriptor<sup>®</sup> (delavirdine)<sup>§||</sup>, Nizoral<sup>®</sup> (ketoconazole), Crixivan<sup>®</sup> (indinavir)<sup>§</sup> and Biaxin<sup>®</sup> (clarithromycin).

Talk to your doctor if you are taking lipid (cholesterol) lowering drugs and Viagra<sup>®</sup> (sildenafil citrate), Levitra<sup>®</sup> (vardenafil), and Cialis<sup>®</sup> (tadalafil).

### ***Does INVIRASE cure HIV/AIDS?***

INVIRASE does not cure AIDS, and it does not prevent you from getting other illnesses that result from advanced HIV infection. In addition, INVIRASE has not been shown to reduce the risk that you may transmit HIV to others through sexual contact or infected blood. You must continue to follow all of your doctor's recommendations for managing your illness.

### ***What else should I discuss with my doctor?***

Inform your doctor:

- If you are pregnant or become pregnant while taking INVIRASE. The effects of INVIRASE on pregnant women or unborn babies are not yet fully known. In addition, experts advise against breast-feeding if you are HIV positive, to reduce the risk of passing the virus to your baby.
- If you are taking anti-HIV medications. Your doctor may want to change one or more of your anti-HIV drugs in order to achieve the best results when you start treatment with INVIRASE.
- If you have diabetes or a family history of diabetes, or if you have hemophilia, hepatitis or other liver disease, your doctor should decide if INVIRASE is right for you.

- If you have ever taken FORTOVASE, discuss with your doctor whether INVIRASE is right for you.

### ***How is INVIRASE supplied?***

INVIRASE is available as light brown and green capsules in a 200-mg strength. INVIRASE comes in bottles of 270 capsules.

INVIRASE is also available as light orange to greyish- or brownish-orange tablets in a 500-mg strength. INVIRASE comes in bottles of 120 tablets.

### ***How should I store INVIRASE?***

INVIRASE capsules and tablets should be stored at room temperature. The bottles should be kept tightly closed.

INVIRASE has been prescribed specifically for you, and only for a particular condition. Do not use it for anything else. Do not give it to anyone else. If you think you have taken more than your prescribed dose, seek medical attention.

**Keep this medication and all other medications out of the reach of children.** Do not keep medicine that is out of date or that you no longer need. Be sure that if you throw any medicine away, it is out of the reach of children.

This provides only a brief summary of product information about INVIRASE. If you have any questions about INVIRASE or HIV, talk to your doctor.

† Below the amount that could be found using a standard test.

‡ Dosages greater than 100 mg twice a day of ritonavir when taken in combination with saquinavir were associated with an increase in side effects.

§ The safety and efficacy of INVIRASE in combination with these drugs has not been established. Dosage adjustments may be required.

|| Use of this combination should be accompanied by close monitoring of liver enzymes.

FORTOVASE and Versed are registered trademarks of Hoffmann-La Roche Inc. Norvir, Rhythmol, and Biaxin are registered trademarks of Abbott Laboratories. Halcion, Mycobutin and Rescriptor are registered trademarks of Pharmacia & Upjohn Co. Hismanal, Propulsid and Nizoral are registered trademarks of Janssen Pharmaceutica Inc. Seldane, Rifadin, Rifamate and Rifater are registered trademarks of Hoechst Marion Roussel. Rimactane and Cafegot are registered trademarks of Novartis Pharmaceuticals Corporation. Viracept is a registered trademark of Agouron Pharmaceuticals Inc. Crixivan is a registered trademark of Merck & Co., Inc. Viagra is a registered trademark of Pfizer, Inc. Levitra is a registered trademark of Bayer Pharmaceuticals Corp. Cialis is a registered trademark of Eli Lilly and Company. Tambocor is a registered trademark of 3M. Pacerone is a registered trademark of Upsher-Smith.

*If you have any questions about INVIRASE, call toll free at 1-800-910-4687.*

Distributed by:



**Pharmaceuticals**

Roche Laboratories Inc.  
340 Kingsland Street  
Nutley, New Jersey 07110-1199

XXXXXXXXXX

XXXXXXXXXX

Revised: Month/Year

Copyright © 1999-XXXX by Roche Laboratories Inc. All rights reserved.