

SOLTAMOX™ Oral Solution  
Rx Only

**WARNING - For Women with Ductal Carcinoma in Situ (DCIS) and Women at High Risk for Breast Cancer:** Serious and life-threatening events associated with tamoxifen in the risk reduction setting (women at high risk for cancer and women with DCIS) include uterine malignancies, stroke and pulmonary embolism. Incidence rates for these events were estimated from the NSABP P-1 trial (see **CLINICAL PHARMACOLOGY, Clinical Studies, Reduction in Breast Cancer Incidence In High Risk Women**).

Uterine malignancies consist of both endometrial adenocarcinoma (incidence rate per 1,000 women-years of 2.20 for tamoxifen vs. 0.71 for placebo) and uterine sarcoma (incidence rate per 1,000 women years of 0.17 for tamoxifen vs. 0.0 for placebo) \*. For stroke, the incidence rate per 1,000 women years was 1.43 for tamoxifen vs. 1.00 for placebo\*\*. For pulmonary embolism, the incidence rate per 1,000 women years was 0.75 for tamoxifen versus 0.25 for placebo \* \*.

Some of the strokes, pulmonary emboli, and uterine malignancies were fatal.

Health care providers should discuss the potential benefits versus the potential risks of these serious events with women at high risk of breast cancer and women with DCIS considering tamoxifen to reduce their risk of developing breast cancer.

The benefits of tamoxifen outweigh its risks in women already diagnosed with breast cancer.


\* Updated long-term follow-up data (median length of follow-up is 6.9 years) from NSABP P-1 study. See **WARNINGS, Effects on the Uterus-Endometrial Cancer and Uterine Sarcoma**.

\*\* See Table 3 under **CLINICAL PHARMACOLOGY, Clinical Studies**.

## DESCRIPTION

SOLTAMOX™ solution, a nonsteroidal antiestrogen, is for oral administration. Each 5 mL solution contains 15.2 mg tamoxifen citrate, equivalent to 10 mg tamoxifen and the following inactive ingredients: ethanol, glycerol, propylene glycol, sorbitol solution, licorice flavor, aniseed flavor, purified water.

The chemical name is (Z)-2-[4-(1,2-diphenyl-1-butenyl) phenoxy]-N, N-dimethylethanamine 2-hydroxy-1,2,3- propanetricarboxylate (1:1). The structural and empirical formulas are:



Tamoxifen citrate has a molecular weight of 563.62, the pKa' is 8.85, the equilibrium solubility in water at 37°C is 0.5 mg/mL and in 0.02 N HCl at 37°C, it is 0.2 mg/mL.

## CLINICAL PHARMACOLOGY

Tamoxifen citrate is a nonsteroidal agent that has demonstrated potent antiestrogenic properties in animal test systems. The antiestrogenic effects may be related to its ability to compete with estrogen for binding sites in target tissues such as breast. Tamoxifen inhibits the induction of rat mammary carcinoma induced by dimethylbenzanthracene (DMBA) and causes the regression of already established DMBA-induced tumors. In this rat model, tamoxifen appears to exert its antitumor effects by binding the estrogen receptors.

In cytosols derived from human breast adenocarcinomas, tamoxifen competes with estradiol for estrogen receptor protein.

### Absorption and Bioavailability

A pharmacokinetic study was performed in healthy perimenopausal and postmenopausal female subjects to evaluate the bioavailability of SOLTAMOX™ (n=30) in comparison with the commercially available tamoxifen citrate tablets (n=33) under fasting conditions. A third arm evaluated the effect of food on SOLTAMOX™ (n=16 evaluable). The rate and extent of absorption of SOLTAMOX™ was found to be bioequivalent to that of tamoxifen citrate tablets under fasting conditions.

In the food effect arm, the  $C_{max}$  and AUC were comparable to the fasting group.  $T_{max}$  was slightly longer in the fed group. There was no difference in bioavailability of SOLTAMOX™ Oral Solution between fed and fasting states, and therefore SOLTAMOX™ can be given without regard to meals.

Following a single oral dose of 20 mg tamoxifen, an average peak plasma concentration of 40 ng/mL (range 35 to 45 ng/mL) occurred approximately 5 hours after dosing. The decline in plasma concentrations of tamoxifen is biphasic with a terminal elimination half-life of about 5 to 7 days. The average peak plasma concentration of N-desmethyl tamoxifen is 15 ng/mL (range 10 to 20 ng/mL). Chronic administration of 10 mg tamoxifen given twice daily for 3 months to patients results in average steady-state plasma concentrations of 120 ng/mL (range 67-183 ng/mL) for tamoxifen and 336 ng/mL (range 148-654 ng/mL) for N-desmethyl tamoxifen. The average steady-state plasma concentrations of tamoxifen and N-desmethyl tamoxifen after administration of 20 mg tamoxifen once daily for 3 months are 122 ng/mL (range 71-183 ng/mL) and 353 ng/mL (range 152-706 ng/mL), respectively. After initiation of therapy, steady-state concentrations for tamoxifen are achieved in about 4 weeks and steady-state concentrations for N-desmethyl tamoxifen are achieved in about 8 weeks, suggesting a half-life of approximately 14 days for this metabolite. In a steady-state, crossover study of 10 mg tamoxifen tablets given twice a day vs. a 20 mg tamoxifen tablet given once daily, the 20 mg tamoxifen tablet was bioequivalent to the 10 mg tamoxifen tablets.

### Metabolism

Tamoxifen is extensively metabolized after oral administration. N-desmethyl tamoxifen is the major metabolite found in patients' plasma. The biological activity of N-desmethyl tamoxifen appears to be similar to that of tamoxifen. 4-Hydroxytamoxifen and a side chain primary alcohol derivative of tamoxifen have been identified as minor metabolites in plasma. Tamoxifen is a substrate of cytochrome P-450 3A, 2C9 and 2D6, and an inhibitor of P-glycoprotein.

## **Excretion**

Studies in women receiving 20 mg of <sup>14</sup>C tamoxifen have shown that approximately 65% of the administered dose was excreted from the body over a period of 2 weeks with fecal excretion as the primary route of elimination. The drug is excreted mainly as polar conjugates, with unchanged drug and unconjugated metabolites accounting for less than 30% of the total fecal radioactivity.

## **Special Populations**

The effects of age, gender and race on the pharmacokinetics of tamoxifen have not been determined. The effects of reduced liver function on the metabolism and pharmacokinetics of tamoxifen have not been determined.

## **Pediatric Patients**

The use of SOLTAMOX™ in pediatric patients has not been evaluated.

## **Drug-drug Interactions**

*In vitro* studies showed that erythromycin, cyclosporine, nifedipine and diltiazem competitively inhibited formation of N-desmethyl tamoxifen with apparent  $K_i$  of 20, 1, 45 and 30  $\mu$ M, respectively. The clinical significance of these *in vitro* studies is unknown.

Tamoxifen reduced the plasma concentration of letrozole by 37% when these drugs were co-administered. Rifampin, a cytochrome P-450 3A4 inducer reduced tamoxifen AUC and  $C_{max}$  by 86% and 55%, respectively. Aminoglutethimide reduces tamoxifen and N-desmethyl tamoxifen plasma concentrations. Medroxyprogesterone reduces plasma concentrations of N-desmethyl, but not tamoxifen.

## **Clinical Studies**

### **Metastatic Breast Cancer:**

*Premenopausal Women (Tamoxifen Citrate vs. Ablation)* - Three prospective, randomized studies (Ingle, Pritchard, Buchanan) compared tamoxifen to ovarian ablation (oophorectomy or ovarian irradiation) in premenopausal women with advanced breast cancer. Although the objective response rate, time to treatment failure, and survival were similar with both treatments, the limited patient accrual prevented a demonstration of equivalence. In an overview analysis of survival data from the 3 studies, the hazard ratio for death (tamoxifen /ovarian ablation) was 1.00 with two-sided 95% confidence intervals of 0.73 to 1.37. Elevated serum and plasma estrogens have been observed in premenopausal women receiving tamoxifen, but the data from the randomized studies do not suggest an adverse effect of this increase. A limited number of premenopausal patients with disease progression during tamoxifen therapy responded to subsequent ovarian ablation.

### **Male Breast Cancer:**

Published results from 122 patients (119 evaluable) and case reports in 16 patients (13 evaluable) treated with tamoxifen have shown that tamoxifen is effective for the palliative treatment of male breast cancer. Sixty-six of these 132 evaluable patients responded to tamoxifen which constitutes a 50% objective response rate.

### **Adjuvant Breast Cancer:**

#### *Overview*

The Early Breast Cancer Trialists' Collaborative Group (EBCTCG) conducted worldwide overviews of systemic adjuvant therapy for early breast cancer in 1985, 1990, and again in

1995. In 1998, 10-year outcome data were reported for 36,689 women in 55 randomized trials of adjuvant tamoxifen using doses of 20-40 mg/day for 1-5+ years. Twenty-five percent of patients received 1 year or less of trial treatment, 52% received 2 years, and 23% received about 5 years. Forty-eight percent of tumors were estrogen receptor (ER) positive (> 10 fmol/mg), 21% were ER poor (< 10 fmol/l), and 31% were ER unknown. Among 29,441 patients with ER positive or unknown breast cancer, 58% were entered into trials comparing tamoxifen to no adjuvant therapy and 42% were entered into trials comparing tamoxifen in combination with chemotherapy vs. the same chemotherapy alone. Among these patients, 54% had node positive disease and 46% had node negative disease.

Among women with ER positive or unknown breast cancer and positive nodes who received about 5 years of treatment, overall survival at 10 years was 61.4% for tamoxifen vs. 50.5% for control (logrank 2p < 0.00001). The recurrence-free rate at 10 years was 59.7% for tamoxifen vs. 44.5% for control (logrank 2p < 0.00001). Among women with ER positive or unknown breast cancer and negative nodes who received about 5 years of treatment, overall survival at 10 years was 78.9% for tamoxifen vs. 73.3% for control (logrank 2p < 0.00001). The recurrence-free rate at 10 years was 79.2% for tamoxifen versus 64.3% for control (logrank 2p < 0.00001).

The effect of the scheduled duration of tamoxifen may be described as follows. In women with ER positive or unknown breast cancer receiving 1 year or less, 2 years or about 5 years of tamoxifen, the proportional reductions in mortality were 12%, 17% and 26%, respectively (trend significant at 2p < 0.003). The corresponding reductions in breast cancer recurrence were 21%, 29% and 47% (trend significant at 2p < 0.00001).

Benefit is less clear for women with ER poor breast cancer in whom the proportional reduction in recurrence was 10% (2p = 0.007) for all durations taken together, or 9% (2p = 0.02) if contralateral breast cancers are excluded. The corresponding reduction in mortality was 6% (NS). The effects of about 5 years of tamoxifen on recurrence and mortality were similar regardless of age and concurrent chemotherapy. There was no indication that doses greater than 20 mg per day were more effective.

#### *Node Positive - Individual Studies:*

Two studies (Hubay and NSABP B-09) demonstrated an improved disease-free survival following radical or modified radical mastectomy in postmenopausal women or women 50 years of age or older with surgically curable breast cancer with positive axillary nodes when tamoxifen was added to adjuvant cytotoxic chemotherapy. In the Hubay study, tamoxifen was added to "low-dose" CMF (cyclophosphamide, methotrexate and fluorouracil). In the NSABP B-09 study, tamoxifen was added to melphalan [L-phenylalanine mustard (P)] and fluorouracil (F).

In the Hubay study, patients with a positive (more than 3 fmol) estrogen receptor were more likely to benefit. In the NSABP B-09 study in women age 50-59 years, only women with both estrogen and progesterone receptor levels 10 fmol or greater clearly benefited, while there was a nonstatistically significant trend toward adverse effect in women with both estrogen and progesterone receptor levels less than 10 fmol. In women age 60-70 years, there was a trend toward a beneficial effect of tamoxifen without any clear relationship to estrogen or progesterone receptor status.

Three prospective studies (ECOG-1178, Toronto, NATO) using tamoxifen adjuvantly as a single agent demonstrated an improved disease-free survival following total mastectomy and axillary dissection for postmenopausal women with positive axillary nodes compared

to placebo/no treatment controls. The NATO study also demonstrated an overall survival benefit.

*Node Negative - Individual Studies:*

NSABP B-14, a prospective, double-blind, randomized study, compared tamoxifen to placebo in women with axillary node-negative, estrogen-receptor positive ( $\geq 10$  fmol/mg cytosol protein) breast cancer (as adjuvant therapy, following total mastectomy and axillary dissection, or segmental resection, axillary dissection, and breast radiation). After five years of treatment, there was a significant improvement in disease-free survival in women receiving tamoxifen. This benefit was apparent both in women under age 50 and in women at or beyond age 50.

One additional randomized study (NATO) demonstrated improved disease-free survival for tamoxifen compared to no adjuvant therapy following total mastectomy and axillary dissection in postmenopausal women with axillary node-negative breast cancer. In this study, the benefits of tamoxifen appeared to be independent of estrogen receptor status.

*Duration of Therapy:*

In the EBCTCG 1995 overview, the reduction in recurrence and mortality was greater in those studies that used tamoxifen for about 5 years than in those that used tamoxifen for a shorter period of therapy.

In the NSABP B-14 trial, in which patients were randomized to tamoxifen 20 mg/day for 5 years vs. placebo and were disease-free at the end of this 5-year period were offered rerandomization to an additional 5 years of tamoxifen or placebo. With 4 years of follow-up after this rerandomization, 92% of the women that received 5 years of tamoxifen were alive and disease-free, compared to 86% of the women scheduled to receive 10 years of tamoxifen ( $p=0.003$ ). Overall survivals were 96% and 94%, respectively ( $p=0.08$ ). Results of the B-14 study suggest that continuation of therapy beyond 5 years does not provide additional benefit.

A Scottish trial of 5 years of tamoxifen vs. indefinite treatment found a disease-free survival of 70% in the five-year group and 61% in the indefinite group, with 6.2 years median follow-up (HR= 1.27, 95% CI: 0.87-1.85).

In a large randomized trial conducted by the Swedish Breast Cancer Cooperative Group of adjuvant tamoxifen 40 mg/day for 2 or 5 years, overall survival at 10 years was estimated to be 80% in the patients in the 5-year tamoxifen group, compared with 74% among corresponding patients in the 2-year treatment group ( $p=0.03$ ). Disease-free survival at 10 years was 73% in the 5-year group and 67% in the 2-year group ( $p=0.009$ ). Compared with 2 years of tamoxifen treatment, 5 years of treatment resulted in a slightly greater reduction in the incidence of contralateral breast cancer at 10 years, but this difference was not statistically significant.

*Contralateral Breast Cancer:*

The incidence of contralateral breast cancer is reduced in breast cancer patients (premenopausal and postmenopausal) receiving tamoxifen compared to placebo. Data on contralateral breast cancer are available from 32,422 out of 36,689 patients in the 1995 overview analysis of the Early Breast Cancer Trialists Collaborative Group (EBCTCG). In clinical trials with tamoxifen of 1 year or less, 2 years, and about 5 years duration, the proportional reductions in the incidence rate of contralateral breast cancer among women

receiving tamoxifen were 13% (NS), 26% (2p = 0.004) and 47% (2p < 0.00001), with a significant trend favoring longer tamoxifen duration (2p = 0.008). The proportional reductions in the incidence of contralateral breast cancer were independent of age and ER status of the primary tumor. Treatment with about 5 years of tamoxifen reduced the annual incidence rate of contralateral breast cancer from 7.6 per 1,000 patients in the control group compared with 3.9 per 1,000 patients in the tamoxifen group.

In a large randomized trial in Sweden (the Stockholm Trial) of adjuvant tamoxifen 40 mg/day for 2-5 years, the incidence of second primary breast tumors was reduced 40% (p < 0.008) on tamoxifen compared to control. In the NSABP B-14 trial in which patients were randomized to tamoxifen 20 mg/day for 5 years vs. placebo, the incidence of second primary breast cancers was also significantly reduced (p < 0.01). In NSABP B-14, the annual rate of contralateral breast cancer was 8.0 per 1,000 patients in the placebo group compared with 5.0 per 1,000 patients in the tamoxifen group, at 10 years after first randomization.

#### Ductal Carcinoma in Situ:

NSABP B-24, a double-blind, randomized trial included women with ductal carcinoma in situ (DCIS). This trial compared the addition of tamoxifen or placebo to treatment with lumpectomy and radiation therapy for women with DCIS. The primary objective was to determine whether 5 years of tamoxifen therapy (20 mg/day) would reduce the incidence of invasive breast cancer in the ipsilateral (the same) or contralateral (the opposite) breast.

In this trial 1,804 women were randomized to receive either tamoxifen or placebo for 5 years: 902 women were randomized to tamoxifen 10 mg tablets twice a day and 902 women were randomized to placebo. As of December 31, 1998, follow-up data were available for 1,798 women and the median duration of follow-up was 74 months.

The tamoxifen and placebo groups were well balanced for baseline demographic and prognostic factors. Over 80% of the tumors were less than or equal to 1 cm in their maximum dimension, were not palpable, and were detected by mammography alone. Over 60% of the study population was postmenopausal. In 16% of patients, the margin of the resected specimen was reported as being positive after surgery. Approximately half of the tumors were reported to contain comedo necrosis.

For the primary endpoint, the incidence of invasive breast cancer was reduced by 43% among women assigned to tamoxifen (44 cases - tamoxifen, 74 cases - placebo; p=0.004; relative risk (RR)=0.57, 95% CI: 0.39-0.84). No data are available regarding the ER status of the invasive cancers. The stage distribution of the invasive cancers at diagnosis was similar to that reported annually in the SEER data base.

Results are shown in Table 1. For each endpoint the following results are presented: the number of events and rate per 1,000 women per year for the placebo and tamoxifen groups; and the relative risk (RR) and its associated 95% confidence interval (CI) between tamoxifen and placebo. Relative risks less than 1.0 indicate a benefit of tamoxifen therapy. The limits of the confidence intervals can be used to assess the statistical significance of the benefits of tamoxifen therapy. If the upper limit of the CI is less than 1.0, then a statistically significant benefit exists.

**Table 1**  
**Major Outcomes of the NSABP B-24 Trial**

Type of Event	Lumpectomy, radiotherapy, and placebo		Lumpectomy, radiotherapy, and tamoxifen		RR	95% CI Limits
	No. of events	Rate per 1000 women per year	No. of events	Rate per 1000 women per year		
<b>Invasive breast cancer (Primary endpoint)</b>	<b>74</b>	<b>16.73</b>	<b>44</b>	<b>9.60</b>	<b>0.57</b>	<b>0.39 to 0.84</b>
-Ipsilateral	47	10.61	27	5.90	0.56	0.33 to 0.91
-Contralateral	25	5.64	17	3.71	0.66	0.33 to 1.27
-Side undetermined	2	-	0	-	-	
<b>Secondary Endpoints</b>						
DCIS	56	12.66	41	8.95	0.71	0.46 to 1.08
-Ipsilateral	46	10.40	38	8.29	0.88	0.51 to 1.25
-Contralateral	10	2.26	3	0.65	0.29	0.05 to 1.13
All Breast Cancer Events	129	29.16	84	18.34	0.63	0.47 to 0.83
-All Ipsilateral events	96	21.70	65	14.19	0.65	0.47 to 0.91
-All Contralateral events	37	8.36	20	4.37	0.52	0.29 to 0.92
Deaths	32		28			
Uterine Malignancies <sup>1</sup>	4		9			
Endometrial Adenocarcinoma <sup>1</sup>	4	0.57	8	1.15		
Uterine Sarcoma <sup>1</sup>	0	0.0	1	0.14		
Secondary primary malignancies (other than endometrial and breast)	30		29			
Stroke	2		7			
Thromboembolic events (DVT, PE)	5		15			

<sup>1</sup>Updated follow-up data (median 8.1 years)

Survival was similar in the placebo and tamoxifen groups. At 5 years from study entry, survival was 97% for both groups.

#### Reduction in Breast Cancer Incidence in High Risk Women:

The Breast Cancer Prevention Trial (BCPT, NSABP P-1) was a double-blind, randomized, placebo-controlled trial with a primary objective to determine whether 5 years of tamoxifen therapy (20 mg/day) would reduce the incidence of invasive breast cancer in women at high risk for the disease (See **INDICATIONS AND USAGE**). Secondary objectives included an evaluation of the incidence of ischemic heart disease; the effects on the incidence of bone fractures; and other events that might be associated with the use of tamoxifen, including: endometrial cancer, pulmonary embolus, deep vein thrombosis, stroke, and cataract formation and surgery (See **WARNINGS**).

The Gail Model was used to calculate predicted breast cancer risk for women who were less than 60 years of age and did not have lobular carcinoma in situ (LCIS). The following risk factors were used: age; number of first-degree female relatives with breast cancer; previous breast biopsies; presence or absence of atypical hyperplasia; nulliparity; age at first live birth; and age at menarche. A 5-year predicted risk of breast cancer of  $\geq 1.67\%$  was required for entry into the trial.

In this trial, 13,388 women of at least 35 years of age were randomized to receive either tamoxifen or placebo for five years. The median duration of treatment was 3.5 years. As

of January 31, 1998, follow-up data is available for 13,114 women. Twenty-seven percent of women randomized to placebo (1,782) and 24% of women randomized to tamoxifen (1,596) completed 5 years of therapy. The demographic characteristics of women on the trial with follow-up data are shown in Table 2.

**Table 2**  
**Demographic Characteristics of Women in the NSABP P-1 Trial**

Characteristic	Placebo		Tamoxifen	
	#	%	#	%
<b>Age (yrs.)</b>				
35-39	184	3	158	2
40-49	2,394	36	2,411	37
50-59	2,011	31	2,019	31
60-69	1,588	24	1,563	24
≥70	393	6	393	6
<b>Age at first live birth (yrs.)</b>				
Nulliparous	1,202	18	1,205	18
12-19	915	14	946	15
20-24	2,448	37	2,449	37
25-29	1,399	21	1,367	21
≥30	606	9	577	9
<b>Race</b>				
White	6,333	96	6,323	96
Black	109	2	103	2
Other	128	2	118	2
<b>Age at menarche</b>				
≥14	1,243	19	1,170	18
12-13	3,610	55	3,610	55
≤11	1,717	26	1,764	27
<b># of first degree relatives with breast cancer</b>				
0	1,584	24	1,525	23
1	3,714	57	3,744	57
2+	1,272	19	1,275	20
<b>Prior Hysterectomy</b>				
No	4,173	63.5	4,018	62.4
Yes	2,397	36.5	2,464	37.7
<b># of previous breast biopsies</b>				
0	2,935	45	2,923	45
1	1,833	28	1,850	28
≥2	1,802	27	1,771	27
<b>History of atypical hyperplasia in the breast</b>				
No	5,958	91	5,969	91
Yes	612	9	575	9
<b>History of LCIS at entry</b>				
No	6,165	94	6,135	94
Yes	405	6	409	6
<b>5-year predicted breast cancer risk (%)</b>				
≤2.00	1,646	25	1,626	25
2.01-3.00	2,028	31	2,057	31
3.01-5.00	1,787	27	1,707	26
≥5.01	1,109	17	1,162	18
<b>Total</b>	<b>6,570</b>	<b>100.0</b>	<b>6,544</b>	<b>100.0</b>

Results are shown in Table 3. After a median follow-up of 4.2 years, the incidence of invasive breast cancer was reduced by 44% among women assigned to tamoxifen (86 cases-tamoxifen, 156 cases-placebo;  $p < 0.00001$ ; relative risk (RR)=0.56, 95% CI: 0.43-0.72). A reduction in the incidence of breast cancer was seen in each prospectively specified age group ( $\leq 49$ , 50-59,  $\geq 60$ ), in women with or without LCIS, and in each of the absolute risk levels specified in Table 3. A non-significant decrease in the incidence of ductal carcinoma in situ (DCIS) was seen (23-tamoxifen, 35-placebo; RR=0.66; 95% CI: 0.39-1.11).

There was no statistically significant difference in the number of myocardial infarctions, severe angina, or acute ischemic cardiac events between the two groups (61-tamoxifen, 59-placebo; RR=1.04, 95% CI: 0.73-1.49).

No overall difference in mortality (53 deaths in tamoxifen group vs. 65 deaths in placebo group) was present. No difference in breast cancer-related mortality was observed (4 deaths in tamoxifen group vs. 5 deaths in placebo group).

Although there was a non-significant reduction in the number of hip fractures (9 on tamoxifen, 20 on placebo) in the tamoxifen group, the number of wrist fractures was similar in the two treatment groups (69 on tamoxifen, 74 on placebo). No information regarding bone mineral density or other markers of osteoporosis is available.

The risks of tamoxifen therapy include endometrial cancer, DVT, PE, stroke, cataract formation and cataract surgery (See **Table 3**). In the NSABP P-1 trial, 33 cases of endometrial cancer were observed in the tamoxifen group vs. 14 in the placebo group (RR=2.48, 95% CI: 1.27-4.92). Deep vein thrombosis was observed in 30 women receiving tamoxifen vs. 19 in women receiving placebo (RR=1.59, 95% CI: 0.86-2.98). Eighteen cases of pulmonary embolism were observed in the tamoxifen group vs. 6 in the placebo group (RR=3.01, 95% CI: 1.15-9.27). There were 34 strokes on the tamoxifen arm and 24 on the placebo arm (RR=1.42; 95% CI 0.82-2.51). Cataract formation in women without cataracts at baseline was observed in 540 women taking tamoxifen vs. 483 women receiving placebo (RR=1.13, 95% CI: 1.00-1.28). Cataract surgery (with or without cataracts at baseline) was performed in 201 women taking tamoxifen vs. 129 women receiving placebo (RR=1.51, 95% CI 1.21-1.89) (See **WARNINGS**).

Table 3 summarizes the major outcomes of the NSABP P-1 trial. For each endpoint, the following results are presented: the number of events and rate per 1,000 women per year for the placebo and tamoxifen groups; and the relative risk (RR) and its associated 95% confidence interval (CI) between tamoxifen and placebo. Relative risks less than 1.0 indicate a benefit of tamoxifen therapy. The limits of the confidence intervals can be used to assess the statistical significance of the benefits or risks of tamoxifen therapy. If the upper limit of the CI is less than 1.0, then a statistically significant benefit exists.

For most participants, multiple risk factors would have been required for eligibility. This table considers risk factors individually, regardless of other co-existing risk factors, for women who developed breast cancer. The 5-year predicted absolute breast cancer risk accounts for multiple risk factors in an individual and should provide the best estimate of individual benefit (See **INDICATIONS AND USAGE**).

**Table 3**  
**Major Outcomes of the NSABP-1 Trial**

	# of Events		Rate/1000 Women/Year		RR	95% CI Limits
<b>Type of Event</b>	<b>Placebo</b>	<b>Tamoxifen</b>	<b>Placebo</b>	<b>Tamoxifen</b>		
<b>Invasive Breast Cancer</b>	156	86	6.49	3.58	0.56	0.43-0.72
Age ≤49	59	38	6.34	4.11	0.65	0.43-0.98
Age 50-59	46	25	6.31	3.53	0.56	0.35-0.91
Age ≥60	51	23	7.17	3.22	0.45	0.27-0.74
<b>Risk Factors for Breast Cancer History, LCIS</b>						
No	140	78	6.23	3.51	0.56	0.43-0.74
Yes	16	8	12.73	6.33	0.50	0.21-1.17
<b>History, Atypical Hyperplasia</b>						
No	138	84	6.37	3.89	0.61	0.47-0.80
Yes	18	2	8.69	1.05	0.12	0.03-0.52
<b>No. First Degree Relatives</b>						
0	32	17	5.97	3.26	0.55	0.30-0.98
1	80	45	5.81	3.31	0.57	0.40-0.82
2	35	18	8.92	4.67	0.52	0.30-0.92
≥3	9	6	13.33	7.58	0.57	0.20-1.59
<b>5-Year Predicted Breast Cancer (as calculated by the Gail Model)</b>						
≤2.00%	31	13	5.36	2.26	0.42	0.22-0.81
2.01-3.00%	39	28	5.25	3.83	0.73	0.45-1.18
3.01-5.00%	36	26	5.37	4.06	0.76	0.46-1.26
≥5.00%	50	19	13.15	4.71	0.36	0.21-0.61
<b>DCIS</b>	35	23	1.47	0.97	0.66	0.39-1.11
<b>Fractures (protocol-specified sites)</b>						
Hip	92 <sup>1</sup>	76 <sup>1</sup>	3.87	3.20	0.61	0.83-1.12
Wrist <sup>2</sup>	20	9	0.84	0.38	0.45	0.18-1.04
<b>Totalschemic Events</b>	59	61	2.47	2.57	1.04	0.71-1.51
Myocardial Infarction	27	27	1.13	1.13	1.00	0.57-1.78
Fatal	8	7	0.33	0.29	0.88	0.27-2.77
Nonfatal	19	20	0.79	0.84	1.06	0.54-2.09
Angina <sup>3</sup>	12	12	0.50	0.50	1.00	0.41-2.44
Acute Ischemic Syndrome <sup>4</sup>	20	22	0.84	0.92	1.11	0.58-2.13
<b>Uterine</b>						
Malignancies (among women with an intact uterus) <sup>10</sup>						
Endometrial	17	57	0.71	2.20		
Adenocarcinoma <sup>10</sup>	0	4	0.0	0.17		
Uterine Sarcoma <sup>10</sup>						
<b>Stroke<sup>5</sup></b>	24	34	1.00	1.43	1.42	0.82-2.51
<b>Transient Ischemic Attack</b>	21	18	0.88	0.75	0.86	0.43-1.70
<b>Pulmonary Emboli<sup>6</sup></b>	6	18	0.25	0.75	3.01	1.15-9.27
<b>Deep-Vein Thrombosis<sup>6</sup></b>	19	30	0.79	1.26	1.59	0.86-2.98
<b>Cataracts Developing on Study<sup>8</sup></b>	483	540	22.51	25.41	1.13	1.00-1.28
<b>Underwent Cataract Surgery<sup>8</sup></b>	63	101	2.83	4.57	1.62	1.18-2.22
<b>Underwent Cataract Surgery<sup>9</sup></b>	1.29	201	5.44	8.56	1.58	1.26-1.97

1 Two women had hip and wrist fractures

2 Includes Colles' and other lower radius fractures

3 Requiring angioplasty or CABG

4 New Q-wave on ECG; no angina or elevation of serum enzymes; or angina requiring hospitalization without surgery

5 Seven cases were fatal; three in the placebo group and four in the tamoxifen citrate group

6 Three cases in the tamoxifen citrate group were fatal

7 All but three cases in each group required hospitalization

8 Based on women without cataracts at baseline (6,230-Placebo, 6,199-tamoxifen citrate)

9 All women (6,707-placebo, 6,681-tamoxifen citrate)

10 Updated long-term follow-up data, (median 6.9 years) from NSABP P-1 study added after cut-off for the other information in this table.

Table 4 describes the characteristics of the breast cancers in the NSABP P-1 trial and includes tumor size, nodal status, ER status. Tamoxifen citrate decreased the incidence of small estrogen receptor positive tumors, but did not alter the incidence of estrogen receptor negative tumors or larger tumors.

**Table 4**  
**Characteristics of Breast Cancer in NSABP P-1 Trial**

<b>Staging Parameter</b>	<b>Placebo N=156</b>	<b>Tamoxifen N=86</b>	<b>Total N=242</b>
<b>Tumor size</b>			
T1	117	60	177
T2	28	20	48
T3	7	3	10
T4	1	2	3
Unknown	3	1	4
<b>Nodal status</b>			
Negative	103	56	159
1-3 positive nodes	29	14	43
≤4 positive nodes	10	12	22
Unknown	14	4	18
<b>Stage</b>			
I	88	47	135
II: node negative	15	9	24
II: node positive	33	22	55
III	6	4	10
IV	21	1	3
Unknown	12	3	15
<b>Estrogen receptor</b>			
Positive	115	38	153
Negative	27	36	63
Unknown	14	12	26

<sup>1</sup> One participant presented with a suspicious bone scan but did not have documented metastases. She subsequently died of metastatic breast cancer.

Interim results from 2 trials in addition to the NSABP P-1 trial examining the effects of tamoxifen in reducing breast cancer incidence have been reported.

The first was the Italian Tamoxifen Prevention trial. In this trial women between the ages of 35 and 70, who had had a total hysterectomy, were randomized to receive 20 mg tamoxifen or matching placebo for 5 years. The primary endpoints were occurrence of, and death from, invasive breast cancer. Women without any specific risk factors for breast cancer were to be entered. Between 1992 and 1997, 5,408 women were randomized. Hormone Replacement Therapy (HRT) was used in 14% of participants. The trial closed in 1997 due to the large number of dropouts during the first year of treatment (26%). After 46 months of follow-up there were 22 breast cancers in women on placebo and 19 in women on tamoxifen. Although no decrease in breast cancer incidence was observed, there was a trend for a reduction in breast cancer among women receiving protocol therapy for at least 1 year (19-placebo, 11-tamoxifen). The small numbers of participants along with the low level of risk in this otherwise healthy group precluded an adequate assessment of the effect of tamoxifen in reducing the incidence of breast cancer.

The second trial, the Royal Marsden Trial (RMT) was reported as an interim analysis. The RMT was begun in 1986 as a feasibility study of whether larger scale trials could be mounted. The trial was subsequently extended to a pilot trial to accrue additional participants to further assess the safety of tamoxifen. Twenty-four hundred and seventy-one women were entered between 1986 and 1996; they were selected on the basis of a family history of breast cancer. HRT was used in 40% of participants. In this trial, with a 70-month median follow-up, 34 and 36 breast cancers (8 noninvasive, 4 on each arm) were observed among women on tamoxifen and placebo, respectively. Patients in this trial were younger than those in the NSABP P-1 trial and may have been more likely to develop ER (-) tumors, which are unlikely to be reduced in number by tamoxifen therapy. Although women were selected on the basis of family history and were thought to have a high risk of breast cancer, few events occurred, reducing the statistical power of the study. These factors are potential reasons why

the RMT may not have provided an adequate assessment of the effectiveness of tamoxifen in reducing the incidence of breast cancer.

In these trials, an increased number of cases of deep vein thrombosis, pulmonary embolus, stroke, and endometrial cancer were observed on the tamoxifen arm compared to the placebo arm. The frequency of events was consistent with the safety data observed in the NSABP P-1 trial.

## **INDICATIONS AND USAGE**

### **Metastatic Breast Cancer**

Tamoxifen citrate is effective in the treatment of metastatic breast cancer in women and men. In premenopausal women with metastatic breast cancer, tamoxifen citrate is an alternative to oophorectomy or ovarian irradiation. Available evidence indicates that patients whose tumors are estrogen receptor positive are more likely to benefit from tamoxifen citrate therapy.

### **Adjuvant Treatment of Breast Cancer**

Tamoxifen citrate is indicated for the treatment of node-positive breast cancer in postmenopausal women following total mastectomy or segmental mastectomy, axillary dissection, and breast irradiation. In some tamoxifen citrate adjuvant studies, most of the benefit to date has been in the subgroup with four or more positive axillary nodes.

Tamoxifen citrate is indicated for the treatment of axillary node-negative breast cancer in women following total mastectomy or segmental mastectomy, axillary dissection, and breast irradiation.

The estrogen and progesterone receptor values may help to predict whether adjuvant tamoxifen citrate therapy is likely to be beneficial.

Tamoxifen citrate reduces the occurrence of contralateral breast cancer in patients receiving adjuvant tamoxifen citrate therapy for breast cancer.

### **Ductal Carcinoma in Situ (DCIS)**

In women with DCIS, following breast surgery and radiation, tamoxifen citrate is indicated to reduce the risk of invasive breast cancer (see **BOXED WARNING** at the beginning of the label). The decision regarding therapy with tamoxifen for the reduction in breast cancer incidence should be based upon an individual assessment of the benefits and risks of tamoxifen therapy.

Current data from clinical trials support five years of adjuvant tamoxifen citrate therapy for patients with breast cancer.

### **Reduction in Breast Cancer Incidence in High Risk Women**

Tamoxifen citrate is indicated to reduce the incidence of breast cancer in women at high risk for breast cancer. This effect was shown in a study of 5 years planned duration with a median follow-up of 4.2 years. Twenty-five percent of the participants received drug for 5 years. The longer term effects are not known. In this study, there was no impact of tamoxifen on overall or breast cancer-related mortality (see **BOXED WARNING** at the beginning of the label).

Tamoxifen citrate is indicated only for high-risk women. "High risk" is defined as women at least 35 years of age with a 5-year predicted risk of breast cancer  $\geq 1.67\%$ , as calculated by the Gail Model.

Examples of combinations of factors predicting a 5-year risk  $\geq 1.67\%$  are:

**Age 35 or older and any of the following combination of factors:**

- One first degree relative with a history of breast cancer, 2 or more benign biopsies, and a history of a breast biopsy showing atypical hyperplasia; or
- At least 2 first degree relatives with a history of breast cancer, and a personal history of at least one breast biopsy; or
- LCIS

**Age 40 or older and any of the following combination of factors:**

- One first degree relative with a history of breast cancer, 2 or more benign biopsies, age at first live birth 25 or older, and age at menarche 11 or younger; or
- At least 2 first degree relatives with a history of breast cancer, and age at first live birth 19 or younger; or
- One first degree relative with a history of breast cancer, and a personal history of a breast biopsy showing atypical hyperplasia.

**Age 45 or older and any of the following combination of factors:**

- At least 2 first degree relatives with a history of breast cancer and age at first live birth 24 or younger; or
- One first degree relative with a history of breast cancer with a personal history of a benign breast biopsy, age at menarche 11 or less and age at first live birth 20 or more.

**Age 50 or older and any of the following combination of factors:**

- At least 2 first degree relatives with a history of breast cancer; or
- History of one breast biopsy showing atypical hyperplasia, and age at first live birth 30 or older and age at menarche 11 or less; or
- History of at least two breast biopsies with a history of atypical hyperplasia, and age at first live birth 30 or more.

**Age 55 or older and any of the following combination of factors:**

- One first degree relative with a history of breast cancer with a personal history of a benign breast biopsy, and age at menarche 11 or less; or
- History of at least 2 breast biopsies with a history of atypical hyperplasia, and age at first live birth 20 or older.

**Age 60 or older and:**

- 5-year predicted risk of breast cancer  $\geq 1.67\%$ , as calculated by the Gail Model.

For women whose risk factors are not described in the above examples, the Gail Model is necessary to estimate absolute breast cancer risk. Health Care Professionals can obtain a Gail Model Risk Assessment Tool by dialing 1-800-284-2480.

There are no data available regarding the effect of tamoxifen citrate on breast cancer incidence in women with inherited mutations (BRCA1, BRCA2).

After an assessment of the risk of developing breast cancer, the decision regarding therapy with tamoxifen citrate for the reduction in breast cancer incidence should be based upon an individual assessment of the benefits and risks of tamoxifen citrate therapy. In the NSABP P-1 trial, tamoxifen citrate treatment lowered the risk of developing breast cancer during

the follow-up period of the trial, but did not eliminate breast cancer risk (See Table 3 in **CLINICAL PHARMACOLOGY**).

## **CONTRAINDICATIONS**

Tamoxifen citrate tablets are contraindicated in patients with known hypersensitivity to the drug or any of its ingredients.

### **Reduction in Breast Cancer Incidence in High Risk Women and Women with DCIS**

Tamoxifen is contraindicated in women who require concomitant coumarin-type anticoagulant therapy or in women with a history of deep vein thrombosis or pulmonary embolus.

## **WARNINGS**

### **Effects in Metastatic Breast Cancer Patients**

As with other additive hormonal therapy (estrogens and androgens), hypercalcemia has been reported in some breast cancer patients with bone metastases within a few weeks of starting treatment with tamoxifen. If hypercalcemia does occur, appropriate measures should be taken and, if severe, tamoxifen should be discontinued.

### **Effects on the Uterus-Endometrial Cancer and Uterine Sarcoma**

An increased incidence of uterine malignancies has been reported in association with tamoxifen treatment. The underlying mechanism is unknown, but may be related to the estrogen-like effect of tamoxifen. Most uterine malignancies seen in association with tamoxifen are classified as adenocarcinoma of the endometrium. However, rare uterine sarcomas, including malignant mixed mullerian tumors, have also been reported. Uterine sarcoma is generally associated with a higher FIGO stage (III/IV) at diagnosis, poorer prognosis, and shorter survival. Uterine sarcoma has been reported to occur more frequently among long-term users ( $\geq 2$  years) of tamoxifen than non-users. Some of the uterine malignancies (endometrial carcinoma or uterine sarcoma) have been fatal.

In the NSABP P-1 trial, among participants randomized to tamoxifen there was a statistically significant increase in the incidence of endometrial cancer (33 cases of invasive endometrial cancer, compared to 14 cases among participants randomized to placebo (RR=2.48, 95% CI: 1.27-4.92). The 33 cases in participants receiving tamoxifen were FIGO Stage 1, including 20 IA, 12 1B, and 1 IC endometrial adenocarcinomas. In participants randomized to placebo, 13 were FIGO Stage I (8 IA and 5 IB) and 1 was FIGO Stage IV. Five women on tamoxifen and 1 on placebo received postoperative radiation therapy in addition to surgery. This increase was primarily observed among women at least 50 years of age at the time of randomization (26 cases of invasive endometrial cancer, compared to 6 cases among participants randomized to placebo (RR=4.50, 95% CI: 1.78-13.16). Among women  $\leq 49$  years of age at the time of randomization there were 7 cases of invasive endometrial cancer, compared to 8 cases among participants randomized to placebo (RR=0.94, 95% CI: 0.28-2.89). If age at the time of diagnosis is considered, there were 4 cases of endometrial cancer among participants  $\leq 49$  randomized to tamoxifen compared to 2 among participants randomized to placebo (RR=2.21, 95% CI: 0.4-12.0). For women  $\geq 50$  at the time of diagnosis, there were 29 cases among participants randomized to tamoxifen compared to 12 among women on placebo (RR=2.5, 95% CI: 1.3-4.9). The risk ratios were similar in the two groups, although fewer events occurred in younger women. Most (29 of 33 cases in the tamoxifen group) endometrial cancers were diagnosed in symptomatic women, although 5 of 33 cases in the tamoxifen group occurred in

asymptomatic women. Among women receiving tamoxifen the events appeared between 1 and 61 months (average=32 months) from the start of treatment.

In an updated review of long-term data (median length of total follow-up is 6.9 years, including blinded follow-up) on 8,306 women with an intact uterus at randomization in the NSABP P-1 risk reduction trial, the incidence of both adenocarcinomas and rare uterine sarcomas was increased in women taking tamoxifen. During blinded follow-up, there were 36 cases of FIGO Stage I endometrial adenocarcinoma (22 were FIGO Stage IA, 13 IB, and 1 IC) in women receiving tamoxifen and 15 cases in women receiving placebo [14 were FIGO Stage I (9 IA and 5 IB), and 1 case was FIGO Stage IV]. Of the patients receiving tamoxifen who developed endometrial cancer, one with Stage IA and 4 with Stage IB cancers received radiation therapy. In the placebo group, one patient with FIGO Stage 1B cancer received radiation therapy and the patient with FIGO Stage IVB cancer received chemotherapy and hormonal therapy. During total follow-up, endometrial adenocarcinoma was reported in 53 women randomized to tamoxifen (30 cases of FIGO Stage IA, 20 were Stage IB, 1 was Stage IC, and 2 were Stage IIIC), and 17 women randomized to placebo (9 cases were FIGO Stage IA, 6 were Stage IB, 1 was Stage IIIC, and 1 was Stage IVB) (incidence per 1,000 women-years of 2.20 and 0.71, respectively). Some patients received post-operative radiation therapy in addition to surgery. Uterine sarcomas were reported in 4 women randomized to tamoxifen (1 was FIGO IA, 1 was FIGO IB, 1 was FIGO IIA, and 1 was FIGO IIIC) and one patient randomized to placebo (FIGO IA); incidence per 1,000 women-years of 0.17 and 0.04, respectively. Of the patients randomized to tamoxifen, the FIGO IA and IB cases were a MMT and sarcoma, respectively; the FIGO II was a MMT; and the FIGO III was a sarcoma; and the one patient randomized to placebo had a MMT. A similar increased incidence in endometrial adenocarcinoma and uterine sarcoma was observed among women receiving tamoxifen in five other NSABP clinical trials.

Any patient receiving or who has previously received tamoxifen who reports abnormal vaginal bleeding should be promptly evaluated. Patients receiving or who have previously received tamoxifen should have annual gynecological examinations and they should promptly inform their physicians if they experience any abnormal gynecological symptoms, e.g., menstrual irregularities, abnormal vaginal bleeding, changes in vaginal discharge, or pelvic pain or pressure.

In the P-1 trial, endometrial sampling did not alter the endometrial cancer detection rate compared to women who did not undergo endometrial sampling (0.6% with sampling, 0.5% without sampling) for women with an intact uterus. There are no data to suggest that routine endometrial sampling in asymptomatic women taking tamoxifen to reduce the incidence of breast cancer would be beneficial.

#### **Non-Malignant Effects on the Uterus**

An increased incidence of endometrial changes including hyperplasia and polyps have been reported in association with tamoxifen treatment. The incidence and pattern of this increase suggest that the underlying mechanism is related to the estrogenic properties of tamoxifen.

There have been a few reports of endometriosis and uterine fibroids in women receiving tamoxifen. The underlying mechanism may be due to the partial estrogenic effect of tamoxifen. Ovarian cysts have also been observed in a small number of premenopausal patients with advanced breast cancer who have been treated with tamoxifen.

Tamoxifen has been reported to cause menstrual irregularity or amenorrhea.

### **Thromboembolic Effects of Tamoxifen**

There is evidence of an increased incidence of thromboembolic events, including deep vein thrombosis and pulmonary embolism, during tamoxifen therapy. When tamoxifen is coadministered with chemotherapy, there may be a further increase in the incidence of thromboembolic effects. For treatment of breast cancer, the risks and benefits of tamoxifen should be carefully considered in women with a history of thromboembolic events.

Data from the NSABP P-1 trial show that participants receiving tamoxifen without a history of pulmonary emboli (PE) had a statistically significant increase in pulmonary emboli (18-tamoxifen, 6-placebo, RR=3.01, 95% CI: 1.15-9.27). Three of the pulmonary emboli, all in the tamoxifen arm, were fatal. Eighty-seven percent of the cases of pulmonary embolism occurred in women at least 50 years of age at randomization. Among women receiving tamoxifen, the events appeared between 2 and 60 months (average = 27 months) from the start of treatment.

In this same population, a non-statistically significant increase in deep vein thrombosis (DVT) was seen in the tamoxifen group (30-tamoxifen, 19-placebo; RR=1.59, 95% CI: 0.86-2.98). The same increase in relative risk was seen in women  $\leq 49$  and in women  $\geq 50$ , although fewer events occurred in younger women. Women with thromboembolic events were at risk for a second related event (7 out of 25 women on placebo, 5 out of 48 women on tamoxifen) and were at risk for complications of the event and its treatment (0/25 on placebo, 4/48 on tamoxifen). Among women receiving tamoxifen, deep vein thrombosis events occurred between 2 and 57 months (average=19 months) from the start of treatment.

There was a non-statistically significant increase in stroke among patients randomized to tamoxifen (24-placebo; 34-tamoxifen; RR=1.42; 95% CI 0.82-2.51). Six of the 24 strokes in the placebo group were considered hemorrhagic in origin and 10 of the 34 strokes in the tamoxifen group were categorized as hemorrhagic. Seventeen of the 34 strokes in the tamoxifen group were considered occlusive and 7 were considered to be of unknown etiology. Fourteen of the 24 strokes on the placebo arm were reported to be occlusive and 4 of unknown etiology. Among these strokes 3 strokes in the placebo group and 4 strokes in the tamoxifen group were fatal. Eighty-eight percent of the strokes occurred in women at least 50 years of age at the time of randomization. Among women receiving tamoxifen, the events occurred between 1 and 63 months (average = 30 months) from the start of treatment.

### **Effects on the Liver: Liver Cancer:**

In the Swedish trial using adjuvant tamoxifen 40 mg/day for 2-5 years, 3 cases of liver cancer have been reported in the tamoxifen treated group vs. 1 case in the observation group (See **PRECAUTIONS,-Carcinogenesis**). In other clinical trials evaluating tamoxifen, no cases of liver cancer have been reported to date.

One case of liver cancer was reported in NSABP P-1 in a participant randomized to tamoxifen.

### **Effects on the Liver: Non-Malignant Effects**

Tamoxifen has been associated with changes in liver enzyme levels, and on rare occasions, a spectrum of more severe liver abnormalities including fatty liver, cholestasis, hepatitis and hepatic necrosis. A few of these serious cases included fatalities. In most reported cases

the relationship to tamoxifen is uncertain. However, some positive rechallenges and dechallenges have been reported.

In the NSABP P-1 trial, few grade 3-4 changes in liver function (SGOT, SGPT, bilirubin, alkaline phosphatase) were observed (10 on placebo and 6 on tamoxifen). Serum lipids were not systematically collected.

### **Other Cancers**

A number of second primary tumors, occurring at sites other than the endometrium, have been reported following the treatment of breast cancer with tamoxifen in clinical trials. Data from the NSABP B-14 and P-1 studies show no increase in other (non-uterine) cancers among patients receiving tamoxifen. Whether an increased risk for other (non-uterine) cancers is associated with tamoxifen is still uncertain and continues to be evaluated.

### **Effects on the Eye**

Ocular disturbances, including corneal changes, decrement in color vision perception, retinal vein thrombosis, and retinopathy have been reported in patients receiving tamoxifen. An increased incidence of cataracts and the need for cataract surgery have been reported in patients receiving tamoxifen.

In the NSABP P-1 trial, an increased risk of borderline significance of developing cataracts among those women without cataracts at baseline (540-tamoxifen; 483-placebo; RR=1.13, 95% CI: 1.00-1.28) was observed. Among these same women, tamoxifen was associated with an increased risk of having cataract surgery (101-tamoxifen; 63-placebo; RR=1.62, 95% CI 1.18-2.22) (See Table 3 in **CLINICAL PHARMACOLOGY**). Among all women on the trial (with or without cataracts at baseline), tamoxifen was associated with an increased risk of having cataract surgery (201- tamoxifen; 129-placebo; RR=1.58, 95% CI 1.26-1.97). Eye examinations were not required during the study. No other conclusions regarding non-cataract ophthalmic events can be made.

### **Pregnancy Category D**

Tamoxifen may cause fetal harm when administered to a pregnant woman. Women should be advised not to become pregnant while taking tamoxifen or within 2 months of discontinuing tamoxifen and should use barrier or nonhormonal contraceptive measures if sexually active. Tamoxifen does not cause infertility, even in the presence of menstrual irregularity. Effects on reproductive functions are expected from the antiestrogenic properties of the drug. In reproductive studies in rats at dose levels equal to or below the human dose, nonteratogenic developmental skeletal changes were seen and were found reversible. In addition, in fertility studies in rats and in teratology studies in rabbits using doses at or below those used in humans, a lower incidence of embryo implantation and a higher incidence of fetal death or retarded in utero growth were observed, with slower learning behavior in some rat pups when compared to historical controls. Several pregnant marmosets were dosed with 10 mg/kg/day (about 2-fold the daily maximum recommended human dose on a mg/m<sup>2</sup> basis) during organogenesis or in the last half of pregnancy. No deformations were seen and, although the dose was high enough to terminate pregnancy in some animals, those that did maintain pregnancy showed no evidence of teratogenic malformations.

In rodent models of fetal reproductive tract development, tamoxifen (at doses 0.002 to 2.4-fold the daily maximum recommended human dose on a mg/m<sup>2</sup> basis) caused changes in both sexes that are similar to those caused by estradiol, ethynylestradiol and

diethylstilbestrol. Although the clinical relevance of these changes is unknown, some of these changes, especially vaginal adenosis, are similar to those seen in young women who were exposed to diethylstilbestrol in utero and who have a 1 in 1,000 risk of developing clear-cell adenocarcinoma of the vagina or cervix. To date, in utero exposure to tamoxifen has not been shown to cause vaginal adenosis, or clear-cell adenocarcinoma of the vagina or cervix, in young women. However, only a small number of young women have been exposed to tamoxifen in utero, and a smaller number have been followed long enough (to age 15-20) to determine whether vaginal or cervical neoplasia could occur as a result of this exposure.

There are no adequate and well-controlled trials of tamoxifen in pregnant women. There have been a small number of reports of vaginal bleeding, spontaneous abortions, birth defects, and fetal deaths in pregnant women. If this drug is used during pregnancy, or the patient becomes pregnant while taking this drug, or within approximately two months after discontinuing therapy, the patient should be apprised of the potential risks to the fetus including the potential long-term risk of a DES-like syndrome.

#### **Reduction in Breast Cancer Incidence in High Risk Women - Pregnancy Category D**

For sexually active women of child-bearing potential, tamoxifen therapy should be initiated during menstruation. In women with menstrual irregularity, a negative  $\beta$ -HCG immediately prior to the initiation of therapy is sufficient (See **PRECAUTIONS Information for Patients - Reduction in Breast Cancer Incidence in High Risk Women**).

### **PRECAUTIONS**

#### **General**

Decreases in platelet counts, usually to 50,000-100,000/mm<sup>3</sup>, infrequently lower, have been occasionally reported in patients taking tamoxifen for breast cancer. In patients with significant thrombocytopenia, rare hemorrhagic episodes have occurred, but it is uncertain if these episodes are due to tamoxifen therapy. Leukopenia has been observed, sometimes in association with anemia and/or thrombocytopenia. There have been rare reports of neutropenia and pancytopenia in patients receiving tamoxifen; this can sometimes be severe.

In the NSABP P-1 trial, 6 women on tamoxifen and 2 on placebo experienced grade 3-4 drops in platelet count ( $\leq 50,000/\text{mm}^3$ ).

#### **Information for Patients**

Patients should be instructed to read the Medication Guide supplied as required by law when tamoxifen citrate is dispensed. The complete text of the Medication Guide is reprinted at the end of this document.

#### **Reduction in Invasive Breast Cancer and DCIS in Women with DCIS**

Women with DCIS treated with lumpectomy and radiation therapy who are considering tamoxifen to reduce the incidence of a second breast cancer event should assess the risks and benefits of therapy, since treatment with tamoxifen decreased the incidence of invasive breast cancer, but has not been shown to affect survival (See Table 1 in **CLINICAL PHARMACOLOGY**).

#### **Reduction in Breast Cancer Incidence in High Risk Women**

Women who are at high risk for breast cancer can consider taking tamoxifen therapy to reduce the incidence of breast cancer. Whether the benefits of treatment are considered to outweigh the risks depends on a woman's personal health history and on how she weighs the

benefits and risks. Tamoxifen therapy to reduce the incidence of breast cancer may therefore not be appropriate for all women at high risk for breast cancer. Women who are considering tamoxifen therapy should consult their health care professional for an assessment of the potential benefits and risks prior to starting therapy for reduction in breast cancer incidence (See **Table 3** in **CLINICAL PHARMACOLOGY**). Women should understand that tamoxifen reduces the incidence of breast cancer, but may not eliminate risk. Tamoxifen decreased the incidence of small estrogen receptor positive tumors, but did not alter the incidence of estrogen receptor negative tumors or larger tumors. In women with breast cancer who are at high risk of developing a second breast cancer, treatment with about 5 years of tamoxifen reduced the annual incidence rate of a second breast cancer by approximately 50%.

Women who are pregnant or who plan to become pregnant should not take tamoxifen to reduce her risk of breast cancer. Effective nonhormonal contraception must be used by all premenopausal women taking tamoxifen and for approximately two months after discontinuing therapy if they are sexually active. Tamoxifen does not cause infertility, even in the presence of menstrual irregularity. For sexually active women of child-bearing potential, tamoxifen therapy should be initiated during menstruation. In women with menstrual irregularity, a negative  $\beta$ -HCG immediately prior to the initiation of therapy is sufficient (See **WARNINGS-Pregnancy Category D**).

Two European trials of tamoxifen to reduce the risk of breast cancer were conducted and showed no difference in the number of breast cancer cases between the tamoxifen and placebo arms. These studies had trial designs that differed from that of NSABP P-1, were smaller than NSABP P-1, and enrolled women at a lower risk for breast cancer than those in P-1.

### **Monitoring During Tamoxifen Therapy**

Women taking or having previously taken tamoxifen should be instructed to seek prompt medical attention for new breast lumps, vaginal bleeding, gynecologic symptoms (menstrual irregularities, changes in vaginal discharge, or pelvic pain or pressure), symptoms of leg swelling or tenderness, unexplained shortness of breath, or changes in vision. Women should inform all care providers, regardless of the reason for evaluation, that they take tamoxifen. Women taking tamoxifen to reduce the incidence of breast cancer should have a breast examination, a mammogram, and a gynecologic examination prior to the initiation of therapy. These studies should be repeated at regular intervals while on therapy, in keeping with good medical practice. Women taking tamoxifen as adjuvant breast cancer therapy should follow the same monitoring procedures as for women taking tamoxifen for the reduction in the incidence of breast cancer. Women taking tamoxifen as treatment for metastatic breast cancer should review this monitoring plan with their care provider and select the appropriate modalities and schedule of evaluation.

### **Laboratory Tests**

Periodic complete blood counts, including platelet counts, and periodic liver function tests should be obtained.

### **Drug Interactions**

When tamoxifen is used in combination with coumarin-type anticoagulants, a significant increase in anticoagulant effect may occur. Where such coadministration exists, careful monitoring of the patient's prothrombin time is recommended.

In the NSABP P-1 trial, women who required coumarin-type anticoagulants for any reason were ineligible for participation in the trial (See **CONTRAINDICATIONS**).

There is an increased risk of thromboembolic events occurring when cytotoxic agents are used in combination with tamoxifen.

Tamoxifen reduced letrozole plasma concentrations by 37%. The effect of tamoxifen on metabolism and excretion of other antineoplastic drugs, such as cyclophosphamide and other drugs that require mixed function oxidases for activation, is not known. Tamoxifen and N-desmethyl tamoxifen plasma concentrations have been shown to be reduced when coadministered with rifampin or aminoglutethimide. Induction of CYP3A4-mediated metabolism is considered to be the mechanism by which these reductions occur; other CYP3A4 inducing agents have not been studied to confirm this effect.

One patient receiving tamoxifen with concomitant phenobarbital exhibited a steady state serum level of tamoxifen lower than that observed for other patients (i.e., 26 ng/mL vs. mean value of 122 ng/mL). However, the clinical significance of this finding is not known. Rifampin induced the metabolism of tamoxifen and significantly reduced the plasma concentrations of tamoxifen in 10 patients. Aminoglutethimide reduces tamoxifen and N-desmethyl tamoxifen plasma concentrations. Medroxyprogesterone reduces plasma concentrations of N-desmethyl, but not tamoxifen.

Concomitant bromocriptine therapy has been shown to elevate serum tamoxifen and N-desmethyl tamoxifen.

#### **Drug/Laboratory Testing Interactions**

During postmarketing surveillance, T4 elevations were reported for a few postmenopausal patients which may be explained by increases in thyroid-binding globulin. These elevations were not accompanied by clinical hyperthyroidism.

Variations in the karyopyknotic index on vaginal smears and various degrees of estrogen effect on Pap smears have been infrequently seen in postmenopausal patients given tamoxifen.

In the postmarketing experience with tamoxifen, infrequent cases of hyperlipidemias have been reported. Periodic monitoring of plasma triglycerides and cholesterol may be indicated in patients with pre-existing hyperlipidemias (See **ADVERSE REACTIONS-Postmarketing experience** section).

#### **Carcinogenesis**

A conventional carcinogenesis study in rats at doses of 5, 20, and 35 mg/kg/day (about one, three and seven-fold the daily maximum recommended human dose on a mg/m<sup>2</sup> basis) administered by oral gavage for up to 2 years revealed a significant increase in hepatocellular carcinoma at all doses. The incidence of these tumors was significantly greater among rats administered 20 or 35 mg/kg/day (69%) compared to those administered 5 mg/kg/day (14%). In a separate study, rats were administered tamoxifen at 45 mg/kg/day (about nine-fold the daily maximum recommended human dose on a mg/m<sup>2</sup> basis); hepatocellular neoplasia was exhibited at 3 to 6 months.

Granulosa cell ovarian tumors and interstitial cell testicular tumors were observed in two separate mouse studies. The mice were administered the trans and racemic forms of

tamoxifen for 13 to 15 months at doses of 5, 20 and 50 mg/kg/day (about one-half, two and five-fold the daily recommended human dose on a mg/m<sup>2</sup> basis).

### **Mutagenesis**

No genotoxic potential was found in a conventional battery of *in vivo* and *in vitro* tests with pro- and eukaryotic test systems with drug metabolizing systems. However, increased levels of DNA adducts were observed by <sup>32</sup>P post-labeling in DNA from rat liver and cultured human lymphocytes. Tamoxifen also has been found to increase levels of micronucleus formation *in vitro* in human lymphoblastoid cell line (MCL-5). Based on these findings, tamoxifen is genotoxic in rodent and human MCL-5 cells.

### **Impairment of Fertility**

Tamoxifen produced impairment of fertility and conception in female rats at doses of 0.04 mg/kg/day (about 0.01-fold the daily maximum recommended human dose on a mg/m<sup>2</sup> basis) when dosed for two weeks prior to mating through day 7 of pregnancy. At this dose, fertility and reproductive indices were markedly reduced with total fetal mortality. Fetal mortality was also increased at doses of 0.16 mg/kg/day (about 0.03-fold the daily maximum recommended human dose on a mg/m<sup>2</sup> basis) when female rats were dosed from days 7-17 of pregnancy. Tamoxifen produced abortion, premature delivery and fetal death in rabbits administered doses equal to or greater than 0.125 mg/kg/day (about 0.05-fold the daily maximum recommended human dose on a mg/m<sup>2</sup> basis). There were no teratogenic changes in either rats or rabbits.

**Pregnancy Category D: See WARNINGS**

### **Nursing Mothers**

Tamoxifen has been reported to inhibit lactation. Two placebo-controlled studies in over 150 women have shown that tamoxifen significantly inhibits early postpartum milk production. In both studies tamoxifen was administered within 24 hours of delivery for between 5 and 18 days. The effect of tamoxifen on established milk production is not known.

There are no data that address whether tamoxifen is excreted into human milk. If excreted, there are no data regarding the effects of tamoxifen in breast milk on the breastfed infant or breastfed animals. However, direct neonatal exposure of tamoxifen to mice and rats (not via breast milk) produced 1) reproductive tract lesions in female rodents (similar to those seen in humans after intrauterine exposure to diethylstilbestrol) and 2) functional defects of the reproductive tract in male rodents such as testicular atrophy and arrest of spermatogenesis.

It is not known if tamoxifen is excreted in human milk. Because of the potential for serious adverse reactions in nursing infants from tamoxifen, women taking tamoxifen should not breast feed.

### **Reduction in Breast Cancer Incidence in High Risk Women and Women with DCIS:**

It is not known if tamoxifen is excreted in human milk. Because of the potential for serious adverse reactions in nursing infants from tamoxifen, women taking tamoxifen should not breast feed.

### **Pediatric Use**

The use of SOLTAMOX™ in pediatric patients has not been evaluated.

### **Geriatric Use**

In the NSABP P-1 trial, the percentage of women at least 65 years of age was 16%. Women at least 70 years of age accounted for 6% of the participants. A reduction in breast cancer incidence was seen among participants in each of the subsets: A total of 28 and 10 invasive breast cancers were seen among participants 65 and older in the placebo and tamoxifen groups, respectively. Across all other outcomes, the results in this subset reflect the results observed in the subset of women at least 50 years of age. No overall differences in tolerability were observed between older and younger patients (See **CLINICAL PHARMACOLOGY - Clinical Studies - Reduction in Breast Cancer Incidence in High Risk Women** section).

In the NSABP B-24 trial, the percentage of women at least 65 years of age was 23%. Women at least 70 years of age accounted for 10% of participants. A total of 14 and 12 invasive breast cancers were seen among participants 65 and older in the placebo and tamoxifen groups, respectively. This subset is too small to reach any conclusions on efficacy. Across all other endpoints, the results in this subset were comparable to those of younger women enrolled in this trial. No overall differences in tolerability were observed between older and younger patients.

### **ADVERSE REACTIONS**

Adverse reactions to tamoxifen are relatively mild and rarely severe enough to require discontinuation of treatment in breast cancer patients.

Continued clinical studies have resulted in further information which better indicates the incidence of adverse reactions with tamoxifen as compared to placebo.

In one single-dose pharmacokinetic study in healthy perimenopausal and postmenopausal female volunteers, throat irritation was reported by 3 of 60 evaluable subjects (5.0%) in the Soltamox™ treatment groups while none of the subjects in the tamoxifen reference group reported this event. All events were mild and occurred within an hour after dosing. All events were resolved within 24 hours.

### **Metastatic Breast Cancer**

Increased bone and tumor pain and, also, local disease flare have occurred, which are sometimes associated with a good tumor response. Patients with increased bone pain may require additional analgesics. Patients with soft tissue disease may have sudden increases in the size of preexisting lesions, sometimes associated with marked erythema within and surrounding the lesions and/or the development of new lesions. When they occur, the bone pain or disease flare are seen shortly after starting tamoxifen and generally subside rapidly.

In patients treated with tamoxifen for metastatic breast cancer, the most frequent adverse reaction to tamoxifen is hot flashes.

Other adverse reactions which are seen infrequently are hypercalcemia, peripheral edema, distaste for food, pruritus vulvae, depression, dizziness, light-headedness, headache, hair thinning and/or partial hair loss, and vaginal dryness.

### **Premenopausal Women**

The following table summarizes the incidence of adverse reactions reported at a frequency of 2% or greater from clinical trials (Ingle, Pritchard, Buchanan) which compared tamoxifen therapy to ovarian ablation in premenopausal patients with metastatic breast cancer.

	TAMOXIFEN All Effects	OVARIAN ABLATION All Effects
Adverse Reactions*	% of Women n =104	% of Women n = 100
Flush	33	46
Amenorrhea	16	69
Altered Menses	13	5
Oligomenorrhea	9	1
Bone Pain	6	6
Menstrual Disorder	6	4
Nausea	5	4
Cough/Coughing	4	1
Edema	4	1
Fatigue	4	1
Musculoskeletal Pain	3	0
Pain	3	4
Ovarian Cyst(s)	3	2
Depression	2	2
Abdominal Cramps	1	2
Anorexia	1	2

\*Some women had more than one adverse reaction.

### Male Breast Cancer

Tamoxifen is well tolerated in males with breast cancer. Reports from the literature and case reports suggest that the safety profile of tamoxifen in males is similar to that seen in women. Loss of libido and impotence have resulted in discontinuation of tamoxifen therapy in male patients. Also, in oligospermic males treated with tamoxifen, LH, FSH, testosterone and estrogen levels were elevated. No significant clinical changes were reported.

### Adjuvant Breast Cancer

In the NSABP B-14 study, women with axillary node-negative breast cancer were randomized to 5 years of tamoxifen 20 mg/day or placebo following primary surgery. The reported adverse effects are tabulated below (mean follow-up of approximately 6.8 years) showing adverse events more common on tamoxifen than on placebo. The incidence of hot flashes (64% vs. 48%), vaginal discharge (30% vs. 15%), and irregular menses (25% vs. 19%) were higher with tamoxifen compared with placebo. All other adverse effects occurred with similar frequency in the 2 treatment groups, with the exception of thrombotic events; a higher incidence was seen in tamoxifen-treated patients (through 5 years, 1.7% vs. 0.4%). Two of the patients treated with tamoxifen who had thrombotic events died.

NSABP B-14 Study		
	TAMOXIFEN (n=1422)	Placebo (n=1437)
	% of Women	
Hot Flashes	64	48
Fluid Retention	32	30
Vaginal Discharge	30	15
Nausea	26	24
Irregular Menses	25	19
Weight Loss (>5%)	23	18
Skin Changes	19	15
Increased SGOT	5	3
Increased Bilirubin	2	1
Increased Creatinine	2	1
Thrombocytopenia*	2	1
Thrombotic Events		
Deep Vein Thrombosis	0.8	0.2
Pulmonary Embolism	0.5	0.2
Superficial Phlebitis	0.4	0.0

\*Defined as a platelet count of <100,000/mm<sup>3</sup>

In the Eastern Cooperative Oncology Group (ECOG) adjuvant breast cancer trial, tamoxifen or placebo was administered for 2 years to women following mastectomy. When compared to placebo, tamoxifen showed a significantly higher incidence of hot flashes (19% vs. 8% for placebo). The incidence of all other adverse reactions was similar in the 2 treatment groups with the exception of thrombocytopenia where the incidence for tamoxifen was 10% vs. 3% for placebo, an observation of borderline statistical significance.

In other adjuvant studies, Toronto and tamoxifen Adjuvant Trial Organization (NATO), women received either tamoxifen or no therapy. In the Toronto study, hot flashes were observed in 29% of patients for tamoxifen vs. 1% in the untreated group. In the NATO trial, hot flashes and vaginal bleeding were reported in 2.8% and 2.0% of women, respectively, for tamoxifen vs. 0.2% for each in the untreated group.

### **Ductal Carcinoma in Situ (DCIS)**

The type and frequency of adverse events in the NSABP B-24 trial were consistent with those observed in the other adjuvant trials conducted with tamoxifen.

### **Reduction in Breast Cancer Incidence in High Risk Women**

In the NSABP P-1 Trial, there was an increase in five serious adverse effects in the tamoxifen group: endometrial cancer (33 cases in the tamoxifen group vs. 14 in the placebo group); pulmonary embolism (18 cases in the tamoxifen group vs. 6 in the placebo group); deep vein thrombosis (30 cases in the tamoxifen group vs. 19 in the placebo group); stroke (34 cases in the tamoxifen group vs. 24 in the placebo group); cataract formation (540 cases in the tamoxifen group vs. 483 in the placebo group) and cataract surgery (101 cases in the tamoxifen group vs. 63 in the placebo group) (See **WARNINGS** and Table 3 in **CLINICAL PHARMACOLOGY**).

The following table presents the adverse events observed in NSABP P-1 by treatment arm. Only adverse events more common on tamoxifen than placebo are shown.

**NSABP P-1 Trial: All Adverse Events**

	TAMOXIFEN (n=6681)	% of Women	Placebo (n=6707)
<b><u>Self Reported Symptoms</u></b>	N=6441 <sup>1</sup>		N=6469 <sup>1</sup>
Hot Flashes	80		68
Vaginal Discharges	55		35
Vaginal Bleeding	23		22
<b><u>Laboratory Abnormalities</u></b>	N=6520 <sup>2</sup>		N=6535 <sup>2</sup>
Platelets decreased	0.7		0.3
<b><u>Adverse Effects</u></b>	N=6492 <sup>3</sup>		N=6484 <sup>3</sup>
<b>Other Toxicities</b>			
Mood	11.6		10.8
Infection/Sepsis	6.0		5.1
Constipation	4.4		3.2
Alopecia	5.2		4.4
Skin	5.6		4.7
Allergy	2.5		2.1

<sup>1</sup>Number with Quality of Life Questionnaires

<sup>2</sup>Number with Treatment Follow-up Forms

<sup>3</sup>Number with Adverse Drug Reaction Forms

In the NSABP P-1 trial, 15.0% and 9.7% of participants receiving tamoxifen and placebo therapy, respectively, withdrew from the trial for medical reasons. The following are the medical reasons for withdrawing from tamoxifen and placebo therapy, respectively: Hot flashes (3.1% vs. 1.5%) and Vaginal Discharge (0.5% vs. 0.1%).

In the NSABP P-1 trial, 8.7% and 9.6% of participants receiving tamoxifen and placebo therapy, respectively, withdrew for non-medical reasons.

On the NSABP P-1 trial, hot flashes of any severity occurred in 68% of women on placebo and in 80% of women on tamoxifen. Severe hot flashes occurred in 28% of women on placebo and 45% of women on tamoxifen. Vaginal discharge occurred in 35% and 55% of women on placebo and tamoxifen respectively; and was severe in 4.5% and 12.3% respectively. There was no difference in the incidence of vaginal bleeding between treatment arms.

**Postmarketing Experience**

Less frequently reported adverse reactions are vaginal bleeding, vaginal discharge, menstrual irregularities, skin rash and headaches. Usually these have not been of sufficient severity to require dosage reduction or discontinuation of treatment. Very rare reports of erythema multiforme, Stevens-Johnson syndrome, bullous pemphigoid, interstitial pneumonitis, and rare reports of hypersensitivity reactions including angioedema have been reported with tamoxifen therapy. In some of these cases, the time to onset was more than one year. Rarely, elevation of serum triglyceride levels, in some cases with pancreatitis, may be associated with the use of tamoxifen (see **PRECAUTIONS, Drug/Laboratory Testing Interactions** section).

**OVERDOSAGE**

Signs observed at the highest doses following studies to determine LD<sub>50</sub> in animals were respiratory difficulties and convulsions.

Acute overdosage in humans has not been reported. In a study of advanced metastatic cancer patients which specifically determined the maximum tolerated dose of tamoxifen in evaluating the use of very high doses to reverse multidrug resistance, acute neurotoxicity

manifested by tremor, hyperreflexia, unsteady gait and dizziness were noted. These symptoms occurred within 3-5 days of beginning tamoxifen and cleared within 2-5 days after stopping therapy. No permanent neurologic toxicity was noted. One patient experienced a seizure several days after tamoxifen was discontinued and neurotoxic symptoms had resolved. The causal relationship of the seizure to tamoxifen therapy is unknown. Doses given in these patients were all greater than 400 mg/m<sup>2</sup> loading dose, followed by maintenance doses of 150 mg/m<sup>2</sup> of tamoxifen given twice a day.

In the same study, prolongation of the QT interval on the electrocardiogram was noted when patients were given doses higher than 250 mg/m<sup>2</sup> loading dose, followed by maintenance doses of 80 mg/m<sup>2</sup> of tamoxifen given twice a day. For a woman with a body surface area of 1.5 m<sup>2</sup> the minimal loading dose and maintenance doses given at which neurological symptoms and QT changes occurred were at least 6 fold higher in respect to the maximum recommended dose.

No specific treatment for overdosage is known; treatment must be symptomatic.

### **DOSAGE AND ADMINISTRATION**

For patients with breast cancer, the recommended daily dose is 20-40 mg. Dosages greater than 20 mg per day should be given in divided doses (morning and evening). A 20 mg dose of SOLTAMOX<sup>TM</sup> is administered as 10 mL (equivalent to 2 teaspoons) of the oral solution.

In three single agent adjuvant studies in women, one 10 mg tamoxifen citrate tablet was administered two (ECOG and NATO) or three (Toronto) times a day for two years. In the NSABP B-14 adjuvant study in women with node-negative breast cancer, one 10 mg tamoxifen citrate tablet was given twice a day for at least 5 years. Results of the B-14 study suggest that continuation of therapy beyond five years does not provide additional benefit (see **CLINICAL PHARMACOLOGY**). In the EBCTCG 1995 overview, the reduction in recurrence and mortality was greater in those studies that used tamoxifen for about 5 years than in those that used tamoxifen for a shorter period of therapy. There was no indication that doses greater than 20 mg per day were more effective. Current data from clinical trials support 5 years of adjuvant tamoxifen citrate therapy for patients with breast cancer.

### **Ductal Carcinoma in Situ (DCIS)**

The recommended dose is tamoxifen citrate 20 mg daily for 5 years.

### **Reduction in Breast Cancer Incidence in High Risk Women**

The recommended dose is tamoxifen citrate 20 mg daily for 5 years. There are no data to support the use of tamoxifen citrate other than for 5 years (See **CLINICAL PHARMACOLOGY-Clinical Studies - Reduction in Breast Cancer Incidence in High Risk Women**).

**HOW SUPPLIED**

SOLTAMOX™ Oral Solution is a sugar-free, clear colorless liquid, with licorice and aniseed odor and taste. It is supplied in a 150mL bottle, each 5mL solution contains 15.2 mg tamoxifen citrate, equivalent to 10 mg tamoxifen. NDC (# *not available at this time*).

Do not store above 25° C (77° F). Store in the original package in order to protect from light. Use within 3 months of opening. Storage: DO NOT freeze or refrigerate.

**R<sub>x</sub> only**

Dated; 8/25/2005

Manufactured for  
Savient Pharmaceuticals, Inc. by:  
Rosemont Pharmaceuticals Ltd.,  
Yorkdale Industrial Park, Braithwaite Street  
Leeds, LS11 9XE, UK

Address medical inquiries to  
Savient Pharmaceuticals, Inc.  
One Tower Center  
Fourteenth Floor  
East Brunswick, NJ 08816  
866-692-6374

© 2004, Savient Pharmaceuticals, Inc.

Printed in USA

## MEDICATION GUIDE

SOLTAMOX™ Oral Solution  
(SOL-ta-mox)  
Generic name: tamoxifen citrate

*Written for women who use SOLTAMOX™ to lower their high chance of getting breast cancer or who have ductal carcinoma in situ (DCIS)*

This Medication Guide discusses only the use of SOLTAMOX™ to lower the chance of getting breast cancer in high-risk women and in women treated for DCIS.

People taking SOLTAMOX™ **to treat** breast cancer have different benefits and different decisions to make than high-risk women or women with ductal carcinoma in situ (DCIS) taking SOLTAMOX™ to reduce the chance of getting breast cancer. If you already have breast cancer, talk with your doctor about how the benefits of treating breast cancer with SOLTAMOX™ compare to the risks that are described in this document.

### **Why should I read this Medication Guide?**

This guide has information to help you decide whether to use SOLTAMOX™ to lower your chance of getting breast cancer.

**You and your doctor should talk about whether the possible benefit of SOLTAMOX™ in lowering your high chance of getting breast cancer is greater than its possible risks.** Your doctor has a special computer program or hand-held calculator to tell if you are in the high-risk group. If you have DCIS and have been treated with surgery and radiation therapy, your doctor may prescribe SOLTAMOX™ to decrease your chance of getting invasive (spreading) breast cancer.

Read this guide carefully before you start SOLTAMOX™. It is important to read the information you get each time you get more medicine. There may be something new. This guide does not tell you everything about SOLTAMOX™ and does **not** take the place of talking with your doctor. Only you and your doctor can determine if SOLTAMOX™ is right for you.

### **What is the most important information I should know about using SOLTAMOX™ to reduce the chance of getting breast cancer?**

SOLTAMOX™ (tamoxifen citrate) is a prescription medicine that is like estrogen (female hormone) in some ways and different in other ways. In the breast, tamoxifen can block estrogen's effects. Because it does this, tamoxifen may block the growth of breast cancers that need estrogen to grow (cancers that are estrogen- or progesterone-receptor positive).

Tamoxifen citrate can lower the chance of getting breast cancer in women with a higher than normal chance of getting breast cancer in the next five years (high-risk women) and women with DCIS. **Because high-risk women don't have cancer yet, it is important to think carefully about whether the possible benefit of tamoxifen in lowering the chance of getting breast cancer is greater than its possible risks.**

This Medication Guide reviews the risks and benefits of using SOLTAMOX™ to reduce the chance of getting breast cancer in high-risk women and women with DCIS. This guide does not discuss the special benefits and decisions for people who already have breast cancer.

### **Why do women and men use SOLTAMOX™?**

SOLTAMOX™ has more than one use. SOLTAMOX™ is used:

1. **to lower the chance** of getting breast cancer in women with a higher than normal chance of getting breast cancer in the next 5 years (high-risk women).
2. **to lower the chance** of getting invasive (spreading) breast cancer in women who had surgery and radiation for ductal carcinoma in situ (DCIS). DCIS means the cancer is only inside the milk ducts.
3. **to treat** breast cancer in women after they have finished early treatment. Early treatment can include surgery, radiation, and chemotherapy. Tamoxifen may keep the cancer from spreading to others parts of the body. It may also reduce the woman's chance of getting a new breast cancer.
4. in women and men, **to treat** breast cancer that has spread to other parts of the body (metastatic breast cancer).

This guide talks only about using SOLTAMOX™ to lower the chance of getting breast cancer (#1 and #2 above).

### **What are the benefits of SOLTAMOX™ to lower the chance of getting breast cancer in high-risk women and in women treated for DCIS?**

A large US study looked at **high-risk women** and compared the ones who took tamoxifen for 5 years with others who took a pill without tamoxifen (placebo). High-risk women were defined as women who have a 1.7% or greater chance of getting breast cancer in the next 5 years, based on a special computer program. In this study:

- Out of every 1,000 high-risk women **who took a placebo**, each year about 7 got breast cancer.
- Out of every 1,000 high-risk women **who took tamoxifen**, each year about 4 got breast cancer.

The study showed that on average, high-risk women who took tamoxifen lowered their chances of getting breast cancer by 44%, from 7 in 1,000 to 4 in 1,000.

Another US study looked at **women with DCIS** and compared those who took tamoxifen for 5 years with others who took a placebo. In this study:

- Out of every 1,000 women with DCIS **who took placebo**, each year about 17 got breast cancer.
- Out of every 1,000 women with DCIS **who took tamoxifen**, each year about 10 got breast cancer.

The study showed that on average, women with DCIS who took tamoxifen lowered their chances of getting invasive (spreading) breast cancer by 43%, from 17 in 1,000 to 10 in 1,000.

**These studies do not mean that taking SOLTAMOX™ will lower your personal chance of getting breast cancer.** We do not know what the benefits will be for any one woman who takes SOLTAMOX™ to reduce her chance of getting breast cancer.

### **What are the risks of SOLTAMOX™?**

In the studies described under "What are the benefits of SOLTAMOX™?", the high-risk women who took tamoxifen got certain side effects at a higher rate than those who took a placebo. **Some of these side effects can cause death.**

In one study, in women who still had their uterus

- Out of every 1,000 women who took a placebo, each year 1 got endometrial cancer (cancer of the lining of the uterus) and none got uterine sarcoma (cancer of the body of the uterus).
- Out of every 1,000 women who took tamoxifen, each year 2 got endometrial cancer and fewer than 1 got uterine sarcoma.

These results show that, on average, in high-risk women **who still had their uterus**, tamoxifen doubled the chance of getting endometrial cancer from 1 in 1,000 to 2 in 1,000, and it increased the chance of getting uterine sarcoma. **This does not mean that taking tamoxifen will double your personal chance of getting endometrial cancer or increase your chance of getting uterine sarcoma.** We do not know what this risk will be for any one woman. The risk is different for women who no longer have their uterus.

**For all women in this study, taking tamoxifen increased the risk of having a blood clot in their lungs or veins, or of having a stroke.** In some cases, women died from these effects.

**Tamoxifen increased the risk of getting cataracts (clouding of the lens of the eye) or needing cataract surgery.** (See "What are the possible side effects of SOLTAMOX™ for more details about side effects.)

### **What don't we know about taking SOLTAMOX™ to reduce the chance of getting breast cancer?**

We don't know

- if tamoxifen lowers the chance of getting breast cancer in women who have abnormal breast cancer genes (BRCA1 and BRCA2)
- if taking tamoxifen for 5 years reduces the number of breast cancers a woman will get in her lifetime or if it only delays some breast cancers
- if tamoxifen helps a woman live longer
- the effects of taking tamoxifen with hormone replacement therapy (HRT), birth control pills, or androgens (male hormones)
- the benefits of taking tamoxifen if you are less than 35 years old

Studies are being done to learn more about the long-term benefits and risks of using tamoxifen to reduce the chance of getting breast cancer.

## **What are the possible side effects of SOLTAMOX™?**

The most common side effect of tamoxifen is hot flashes. This is not a sign of a serious problem.

The next most common side effect is vaginal discharge. If the discharge is bloody, it could be a sign of a serious problem. [See "Changes in the lining (endometrium) or body of your uterus" below.]

Less common but serious side effects of tamoxifen are listed below. These can occur at any time. Call your doctor right away if you have any signs of side effects listed below:

- **Changes in the lining (endometrium) or body of your uterus.** These changes may mean serious problems are starting, including cancer of the uterus. The signs of changes in the uterus are:
  - Vaginal bleeding or bloody discharge that could be a rusty or brown color. You should call your doctor even if only a small amount of bleeding occurs.
  - Change in your monthly bleeding, such as in the amount or timing of bleeding or increased clotting.
  - Pain or pressure in your pelvis (below your belly button).
- **Blood clots in your veins or lungs.** These can cause serious problems, including death. You may get clots up to 2-3 months after you stop taking tamoxifen. The signs of blood clots are:
  - sudden chest pain, shortness of breath, coughing up blood
  - pain, tenderness, or swelling in one or both of your legs
- **Stroke.** Stroke can cause serious medical problems, including death. The signs of stroke are:
  - sudden weakness, tingling, or numbness in your face, arm or leg, especially on one side of your body
  - sudden confusion, trouble speaking or understanding
  - sudden trouble seeing in one or both eyes
  - sudden trouble walking, dizziness, loss of balance or coordination
  - sudden severe headache with no known cause
- **Cataracts or increased chance of needing cataract surgery.** The sign of these problems is slow blurring of your vision.
- **Liver problems, including jaundice.** The signs of liver problems include lack of appetite and yellowing of your skin or whites of your eyes.

Another less serious side effect that has been reported occasionally with SOLTAMOX™ is mild, temporary throat irritation. These are not all the possible side effects of tamoxifen. For a complete list, ask your doctor or pharmacist.

### **Who should not take SOLTAMOX™?**

#### **Do not take tamoxifen for any reason if you**

- **Are pregnant or plan to become pregnant while taking tamoxifen or during the 2 months after you stop taking tamoxifen. Tamoxifen may harm your unborn baby.** It takes about 2 months to clear tamoxifen from your body. To be sure you are not pregnant, you can start taking tamoxifen while you are having your menstrual period. Or, you can take a pregnancy test to be sure you are not pregnant before you begin.
- **Are breast-feeding.** We do not know if tamoxifen can pass through your milk and harm your baby.
- **Have had an allergic reaction to tamoxifen or SOLTAMOX™** (the other name for tamoxifen), or to any of its inactive ingredients.

**If you get pregnant while taking SOLTAMOX™, stop taking it right away and contact your doctor. SOLTAMOX™ may harm your unborn baby.**

#### **Do not take SOLTAMOX™ to lower your chance of getting breast cancer if**

- You ever had a blood clot that needed medical treatment.
- You are taking medicines to thin your blood, like warfarin, (also called Coumadin®\*).
- Your ability to move around is limited for most of your waking hours.
- You are at risk for blood clots. Your doctor can tell you if you are at high risk for blood clots.
- You do not have a higher than normal chance of getting breast cancer. Your doctor can tell you if you are a high-risk woman.

### **How should I take SOLTAMOX™?**

- Take 2 tsp SOLTAMOX™ once or twice a day, as directed by your physician. You can take SOLTAMOX™ with or without food. Take your medicine every day. It may be easier to remember if you take it at the same time each day.
- If you forget a dose, take it when you remember, then take the next dose as usual. If it is almost time for your next dose or you remember at your next dose, do not take more than 2 tsp at one time to make up a missed dose.
- Take SOLTAMOX™ for 5 years, unless your doctor tells you otherwise.

### **What should I avoid while taking SOLTAMOX™?**

- **Do not become pregnant while taking tamoxifen or for 2 months after you stop.** Tamoxifen can stop hormonal birth control methods from working. Hormonal methods include birth control pills, patches, injections, rings and implants. Therefore, while taking tamoxifen, **use birth control methods that**

**don't use hormones**, such as condoms, diaphragms with spermicide, or plain IUD's. If you get pregnant, stop taking tamoxifen right away and call your doctor.

- **Do not breast feed.** We do not know if tamoxifen can pass through your milk and if it can harm the baby.

#### **What should I do while taking SOLTAMOX™?**

- Have regular gynecology check-ups ("female exams"), breast exams and mammograms. Your doctor will tell you how often. These will check for signs of breast cancer and cancer of the endometrium (lining of the uterus). Because tamoxifen does not prevent all breast cancers, and you may get other types of cancers, you need these exams to find any cancers as early as possible.
- Because tamoxifen can cause serious side effects, pay close attention to your body. Signs you should look for are listed in "What are the possible side effects of SOLTAMOX™?"
- Tell all of the doctors that you see that you are taking SOLTAMOX™.
- Tell your doctor right away if you have any new breast lumps.

#### **General information about the safe and effective use of SOLTAMOX™**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Your doctor has prescribed SOLTAMOX™ only for you. Do not give it to other people, even if they have a similar condition, because it may harm them. Do not use it for a condition for which it was not prescribed.

This Medication Guide is a summary of information about SOLTAMOX™ for women who use SOLTAMOX™ to lower their high chance of getting breast cancer or who have DCIS. If you want more information about SOLTAMOX™, ask your doctor or pharmacist. They can give you information about SOLTAMOX™ that is written for health professionals. For more information about SOLTAMOX™ or breast cancer, please visit [www.savientpharma.com](http://www.savientpharma.com) or call 1-866-692-6374.

Ingredients: tamoxifen, ethanol, glycerol, propylene glycol, sorbitol solution, licorice flavor, aniseed flavor, purified water.

\*Coumadin® is a registered trademark of Bristol-Myers Squibb Pharmaceuticals.

Savient Pharmaceuticals, Inc., East Brunswick, NJ 08816 Rev xx-0x

Printed in USA ©200x Savient Pharmaceuticals, Inc.

This Medication Guide has been approved by the U.S. Food and Drug Administration

**SOLTAMOX™**

**TAMOXIFEN CITRATE**

**Oral Solution**

**(equivalent to)**

**10mg/5mL**

**Tamoxifen**

**New Drug Application [505(b)(2)]**

**Section 2**

**Labeling**

<b>Application Number:</b>	<b>21-807</b>
Pharmacologic category:	A Nonsteroidal Antiestrogen
Product Name:	Soltamox™
Indication:	Treatment of Breast Cancer
Document status:	Draft
Document date:	XX XX 2004
Number of pages:	XX

Property of Savient Pharmaceuticals, Inc.

**Confidential**

May not be divulged, published or otherwise disclosed  
to any other parties without the prior written permission of Savient Pharmaceuticals, Inc.

SOLTAMOX™

Tamoxifen Citrate Oral Solution Equivalent To 10 mg/5mL Tamoxifen

NDA 505(b)(2) #21-807

Section 2: Labeling

---

<p>Each 5mL dose of oral solution contains Tamoxifen Citrate USP 15.2mg (equivalent to 10mg tamoxifen). The product also contains ethanol (19% v/v), glycerol, propylene glycol, sorbitol solution, licorice and aniseed flavors and purified water.</p> <p><b>Storage:</b> Store at Controlled Room Temperature 15° – 25 °C (59 ° - 77° F). Store in original package in order to protect from light. Use within 3 months of opening.</p> <p><b>Pharmacist:</b> Dispense only in original container with calibrated cup.</p> <div data-bbox="499 938 690 987" style="border: 1px solid black; text-align: center; padding: 5px; margin: 10px auto; width: fit-content;">BARCODE</div>	<p><b>NDC#</b></p> <p><b>SOLTAMOX™</b> <b>TAMOXIFEN CITRATE</b> <b>Oral Solution</b> <b>(equivalent to)</b> <b>10mg/5mL</b> <b>Tamoxifen</b> <b>Rx Only</b> <b>Sugar Free</b> <b>150 mL</b> <b>PATIENT PACK</b></p> 	<p>KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.</p> <p><b>Dosage:</b> As prescribed by the physician. To be taken orally.</p> <p>See Package Insert for Complete Prescribing Information.</p> <p>Manufactured by: Rosemont Pharmaceuticals Ltd., Yorkdale Industrial Park, Braithwaite Street Leeds, LS11 9XE, UK</p> <p>Manufactured for: Savient Pharmaceuticals, Inc. East Brunswick, NJ 08816, USA</p> <p>LOT : MFG : EXP :</p>
--	--	--


**DRAFT CONTAINER LABEL**

SOLTAMOX™

Tamoxifen Citrate Oral Solution Equivalent To 10mg/5mL Tamoxifen

NDA 505(b)(2) #21-807

Section 2: Labeling

<p><b>Lot:</b> <b>Mfg:</b> <b>Exp:</b></p> <p>TAMOXIFEN CITRATE <b>Oral Solution</b> (equivalent to) <b>10mg/5mL</b> Tamoxifen</p> <p><b>Rx only</b></p> 			
<p><b>NDC#</b></p> <p><b>SOLTAMOX™</b> <b>TAMOXIFEN CITRATE</b> <b>Oral Solution</b> (equivalent to) <b>10mg/5 mL</b> <b>Tamoxifen</b></p> <p><b>Rx only</b> <b>150 mL</b> <b>Patient pack</b></p>  	<p>Each 5mL dose of oral solution contains Tamoxifen Citrate USP 15.2mg (equivalent to 10mg tamoxifen). The product also contains ethanol (19% v/v), glycerol, propylene glycol, sorbitol solution, licorice and aniseed flavors and purified water.</p> <p><b>Storage:</b> Store at Controlled Room Temperature 15°–25° C (59– 77° F). Store in the original package in order to protect from light.</p> <p>Use within 3 months of opening.</p> <p><b>Pharmacist:</b> Dispense only in original container with the calibrated cup.</p> <div data-bbox="751 1226 1012 1299" style="border: 1px solid black; text-align: center; padding: 5px;"><b>BAR CODE</b></div>	<p><b>NDC#</b></p> <p><b>SOLTAMOX™</b> <b>TAMOXIFEN CITRATE</b> <b>Oral Solution</b> (equivalent to) <b>10mg/5mL</b> <b>Tamoxifen</b></p> <p><b>Rx only</b> <b>150 mL</b> <b>Patient Pack</b></p>  	<p><b>KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.</b></p> <p><b>Dosage:</b> As prescribed by the physician. To be taken orally.</p> <p>See Package Insert for Complete Prescribing Information.</p> <p>Manufactured by: Rosemont Pharmaceuticals Ltd., Yorkdale Industrial Park, Braithwaite Street Leeds, LS11 9XE, UK</p> <p>Manufactured for: Savient Pharmaceuticals, Inc. East Brunswick, NJ 08816, USA</p>

Confidential Information Proprietary to Savient Pharmaceuticals, Inc.

**DRAFT CARTON LABEL**

SOLTAMOX™  
Tamoxifen Citrate Oral Solution Equivalent To 10 mg/5mL Tamoxifen  
NDA 505(b)(2) #21-807  
Section 2: Labeling

---