



PMS-186C

PMS-116C

PMS-301C

Black

PMS-300C

Sep. 05, 2002
 Revised by I.Horii/Fujisawa Japan

CrossTech-C45422-Proof 1
 Form M03-PS:C1
 Fujisawa-P.O. 21321-Job No. 62922
 3/10/05-jc-Q4.11

Fonts: Helvetica (A)



NDC 0469-3250-10

Mycamine™
(micafungin sodium)
for injection

50 mg/vial

50 mg X 10 Vials

325010



NDC 0469-3250-10

Mycamine™
(micafungin sodium)
for injection
For Intravenous Infusion Only

50 mg/vial

Once reconstituted with 5 mL of 0.9% sodium chloride injection
(without bacteriostatic agent), each mL contains 10 mg/mL.
Requires further dilution prior to infusion. See package insert.

Single Use Vial. Discard unused portion.

Rx Only

50 mg X 10 Vials

325010



Mycamine™
(micafungin sodium)
for injection

50 mg/vial

Made in Japan
Mid. for:
Fujisawa Healthcare, Inc.
Deerfield, IL 60015-2548

50 mg X 10 Vials



NDC 0469-3250-10

Mycamine™
(micafungin sodium)
for injection
For Intravenous Infusion Only

50 mg/vial


Sterile, Nonpyrogenic

Each vial contains: Micafungin sodium 50 mg;
lactose, NF 200 mg with citric acid and/or sodium
hydroxide for pH adjustment.

Store at 25°C (77°F) excursions 15°-30°C (59°-86°F).
[See USP Controlled Room Temperature] Protect from light.

50 mg X 10 Vials

325010



(01) 2 0304693 25010 5



Mycamine™
(micafungin sodium)
for injection

50 mg/vial

Preservative free. Discard unused portion.

50 mg X 10 Vials



PMS-186C PMS-116C PMS-301C PMS-300C Black


CrossTech-C45421-Proof 1
Form M03-PS:C1
Fujisawa-P.O. 21321-Job No. 62923
3/10/05-jc-Q4.11

Fonts: Helvetica (A)

(b) (4)

CrossTech
COMMUNICATIONS INC

312 382-0111 FAX 382-0004





PMS-186C

PMS-116C

PMS-301C

PMS-300C

PMS-WHITE
BACKGROUND

Black



CrossTech-B45420-Proof A7
 Form M01-PS:B1
 Fujisawa-P.O. 21321-Job No. 40927
 2/24/05-hc-Q4.11

Fonts: Helvetica (A)
 Typesmiths Pi Font

