

In case of overdose, get medical help or contact a Poison Control Center right away.

Retain outer carton for full product uses, directions and warnings. Discard POPPAC after use.

Tamper Evident Feature: Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken.

Distributed By:

GlaxoSmithKline Consumer Healthcare, L.P.
Moon Township, PA 15108

Made in the U.S.A.

©2006 GlaxoSmithKline

00000XX

28 LOZENGES



To open vial, push in child resistant band on the POPPAC™ with thumb.



Flip up the top of the POPPAC™ and remove lozenge.

A small amount of powder on opening of the POPPAC™ is normal.

NL 20 POPPAC™

Commit

nicotine polacrilex lozenge, 2mg
STOP SMOKING AID

MINT FLAVOR

Directions:

Not for Individual Sale

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours


Place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge. Occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 20 - 30 minutes). Do not eat or drink 15 minutes before using or while the lozenge is in your mouth.

Do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.

Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash.




33



FOR MORE INFORMATION, PLEASE VISIT www.quit.com (it's just so easy to quit)

VARNISH KO

LOT EXP



FROM THE MARKETERS OF
Nicorette

Commit[®]
nicotine polacrilex lozenge, 2mg
STOP SMOKING AID

Includes Usage Guide

2 mg


112 LOZENGES
in 4 Containers of 28
2mg EACH

VARNISH KO

Glue Flap - NO COPY OR VARNISH


VARNISH KO

VARNISH KO



0000000000

PLACE ANTI-THEFT STICKER HERE



EAS TAGGED
0 0000-0000-00 0

VARNISH KO

Drug Facts (continued)

• If you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 4	Weeks 5 to 8	Weeks 9 to 12
1 lozenge every 1 to 3 hours	1 lozenge every 3 to 4 hours	1 lozenge every 4 to 8 hours

• Nicotine lozenge is a medicine and must be used a certain way to get the best results.

• Place the lozenge in your mouth and use the lozenge to slowly dissolve (about 20–30 minutes). Minimize swallowing. Do not chew or swallow the lozenge.

• You may feel a warm or tingling sensation.

• Occasionally, move the lozenge from one side of your mouth to the other until completely dissolved (about 20–30 minutes).

• Do not eat or drink 15 minutes before using or while the lozenge is in your mouth.

• To improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks.

• Do not use more than one lozenge at a time or continuously. Use one lozenge after another since this may cause jaw pain, heartburn, nausea or other side effects.

• Do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.

• Stop using the nicotine lozenge at the end of 12 weeks. If you still feel the need to use nicotine lozenges, talk to your doctor.

Other information

- each lozenge contains sodium, 15mg
- Phenylephrine: Contains Phenylephrine 0.4 mg per lozenge
- dose of 0.25–0.75, 0.5–1.75
- protein from light

Inactive ingredients saccharin, aspartame, calcium polycrystalline, corn syrup solids, beeswax, butane, magnesium stearate, indigo carmine, menthyl rubinate, menthyl sodium alginate, sodium carbonate, soy protein, triethyl citrate, xanthan gum

Questions or comments?
Call 1-800-QUIT-NOW (1-800-488-7463) (10am–6pm EST weekdays)

COMMITTED QUITTERS POPPAC™ and associated logo designs and marks made these designs are trademarks owned and/or licensed to GlaxoSmithKline or its affiliated companies.


Distributed by:
GlaxoSmithKline Consumer Products, L.P.
Morris Township, PA 15109, Made in the U.S.A.
©2005 GlaxoSmithKline 003860X

Must be safe to those under 18 years of age
Proof of age required
Must be safe in vending machines or from any source where proof of age cannot be verified

TAMPER EVIDENT FEATURE: Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken. Retain outer carton for full product uses, directions and warnings.

TO INCREASE YOUR SUCCESS IN QUITTING:

- You must be motivated to quit.
- Use Enough – Use at least 9 lozenges of Commit per day during the first six weeks.
- Use Long Enough – Use Commit for the full 12 weeks.
- Use With a Support Program as directed in the enclosed User's Guide.



Commit POPPAC™

To open, twist, push in (do not rotate) band on the POPPAC™ with thumb.

Flip up the top of the POPPAC™ and remove lozenge.

A small amount of powder on opening of the POPPAC™ is normal

Drug Facts

Active ingredient (in each lozenge)	Purpose
Nicotine polacrilex, 2mg	Stop smoking aid

Use – used as withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings

• If you are pregnant or breastfeeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your baby. Try to stop smoking without using any nicotine replacement product. This medicine is believed to be safer than smoking. However, for each hour smoked from this medicine are not 100% hours.

Do not use

- if you continue to smoke, chew tobacco, use snuff, or use a nicotine patch or other nicotine-containing products.

Ask a doctor before use if you have

- a sudden-onset chest pain
- heart disease, recent heart attack, or irregular heartbeat. Medicine can increase your heart rate.
- high blood pressure not controlled with medication. Medicine can increase your blood pressure.
- stomach pain or diarrhea

Ask a doctor or pharmacist before use if you are

- using a tricyclic antidepressant or other smoking drug
- taking prescription medication for depression or anxiety. Your prescription dose may need to be adjusted.


Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat

Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. If you are pregnant, get medical help or contact a Poison Control Center right away.

Directions

- If you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- stop smoking completely when you begin using the lozenge
- if you smoke your first cigarette within 30 minutes of waking up, use 2mg nicotine lozenge.



FREE DELIVERED TO YOUR HOME

FROM THE MARKETERS OF **Nicorette**

Commit
nicotine polacrilex lozenge, 2mg
STOP SMOKING AID

Includes User's Guide

MINT FLAVOR

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP. If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Commit 4mg Lozenge.

2 mg

168 LOZENGES (6 POPPAC Containers of 28)

2mg EACH

Drug Facts (continued)

- If you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 1	Weeks 2 to 5	Weeks 6 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- nicotine lozenge is a medicine and must be used in a certain way to get the best results
- Place the lozenge in your mouth and allow the lozenge to slowly dissolve about 15-30 minutes. Minimize swallowing. Do not chew or swallow lozenge.
- You may feel a warm or tingling sensation.
- Occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 15-30 minutes).
- Do not eat or drink 15 minutes before using or while the lozenge is in your mouth.
- To improve your chances of quitting, use at least 16 lozenges per day for the first 6 weeks.
- Do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you nausea, heartburn, nausea or other side effects.
- Do not use more than 16 lozenges in 4 hours. Do not use more than 16 lozenges per day.
- Stop using the nicotine lozenge at the end of 12 weeks. If you still feel the need to use nicotine lozenges, talk to your doctor.

Other information

- each lozenge contains sodium, 10mg
- Pharmaceuticals: Contains Phenylephrine 5mg per lozenge
- store at 20°-25°C (68°-77°F)
- protect from light

Inactive ingredients include: aspartame, sodium polycrystalline, corn syrup solids, sucrose, butane, magnesium stearate, mallic acid, menthyl potassium benzoate, sodium alginate, sodium carbonate, soy protein, trimethylolpropane, xanthan gum

Questions or comments?
Call 1-800-368-6973 (1-800-NICOTINE) 10:00am - 4:00pm EST weekdays

COMMIT, COMMITTED TO QUITTING, POPPAC™, and associated logo designs and overall theme design are trademarks owned and/or licensed to Ciba-Geigy/Novartis or its affiliated companies.

Ciba-Geigy/Novartis Consumer Healthcare, LLC
Novartis Township, PA 15088. Made in the USA.
©2008 Ciba-Geigy/Novartis 00000000

Not for sale to those under 18 years of age


No proof of age required

Alert for side in vending machines or from any source where proof of age cannot be verified

TAMPER EVIDENT FEATURE: Do not use if clear neckband printed "SEALED FOR YOUR PROTECTION" is missing or broken. Retain outer carton for full product uses, directions and warnings.


TO INCREASE YOUR SUCCESS IN QUITTING:

- You must be motivated to quit.
- Use Enough** - Use at least 16 lozenges of Commit per day during the first six weeks.
- Use Long Enough** - Use Commit for the full 12 weeks.
- Use With a Support Program** as directed in the enclosed User's Guide.



POP
PAC™

To open the unit, push in child resistant band on the POPPAC™ with thumb.



POP
PAC™

To pop the top of the POPPAC™ and remove lozenge.

A small amount of powder on opening of the POPPAC™ is normal

VARNISH KO

EXP LOT

VARNISH KO

VARNISH KO

0000000000

EAS TAGGED

0 0000-0000-00 0

PLACE ANTI-THEFT STICKER HERE

THEFT SURVEILLANCE TAG AREA

GLUE FLAP - NO COPY OR VARNISH

Drug Facts

Active Ingredient (in each lozenge) Nicotine polacrilex, 2mg

Purpose Stop smoking aid

Use - relieve withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breastfeeding, only use this medicine on the advice of your health care provider. Smoking can harm your baby. Try to stop smoking without using any nicotine replacement medicine. This medicine is not used to be used then smoking. However, the risks to your child from this medicine are not fully known.

Do not use

- If you continue to smoke, chew tobacco, use snuff, or use a nicotine patch or other nicotine containing products

Ask a doctor before use if you have

- a sudden increase in blood pressure
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, tremor, weakness, and fainting

Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in a paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- If you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- stop smoking completely when you begin using the lozenge
- If you smoke your first cigarette within 30 minutes of waking up, use 4mg nicotine lozenge.

In case of overdose, get medical help or contact a Poison Control Center right away.

Retain outer carton for full product uses, directions and warnings. Discard POPPAC after use.

Tamper Evident Feature: Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken.

Distributed By:
GlaxoSmithKline Consumer Healthcare, L.P.
Moon Township, PA 15108

Made in the U.S.A.
©2006 GlaxoSmithKline
00000XX

28 LOZENGES



To open vial, push in child resistant band on the POPPAC™ with thumb.



Flip up the top of the POPPAC™ and remove lozenge.

A small amount of powder on opening of the POPPAC™ is normal.



MINT FLAVOR

Directions:

Not for Individual Sale

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

Place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge. Occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 20 - 30 minutes). Do not eat or drink 15 minutes before using or while the lozenge is in your mouth.

Do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.

Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. ↓



34

FOR MORE INFORMATION, PLEASE VISIT www.committedquitters.com (© 2006 GlaxoSmithKline)


FREE RECIPIENT-SPECIFIC NICHOTINE PATCHES
FOR COMMITTED QUITTERS™

VARNISH KO

LOT

EXP

FROM THE MARKETERS OF
Nicorette®



Commit®
nicotine polacrilex lozenge, 4mg
STOP SMOKING AID

Includes User's Guide

MINT FLAVOR


4 mg

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.
If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Commit® 2mg Lozenge.

112 LOZENGES
(4 Poppac® Containers of 28)
4mg EACH

VARNISH KO

VARNISH KO




0 000000000

PLACE ANTI-THEFT STICKER HERE

THFT SUPPLIANCE TAG AREA

EAS TAGGED



0 0000-0000-00 0

VARNISH KO

VARNISH KO

not for sale to those under 18 years of age


not for sale in vending machines or from any source where proof of age cannot be verified

TAMPER EVIDENT FEATURE: Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken.

Retain outer carton for full product uses, directions and warnings.


TO INCREASE YOUR SUCCESS IN QUITTING:

- You must be motivated to quit.
- Use Enough** - Use at least 8 lozenges of Commit® per day during the first six weeks.
- Use Long Enough** - Use Commit® for the full 12 weeks.
- Use With a Support Program** as directed in the enclosed User's Guide.



Commit® POPPAC™

To open vial, push in child resistant band on the POPPAC™ with thumb.



Flip up the top of the POPPAC™ and remove lozenge.

A small amount of powder on opening of the POPPAC™ is normal

VARNISH KO

VARNISH KO

Drug Facts (continued)

It is not clear how long it takes to get the full benefit of the lozenge. It may take up to 12 weeks to see the full benefit.

Weeks 1 to 6	Weeks 7 to 8	Weeks 9 to 12
1 lozenge every 1 to 2 hours	2 lozenges every 2 to 3 hours	1 lozenge every 3 to 4 hours

- nicotine lozenge is a medicine and must be used a certain way to get the best results
- place the lozenge in your mouth and allow the lozenge to slowly dissolve (about 20-30 minutes). Minimize swallowing. Do not chew or swallow the lozenge.
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 10-15 minutes)
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth to improve your chances of quitting, use at least 8 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you dizziness, headache, nausea or other side effects
- do not use more than 8 lozenges in 6 hours. Do not use more than 28 lozenges per day.
- when using the nicotine lozenge at the end of 12 weeks, if you still need the need to use nicotine lozenges, talk to your doctor.

Other information

- each lozenge contains sodium, 15mg
- Phenylephrine HCl; Contains Phenylephrine 0.4 mg per lozenge
- store at 20°-25°C (68°-77°F)
- protect from light

Inactive ingredients sucralose, sorbitol, calcium polycrystalline, gum arabic, sodium benzoate, magnesium stearate, hydroxypropyl methylcellulose, potassium bicarbonate, sodium alginate, sodium carbonate, zinc stearate, hydroxyethylcellulose, croscarmellose sodium

Questions or comments?
call toll free 1-800-4-A-SMOKER (1-800-427-7662) (9:30am-4:30pm EST weekdays)

COMMITTED QUITTERS™ POPPAC™ and associated logo designs and overall trade dress designs are trademarks owned and/or licensed to GlaxoSmithKline or its affiliated companies.

Distributed by:
GlaxoSmithKline Consumer Healthcare, L.P.
Blow, Pennsylvania, PA 19106, Made in the U.S.A. 000000X

Drug Facts

Active ingredient (in each lozenge) Nicotine polacrilex, 4mg **Purpose**
Stop smoking aid

Use - reduce withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breastfeeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking, however, the risks to your child from this medicine are not fully known.

Do not use

- if you continue to smoke, chew tobacco, use snuff, or use a nicotine patch or other nicotine-containing products

Ask a doctor before use if you have

- a sudden-onset chest pain
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes

Ask a doctor or pharmacist before use if you are

- using a non-medicated stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Do not use and ask a doctor if


- mouth problems occur
- persistent halitosis or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, dizziness, weakness and rapid heartbeat

Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using the product, read the enclosed User's Guide for complete directions and other important information
- stop smoking completely when you begin using the lozenge
- if you smoke your first cigarette more than 30 minutes after waking up, use 2mg nicotine lozenge

GLUE FLAP - NO COPY OR VARNISH



VARNISH KO

LOT

EXP

VARNISH KO

VARNISH KO

Drug Facts (continued)

• You should stop your cigarette within 30 minutes of waking up, use 1mg nicotine lozenge according to the following 12 week schedule:

Week 1 to 1	Week 2 to 2	Week 3 to 12
1 lozenge every 1-2 hours	1 lozenge every 2-3 hours	1 lozenge every 3-4 hours

• **Warnings:** Before the medicine will work, you must smoke to get the full benefit.

• **Directions:** Place 1 lozenge in your mouth and chew the lozenge to break it into small pieces (about 20-30 minutes). Alternate chewing lozenges. Do not chew or swallow lozenges. You may feel a warm or tingling sensation.

• **Other information:** Place the lozenge 1/2 to one side of your mouth in the other side of your mouth. Do not use more than 1 lozenge at a time. Do not use more than 20 lozenges per day.

• **Do not use more than one lozenge at a time and do not use more than one lozenge after another since this may cause jaw soreness, tooth or gum pain or other side effects. Do not use more than 18 lozenges in 4 hours. Do not use more than 20 lozenges per day.**

• **Child-Resistant Packaging:** At the end of 12 weeks, you will find the need to use lozenges less often during your lozenge.

Other information:

- each lozenge contains nicotine, 4mg
- **Warnings:** Contains Nicotine. Nicotine is a drug. It can be addictive. It can also be harmful. It can cause dizziness, upset stomach, loss of appetite, dry mouth, blurred vision, headache, nervousness, heart palpitations, dizziness, rash, allergic reactions, numbness and tingling, and other side effects.
- **Directions:** See directions on the label.

Contains: COMMITTED TO HELPING SMOKERS QUIT. FREE. (To learn how, visit www.quitnow.com)

Other information:

- **Warnings:** Contains Nicotine. Nicotine is a drug. It can be addictive. It can also be harmful. It can cause dizziness, upset stomach, loss of appetite, dry mouth, blurred vision, headache, nervousness, heart palpitations, dizziness, rash, allergic reactions, numbness and tingling, and other side effects.
- **Directions:** See directions on the label.

Contains: COMMITTED TO HELPING SMOKERS QUIT. FREE. (To learn how, visit www.quitnow.com)

Other information:

- **Warnings:** Contains Nicotine. Nicotine is a drug. It can be addictive. It can also be harmful. It can cause dizziness, upset stomach, loss of appetite, dry mouth, blurred vision, headache, nervousness, heart palpitations, dizziness, rash, allergic reactions, numbness and tingling, and other side effects.
- **Directions:** See directions on the label.

Contains: COMMITTED TO HELPING SMOKERS QUIT. FREE. (To learn how, visit www.quitnow.com)

Other information:


- **Warnings:** Contains Nicotine. Nicotine is a drug. It can be addictive. It can also be harmful. It can cause dizziness, upset stomach, loss of appetite, dry mouth, blurred vision, headache, nervousness, heart palpitations, dizziness, rash, allergic reactions, numbness and tingling, and other side effects.
- **Directions:** See directions on the label.

Contains: COMMITTED TO HELPING SMOKERS QUIT. FREE. (To learn how, visit www.quitnow.com)

Other information:

- **Warnings:** Contains Nicotine. Nicotine is a drug. It can be addictive. It can also be harmful. It can cause dizziness, upset stomach, loss of appetite, dry mouth, blurred vision, headache, nervousness, heart palpitations, dizziness, rash, allergic reactions, numbness and tingling, and other side effects.
- **Directions:** See directions on the label.

FROM THE MANUFACTURER OF
Nicorette®



Commit
nicotine polacrilex lozenge, 4mg
STOP SMOKING AID

Includes User's Guide
MINT FLAVOR

4 mg

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.
If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Commit® 2mg Lozenge.

168 LOZENGES
(18 POPPAC® COUNTERS FOR 28) 4mg EACH

Not for sale to those under 18 years of age.
Not for sale to those under 18 years of age.
Not for sale to those under 18 years of age.


Do not use if your medicine pointed "SEALED FOR SAFETY" is missing or broken.

Retain your carton for 180 days. Directions and warnings.

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use Enough - Use at least 3 lozenges of Commit® per day during the first six weeks.
3. Use Long Enough - Use Commit® for the full 12 weeks.
4. Use With a Support Program as directed in the enclosed User's Guide.

Commit® POPPAC™



In separate pouch in child-resistant carton on the POPPAC™ with lozenge.

Fill up the top of the POPPAC™ and remove lozenge.

A small amount of powder on opening of the POPPAC™ is normal

Drug Facts

Active ingredient (in each lozenge)
Nicotine polacrilex, 4mg

Warnings

• If you are pregnant or breastfeeding, only use this medicine as the advice of your health care provider. Smoking can harm your unborn child. Try to quit smoking without using any nicotine replacement products. Your medicine is labeled to be used after your smoking. However, the data for use after 12 weeks are not known.

Do not use

- if you continue to smoke, chew tobacco, use snuff, or use a mouth pouch or other nicotine-containing product.
- if you have had a heart attack, chest pain, or other heart condition.
- if you have had a stroke, or if you have had a seizure.
- if you have had a recent heart attack, or if you have had a heart condition.
- if you have had a recent stroke, or if you have had a seizure.
- if you have had a recent heart attack, or if you have had a heart condition.
- if you have had a recent stroke, or if you have had a seizure.

Do not use

- if you have had a heart attack, chest pain, or other heart condition.
- if you have had a stroke, or if you have had a seizure.
- if you have had a recent heart attack, or if you have had a heart condition.
- if you have had a recent stroke, or if you have had a seizure.

Do not use

- if you have had a heart attack, chest pain, or other heart condition.
- if you have had a stroke, or if you have had a seizure.
- if you have had a recent heart attack, or if you have had a heart condition.
- if you have had a recent stroke, or if you have had a seizure.

Do not use

- if you have had a heart attack, chest pain, or other heart condition.
- if you have had a stroke, or if you have had a seizure.
- if you have had a recent heart attack, or if you have had a heart condition.
- if you have had a recent stroke, or if you have had a seizure.

Do not use

- if you have had a heart attack, chest pain, or other heart condition.
- if you have had a stroke, or if you have had a seizure.
- if you have had a recent heart attack, or if you have had a heart condition.
- if you have had a recent stroke, or if you have had a seizure.

Do not use

- if you have had a heart attack, chest pain, or other heart condition.
- if you have had a stroke, or if you have had a seizure.
- if you have had a recent heart attack, or if you have had a heart condition.
- if you have had a recent stroke, or if you have had a seizure.

Do not use

- if you have had a heart attack, chest pain, or other heart condition.
- if you have had a stroke, or if you have had a seizure.
- if you have had a recent heart attack, or if you have had a heart condition.
- if you have had a recent stroke, or if you have had a seizure.

Do not use

- if you have had a heart attack, chest pain, or other heart condition.
- if you have had a stroke, or if you have had a seizure.
- if you have had a recent heart attack, or if you have had a heart condition.
- if you have had a recent stroke, or if you have had a seizure.

Do not use

- if you have had a heart attack, chest pain, or other heart condition.
- if you have had a stroke, or if you have had a seizure.
- if you have had a recent heart attack, or if you have had a heart condition.
- if you have had a recent stroke, or if you have had a seizure.

Do not use

- if you have had a heart attack, chest pain, or other heart condition.
- if you have had a stroke, or if you have had a seizure.
- if you have had a recent heart attack, or if you have had a heart condition.
- if you have had a recent stroke, or if you have had a seizure.


Do not use

- if you have had a heart attack, chest pain, or other heart condition.
- if you have had a stroke, or if you have had a seizure.
- if you have had a recent heart attack, or if you have had a heart condition.
- if you have had a recent stroke, or if you have had a seizure.

VARNISH KO


VARNISH KO

VARNISH KO




0000000000

PLACE ANTI-THEFT STICKER HERE



0 0000-0000-00 0

EAS TAGGED



0 0000-0000-00 0

VARNISH KO

VARNISH KO

VARNISH KO

GLUE FLAP - NO COPY OR VARNISH