

NDC 010888-0000-00

See new warnings information.

Migraine Relief

Ibuprofen Capsules,
200 mg

Pain Reliever (NSAID)

40 Softgels*
*liquid filled capsules

Active ingredient (in each capsule)

Ibuprofen 200 mg (NSAID)
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Use

• treats migraine

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people

- who are allergic to aspirin. Symptoms may include: hives, facial swelling, skin raddening, asthma (wheezing), blisters
- facial swelling
- rash
- asthma (wheezing)
- blisters

If allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:

- are age 60 or older.
- have had stomach ulcers or bleeding problems
- take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

Do not use if you have ever had an allergic reaction to any other pain reliever (fever reducer

or night anesthetic) or after heart surgery.

Ask a doctor before use if you have: • never had migraines diagnosed by a health

care professional • been using usual migraine medicine • had a headache

of your life • fever and stiff neck • headaches beginning after or caused by head injury

• motion, coughing or bending • experienced your first headache after the age of 50 • daily

headaches • a migraine so severe as to require bed rest • problems or serious side effects

from taking pain relievers or fever reducers • stomach problems that last or come back,

such as heartburn, upset stomach, or stomach pain • ulcers • bleeding problems • high

blood pressure • kidney or liver problems • heart or lung disease • reached age 60 or older

Ask a doctor or pharmacist before use if you are: • taking any other drug containing an NSAID (prescription or nonprescription) • taking a

blood thinning (anticoagulant) or steroid drug • under a doctor's care for any serious

condition • taking any other drug

When using this product take with food or milk if stomach upset occurs • long term

continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if: • migraine pain is not relieved or gets worse after first

dose • you feel faint, vomit blood, or have bloody or black stools. These are signs of

stomach bleeding • pain gets worse or lasts more than 10 days • any new symptoms

appear

• pregnant or breast-feeding, ask a health professional before use. It is especially

important to avoid using the last 3 weeks of pregnancy unless clearly

indicated to do so by a doctor because it may cause problems in the unborn child or

complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a

Poison Control Center right away.

Directions

• do not take more than directed

adults.

• take 2 capsules with a glass of water

• if symptoms persist or worsen, ask your doctor

• do not take more than 2 capsules in 24 hours, unless directed by a doctor

• do not take longer than 10 days, unless directed by a doctor (see Warnings)

• under 18 years of age, ask a doctor

Other information • read all warnings and directions before use. Keep carton,

• store at 20°-25°C (68°-77°F) • avoid high humidity and excessive heat above 40°C

(104°F) • do not use if temperature is not under bottle cap imprinted with "Sealed for

Your Protection" is broken or missing

Inactive Ingredients gelatin, mannitol, pharmaceutical ink, polyethylene

glycol, povidone, purified water, sorbitol, vitamin E, polyethylene glycol succinate

and may contain FD&C Blue #1.

Questions or comments? Call toll free 1-800-447-1140

Manufactured by:

Banner Pharmaceuticals, Inc.

4125 Premier Drive, High Point, NC 27265

Lot No.

Exp. Date:

(b) (4)

Customer: BANNER

Product Number: MIGRAINE NSAID 40CT

Revision Level: _____

Size: 2.125 X 6.25

Artist: JLS PP# 25527BANN

Proof Date: 11/11/2005

Proofread: JLS _____

Artist Q.C.

Pharmalabel

7915 INDUSTRIAL VILLAGE ROAD, GREENSBORO, NC 27409

PHONE 336-855-7142 • FAX 336-854-3753

www.pharmalabel.com

1st Proof 2nd Proof 3rd Proof 4th Proof _____ Other


Revise and resubmit proof

PHARMALABEL provides this proof to represent the image prepared for you according to the instructions provided. PHARMALABEL is not responsible for errors or omissions unless clearly indicated on this proof.

This proof does not accurately represent the colors that will print in the finished product. Any colors that are referenced on this proof are representative only. The finished product will be printed to match the referenced Pantone Color or any approved match standard provided by the customer.

Approved By: _____

Date: _____



Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
 • do not take more than directed
 • adults:
 • take 2 capsules with a glass of water
 • if symptoms persist or worsen, ask your doctor
 • do not take more than 2 capsules in 24 hours, unless directed by a doctor
 • do not take longer than 10 days, unless directed by a doctor (see Warnings)
 • under 18 years of age: ask a doctor
Other information
 • read all warnings and directions before use. Keep carton
 • store at 20-25°C (68-77°F)

Drug Facts (continued)
 • avoid high humidity and excessive heat above 40°C (104°F)
 • do not use if Tamper-Evident seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing
Inactive ingredients
 gelatin, mannitol, pharmaceutical ink, polyethylene glycol, povidone, purified water, sorbitan, sorbitol, vitamin E polyethylene glycol succinate and may contain FD&C Blue #1.
Questions or Comments?
 call toll free 1-800-447-1140

Drug Facts
Active ingredient (in each capsule) Purpose
 Ibuprofen 200 mg (NSAID)*Pain reliever
 *nonsteroidal anti-inflammatory drug
Use • treats migraine
Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 • hives • facial swelling • asthma (wheezing)
 • shock • skin reddening • rash • blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:
 • are age 60 or older
 • have had stomach ulcers or bleeding problems

• take a blood thinning (anticoagulant) or steroid drug
 • take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
 • have 3 or more alcoholic drinks every day while using this product
 • take more or for a longer time than directed
Do not use
 • if you have ever had an allergic reaction to any other pain reliever/fever reducer
 • right before or after heart surgery
Ask a doctor before use if you have
 • never had migraines diagnosed by a health professional
 • a headache that is different from your usual migraines
 • the worst headache of your life
 • fever and stiff neck
 • headaches beginning after or caused by head injury, exertion, coughing or bending
 • experienced your first headache after the age of 50
 • daily headaches
 • a migraine so severe as to require bed rest

Drug Facts (continued)
 • problems or serious side effects from taking pain relievers or fever reducers
 • stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
 • ulcers
 • bleeding problems
 • high blood pressure
 • heart or kidney disease
 • taken a diuretic
 • reached age 60 or older
Ask a doctor or pharmacist before use if you are
 • taking any other drug containing an NSAID (prescription or nonprescription)
 • taking a blood thinning (anticoagulant) or steroid drug
 • under a doctor's care for any serious condition
 • taking any other drug
When using this product
 • take with food or milk if stomach upset occurs

Drug Facts (continued)
 • long term continuous use may increase the risk of heart attack or stroke
Stop use and ask a doctor if
 • migraine pain is not relieved or gets worse after first dose
 • you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
 • pain gets worse or lasts more than 10 days
 • stomach pain or upset gets worse or lasts
 • any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

DIE 0437C
 9.4375 x 8.59375 Flat
 2.03125 x 2.03125 x 3.875

1st Proof 2nd Proof 3rd Proof 4th Proof Other
Revise and resubmit proof
 PHARMALABEL provides this proof to represent the image prepared for you according to the instructions provided. PHARMALABEL is not responsible for errors or omissions unless clearly indicated on this proof.
 This proof does not accurately represent the colors that will print in the finished product. Any colors that are referenced on this proof are representative only. The finished product will be printed to match the referenced Pantone Color or any approved match standard provided by the customer.

Proof Date: 11/15/05
 Proofread: JLS Artist
Pharmalabel
 7915 INDUSTRIAL VILLAGE ROAD, GREENSBORO, NC 27409
 PHONE 336-855-7142 • FAX 336-854-3753
 www.pharmalabel.com



Customer: BANNER
 Product Number: MIGRAINE 200MG 4OCT
 Revision Level: REV 11/05
 Size: DIE 0437C
 Artist: JLS PP# 25528BANN

Approved By: _____
 Date: _____

BLACK
 VARNISH

(b) (4)

DIE 0437C
 9.4375 x 8.59375 Flat
 2.03125 x 2.03125 x 3.875



Manufactured by: Banner Pharmaceuticals Inc.
 4125 Premier Drive, High Point, NC 27265

Exp. Date: _____
 Lot No.: _____
 Rev # 11/05

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
 • do not take more than directed
 • adults:
 • take 2 capsules with a glass of water
 • if symptoms persist or worsen, ask your doctor
 • do not take more than 2 capsules in 24 hours, unless directed by a doctor
 • do not take longer than 10 days, unless directed by a doctor (see Warnings)
 • under 18 years of age: ask a doctor
Other information
 • read all warnings and directions before use. Keep carton
 • store at 20-25°C (68-77°F)

Drug Facts (continued)
 • avoid high humidity and excessive heat above 40°C (104°F)
 • do not use if Tamper-Evident seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing
Inactive ingredients
 gelatin, mannitol, pharmaceutical ink, polyethylene glycol, povidone, purified water, sorbitan, sorbitol, vitamin E polyethylene glycol succinate and may contain FD&C Blue #1.
Questions or Comments?
 call toll free 1-800-447-1140

NDC 010888-0000-00
See new warnings information
Migraine Relief
Ibuprofen Capsules, 200 mg
Pain Reliever (NSAID)
 40 Softgels*
 *liquid filled capsules

Drug Facts
Active ingredient (in each capsule) Purpose
 Ibuprofen 200 mg (NSAID)*Pain reliever
 *nonsteroidal anti-inflammatory drug
Use • treats migraine
Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 • hives • facial swelling • asthma (wheezing)
 • shock • skin reddening • rash • blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:
 • are age 60 or older
 • have had stomach ulcers or bleeding problems

• take a blood thinning (anticoagulant) or steroid drug
 • take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
 • have 3 or more alcoholic drinks every day while using this product
 • take more or for a longer time than directed
Do not use
 • if you have ever had an allergic reaction to any other pain reliever/fever reducer
 • right before or after heart surgery
Ask a doctor before use if you have
 • never had migraines diagnosed by a health professional
 • a headache that is different from your usual migraines
 • the worst headache of your life
 • fever and stiff neck
 • headaches beginning after or caused by head injury, exertion, coughing or bending
 • experienced your first headache after the age of 50
 • daily headaches
 • a migraine so severe as to require bed rest

Drug Facts (continued)
 • problems or serious side effects from taking pain relievers or fever reducers
 • stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
 • ulcers
 • bleeding problems
 • high blood pressure
 • heart or kidney disease
 • taken a diuretic
 • reached age 60 or older
Ask a doctor or pharmacist before use if you are
 • taking any other drug containing an NSAID (prescription or nonprescription)
 • taking a blood thinning (anticoagulant) or steroid drug
 • under a doctor's care for any serious condition
 • taking any other drug
When using this product
 • take with food or milk if stomach upset occurs

Drug Facts (continued)
 • long term continuous use may increase the risk of heart attack or stroke
Stop use and ask a doctor if
 • migraine pain is not relieved or gets worse after first dose
 • you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
 • pain gets worse or lasts more than 10 days
 • stomach pain or upset gets worse or lasts
 • any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Ibuprofen Capsules, 200 mg
 Pain Reliever(NSAID)
 40 Softgels*
 *liquid filled capsules

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION



Customer: BANNER
 Product Number: MIGRAINE 200MG 40CT
 Revision Level: REV 11/05
 Size: DIE 0437C
 Artist: JLS PP# 25528BANN

Proof Date: 11/15/05 Q.C.
 Proofread: JLS Artist
Pharmalabel
 7915 INDUSTRIAL VILLAGE ROAD, GREENSBORO, NC 27409
 PHONE 336-855-7442 • FAX 336-854-3753
 www.pharmalabel.com

1st Proof 2nd Proof 3rd Proof 4th Proof
 Revise and resubmit proof
 Other
 PHARMALABEL provides this proof to represent the image prepared for you according to the instructions given on the PHARMALABEL. PHARMALABEL is not responsible for errors or omissions unless clearly indicated on this proof.
 This proof does not accurately represent the colors that will print in the finished product. Any colors that are referenced on this proof are representative only. The finished product will be printed to match the referenced Pantone Color or any approved match standard provided by the customer.

BLACK
 VARNISH

Approved By: _____
 Date: _____

Headings= 7 pt

Body Copy= 4.5pt

Sub Headings= 4.5 pt

4.5 pt

6 pt

4.5 pt

4.5 pt

7 pt

8 pt

16 pt

9 pt

9 pt

10 pt

8 pt

NDC 010888-0000-00

Pain reliever

Purpose

Active ingredient (in each capsule)
Ibuprofen 200 mg (NSAID) nonsteroidal anti-inflammatory drug

Use
Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 • rash
 • skin redness
 • hives (weal/itch)
 • asthma (wheezing)
 • blisters
 • shock

Warnings
 Allergic reaction occurs. Stop use and seek medical help right away.
 Stomach bleeding warnings: This product contains a nonsteroidal anti-inflammatory drug (NSAID) which may increase the risk of bleeding problems, such as:
 • taking other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
 • taking other drugs containing aspirin
 • taking other drugs containing aspirin for a longer time than directed
 • taking more or for a longer time than directed
 • taking other drugs containing aspirin
 • taking other drugs containing aspirin

Do not use if you have ever had an allergic reaction to other pain reliever/fever reducers.

Ask a doctor before use if you have:
 • a headache that is different from your usual migraines • the worst headache of your life • fever and stiff neck • headaches beginning after or caused by head injury, sinus infection, or a cold
 • a stomach ulcer, indigestion, or heartburn • a history of stomach ulcers, bleeding, or other problems from taking pain relievers or fever reducers • stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain • ulcers • bleeding problems • high blood pressure • kidney disease • liver disease • asthma • a history of asthma
 • taking any other drug containing an NSAID (prescription or nonprescription) • taking a blood thinning (anticoagulant) or steroid drug • under a doctor's care for any serious condition
 • taking any other drug containing aspirin

When using this product buy with food or milk if stomach upset occurs • long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if • migraine pain is not relieved or gets worse after first dose • you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding. • pain gets worse or lasts more than 10 days • any new symptoms

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a poison expert. Center right only.

Directions
 • do not take more than directed
 • take 2 capsules with a glass of water
 • if symptoms persist or worsen, ask your doctor
 • do not take more than 2 capsules in 24 hours, unless directed by a doctor
 • do not take longer than 10 days, unless directed by a doctor (see Warnings) under 10 years of age; ask a doctor

Other information • read all warnings and directions before use. Keep carton, capsules at 20-25°C (68°-77°F); avoid high humidity and excessive heat above 40°C (104°F). Store in original container. Do not use if you see mold, or see under bottle cap impaired with "Sealed for Your Protection" is broken or missing

Inactive Ingredients methylcellulose, hydroxypropyl methylcellulose, croscarmellose sodium, hydroxypropyl methylcellulose, polyethylene glycol succinate, and may contain FD&C Blue #1.

Questions or comments? Call toll free 1-800-447-1140

Manufactured by:
Banner Pharmaceuticals, Inc.
4125 Premier Drive, High Point, NC 27655

Lot No. _____
Exp. Date: _____

See new warnings information.

Migraine Relief

Ibuprofen Capsules, 200 mg

Pain Reliever (NSAID)

40 Softgels*
*liquid filled capsules

(B) (4)

Customer: BANNER

Product Number: MIGRAINE NSAID 40CT

Revision Level: SPEC

Size: 2.125 X 6.25

Artist: JLS PP# 25527BANN

Proof Date: 11/11/2005

Proofread: JLS _____
Artist Q.C.

Pharmalabel

7915 INDUSTRIAL VILLAGE ROAD, GREENSBORO, NC 27409
PHONE 336-855-7142 • FAX 336-854-3753
www.pharmalabel.com

1st Proof 2nd Proof 3rd Proof 4th Proof _____ Other _____

Revise and resubmit proof

PHARMALABEL provides this proof to represent the image prepared for you according to the instructions provided. PHARMALABEL is not responsible for errors or omissions unless clearly indicated on this proof.
 This proof does not accurately represent the colors that will print in the finished product. Any colors that are referenced on this proof are representative only. The finished product will be printed to match the referenced Pantone Color or any approved match standard provided by the customer.

BLACK

Approved By: _____

Date: _____

**This is a representation of an electronic record that was signed electronically and
this page is the manifestation of the electronic signature.**

/s/

Andrea Segal
3/13/2006 10:26:52 AM