

Drug Facts

Active ingredient (in each tablet)
Loperamide HCl 2 mg.....**Purpose**
Anti-diarrheal

Use
controls symptoms of diarrhea, including Traveler's Diarrhea

Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use if you have bloody or black stool

Ask a doctor before use if you have
■ fever ■ mucus in the stool ■ a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product ■ tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if
■ symptoms get worse
■ diarrhea lasts for more than 2 days
■ you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, call your nearest medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

■ take only on an empty stomach (1 hour before or 2 hours after a meal)

■ find right dose on chart. If possible, use weight to dose, otherwise, use age.

adults and children 12 years and over	chew 2 tablets after the first loose stool; chew 1 tablet after each subsequent loose stool; but no more than 4 tablets in 24 hours
children 9-11 years (60-95 lbs)	chew 1 tablet after the first loose stool; chew 1/2 tablet after each subsequent loose stool; but no more than 3 tablets in 24 hours

Drug Facts (continued)

children 6-8 years (40-59 lbs)	chew 1 tablet after the first loose stool; chew 1/2 tablet after each subsequent loose stool; but no more than 2 tablets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Other information
■ store between 20-25°C (68-77°F)
■ do not use if carton is open or if pitted foil seal under bottle cap is open or torn
■ see bottom panel for lot number and expiration date

Inactive ingredients
acetaminophen, potassium, basic polyurethane, cellulose, cellulose acetate, croscollon, croscollon, D&C yellow #10, dextrose, FD&C blue #1, flavors, magnesium stearate, sucralose

Questions or comments?
call 1-877-895-3465 (English) or 1-888-466-9746 (Spanish)

3 0045-0397-20 1

Imodium[®] A-D Loperamide HCl / Anti-Diarrheal

EZ CHEWS

20 Chewable CoolMint Flavored Tablets

NDC 50580-397-20

NEW!

Anti-Diarrheal

Controls the symptoms of diarrhea

20 Chewable CoolMint Flavored Tablets

Each chewable tablet contains 2 mg loperamide HCl

McNeil Consumer Healthcare
DIVISION OF MCNEIL-PPC, INC.
FORT WASHINGTON, PA 19034 USA
© MCN-PPC, Inc. '06
www.imodium.com

Imodium[®] A-D Loperamide HCl / Anti-Diarrheal

EZ CHEWS

20 Chewable CoolMint Flavored Tablets

Actual Size

00000000

00000000

20-ct bottle label:

Active ingredient (in each tablet)
Loperamide HCl 2 mg.....Anti-diarrheal

Use controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever

NDC 50580-397-20

Imodium®

A-D Loperamide HCl/
Anti-Diarrheal

EZ CHEWS
Cool Mint Flavored Tablets

20 Chewable

had a rash or other allergic reaction to loperamide HCl. Do not use if you have bloody or black stool. Ask a doctor before use if you have a fever, mucus in the stool, a history of liver disease. Ask a doctor or pharmacist before use if you are taking antibiotics. When using this product, tiredness,

XXXXXXX

drowsiness or dizziness may occur. Be careful when driving or operating machinery. **Stop use and ask a doctor if** symptoms get worse • diarrhea lasts for more than 2 days • you get abdominal swelling or bloating. These may be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) **Directions • drink plenty of clear fluids to help prevent dehydration caused by diarrhea •** take only on an empty stomach (1 hour before or 2 hours after a meal) • find right dose on chart, if possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	chew 2 tablets after the first loose stool; chew 1 tablet after each subsequent loose stool; but no more than 4 tablets in 24 hours
children 9-11 years (60-95 lbs)	chew 1 tablet after the first loose stool; chew 1/2 tablet after each subsequent loose stool; but no more than 3 tablets in 24 hours
children 6-8 years (48-59 lbs)	chew 1 tablet after the first loose stool; chew 1/2 tablet after each subsequent loose stool; but no more than 2 tablets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Other information • store between 20-25°C (68-77°F) • **do not use if printed foil seal under bottle cap is open or torn**

Questions or comments? call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish)

DISTRIBUTED BY: MCNEIL-PPC, INC., FORT WASHINGTON, PA 19034 USA © MCN-PPC, Inc. '06