

<b>Drug Facts</b>	
<b>Active ingredients</b> Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	<b>Purpose</b> Antiseptic Antiseptic
<b>Use</b> ■ for the preparation of the patient's skin prior to surgery	
<b>Warnings</b> <b>For external use only.</b> <b>Flammable, keep away from fire or flame. To reduce the risk of fire:</b> ■ solution contains alcohol and gives off <b>flammable vapors</b> ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin) ■ avoid getting solution into hairy areas. Solution may take much longer to dry or may not dry completely. ■ do not allow solution to pool ■ remove wet materials from prep area <b>Do not use</b> ■ in children less than 2 months of age because of the potential for excessive skin irritation and increased drug absorption ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser <b>When using this product</b> ■ keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. <b>Stop use and ask a doctor if</b> ■ irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> ■ to reduce the risk of fire the following strategies are recommended: ■ use in a well ventilated area ■ avoid getting solution into hairy areas. If this occurs, wipe hair with towel. Solution may take much longer to dry or may not dry completely. ■ do not allow solution to pool ■ tuck prep towels to absorb solution, and then remove ■ remove wet material from prep area ■ maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm <sup>2</sup> ). Discard the applicator after a single use. ■ pinch the wings on the applicator to break the ampule and release the antiseptic. Do not touch the sponge. Wet the sponge by repeatedly pressing and releasing the sponge against the treatment area until liquid is visible on the skin. ■ <b>dry surgical sites</b> (such as abdomen or arm): Use repeated back-and-forth strokes of the sponge for approximately 30 seconds. Completely wet the treatment area with antiseptic. Allow the area to air dry for approximately 30 seconds. Do not blot or wipe away. ■ <b>moist surgical sites</b> (such as the inguinal fold): Use repeated back-and-forth strokes of the sponge for approximately 2 minutes. Completely wet the treatment area with antiseptic. Allow the area to air dry for approximately one (1) minute. Do not blot or wipe away.	
<b>Other information</b> ■ store between 15-30 °C (59-86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)	
<b>Inactive ingredients</b> ■ FD&C green #3 dye ■ USP purified water	
<b>Questions?</b> Call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST) ■ www.chloraprep.com	

**ChloroPrep With Tint**  
Chlorhexidine Gluconate 2% w/v and Isopropyl Alcohol 70% v/v  
Patient Preparation Skin Preparation - 5-mL Applicator  
5-mL Applicators - 25 qty.

**Drug Facts**

**Warnings**

**Directions**

**Other information**

**Inactive ingredients**

**Questions?** Call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST) ■ www.chloraprep.com

Form for tracking ChloroPrep With Tint 5-mL Applicators usage. Includes a table for recording Date, Qty, Lot #, and What was changed?

Date	Qty	Lot #	What was changed?




Color key:  
 Green: PAB 017C  
 Yellow: PAB 100C  
 Red: PAB 100C  
 Black: PAB 200 (2004)

**enturla**  
ChloroPrep With Tint  
5-mL Applicators - 25 qty.

**enturla**  
ChloroPrep With Tint  
5-mL Applicators - 25 qty.

**enturla**  
ChloroPrep With Tint  
5-mL Applicators - 25 qty.

**enturla**  
ChloroPrep With Tint  
5-mL Applicators - 25 qty.

<b>WARNING</b>		Cat No. 260425	NDC #054365-400-10
<b>Flammable</b>		<b>ChloroPrep<sup>®</sup></b> <b>With Tint</b>	
	Keep away from fire or flame. To reduce risk of fire:	Chlorhexidine Gluconate 2% w/v Isopropyl Alcohol 70% v/v • 3.0-mL Applicator Patient Preoperative Skin Preparation	
	<ul style="list-style-type: none"><li>• solution contains alcohol and gives off flammable vapors</li><li>• do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin)</li><li>• avoid getting solution into hairy areas. Solution may take much longer to dry or may not dry completely.</li><li>• do not allow solution to pool</li><li>• remove wet material from prep area</li></ul>	External Use Only	Professional Use Only. Sterile Contents:* Applicator (1)
<small>Artwork 6-775425 • Rev. A • Issue Date 12/05</small>		STERILE EO	 Do Not Reuse 
		<b>See Drug Facts for full labeling information</b>	
		<small>* Sterility of contents guaranteed unless package is damaged or open.</small>	

Cat. No. 260425 NDC #054365-400-10  
 Single Use • Applicator is STERILE if package is intact

## ChloroPrep® With Tint

Chlorhexidine Gluconate 2% w/v and Isopropyl Alcohol 70% v/v  
 Patient Preoperative Skin Preparation • 3.0-ml Applicator

**WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.**

### Drug Facts

Active ingredients	Purpose
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

**Use** for the preparation of the patient's skin prior to surgery

### Warnings

**For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire:**

- solution contains alcohol and gives off **flammable vapors**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin)
- avoid getting solution into hairy areas. Solution may take much longer to dry or may not dry completely.
- do not allow solution to pool
- remove wet materials from prep area

### Do not use

- in children less than 2 months of age because of the potential for excessive skin irritation and increased drug absorption
- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

### Directions

- to reduce the risk of fire the following strategies are recommended:
  - use in a well ventilated area
  - avoid getting solution into hairy areas. If this occurs, wipe hair with towel. Solution may take much longer to dry or may not dry completely.
  - do not allow solution to pool
  - tuck prep towels to absorb solution, and then remove
  - remove wet material from prep area
- maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm<sup>2</sup>). Discard the applicator after a single use.
- pinch the wings on the applicator to break the ampule and release the antiseptic. Do not touch the sponge. Wet the sponge by repeatedly pressing and releasing the sponge against the treatment area until liquid is visible on the skin.
- **dry surgical sites** (such as abdomen or arm): Use repeated back-and-forth strokes of the sponge for approximately 30 seconds. Completely wet the treatment area with antiseptic. Allow the area to air dry for approximately **30 seconds**. Do not blot or wipe away.
- **moist surgical sites** (such as the inguinal fold): Use repeated back-and-forth strokes of the sponge for approximately 2 minutes. Completely wet the treatment area with antiseptic. Allow the area to air dry for approximately one **(1) minute**. Do not blot or wipe away.

### Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

### Inactive ingredients

- FD&C green #3 dye ■ USP purified water

### Questions?

www.chloroprep.com  
 Call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)