

Swabstick Package Label

NDC 10819-4075-2

REORDER NO. S32450

Chlorascrub™ Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation

*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

- For external use only ■ Use for the preparation of the patient's skin prior to surgery or injection ■ Latex free
- Helps reduce bacteria that potentially can cause skin infection ■ For hospital and professional use only

IMPORTANT: See insert for complete *Drug Facts* and safety information.
One Swabstick, 0.054 fl.oz. (1.6 mL) Each



Made in U.S.A.

CONTENTS: One swabstick premoistened with 1.6 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water. **DIRECTIONS:** Maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm). Tear open the pouch to expose the swabstick. Remove the swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply swabstick to the procedure site. Place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute. Turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute. Allow the prepped area to air dry for one and one half minutes (1.5 minutes). Do not blot or wipe dry. Discard after a single use. **WARNINGS:** For external use only. Flammable, keep away from fire or flame. Do not use with electrocautery procedures. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

IMPORTANT: See insert for complete *Drug Facts* and safety information.

PDI®, The Healthcare Division of Nice-Pak Products, Inc., Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • www.pdipi.com
Chlorascrub™ is under license by American Solumed Inc.

EXP.

THIS SIDE UP
TEAR OPEN HERE

2AS324

Maxi Swabstick Package Label

NDC 10819-4075-3

REORDER NO. S27350

Chlorascrub™ Maxi Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation

*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

- For external use only ■ Use for the preparation of the patient's skin prior to surgery or injection
- Helps reduce bacteria that potentially can cause skin infection
- For hospital and professional use only ■ Latex free

IMPORTANT: See insert for complete *Drug Facts* and safety information.

One Maxi Swabstick, 0.172 fl.oz. (5.1 mL) Each



Made in U.S.A.

CONTENTS: One Maxi Swabstick premoistened with 5.1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water.

DIRECTIONS: Maximum treatment area for one maxi swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm). Tear open the pouch to expose the swabstick. Remove swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply swabstick to the procedure site. Place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute. Turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute. Allow the prepped area to air dry for one and one half minutes (1.5 minutes). Do not blot or wipe dry. Discard after a single use.

WARNINGS: For external use only. Flammable, keep away from fire or flame. Do not use with electrocautery procedures. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

IMPORTANT: See insert for complete *Drug Facts* and safety information.

PDI®, The Healthcare Division of Nice-Pak Products, Inc., Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • www.pdipi.com
Chlorascrub™ is under license by American Solumed Inc.

EXP.

THIS SIDE UP
TEAR OPEN HERE

2AS273

Swabstick Package Insert

REORDER NO. S32450
NDC 10819-4075-2

Chlorascrub™ Swabstick
[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

Drug Facts		Drug Facts (continued)	
Active ingredients	Purpose	Directions	
Chlorhexidine gluconate 3.15% w/v.....	Antiseptic	<ul style="list-style-type: none"> ■ maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm) ■ tear open the pouch to expose the swabstick ■ remove the swabstick from package. Do not touch the foam tip. ■ prior to surgery or injection, apply swabstick to the procedure site ■ place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute ■ turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute ■ allow the prepped area to air dry for one and one half minutes (1.5 minutes) ■ do not blot or wipe dry ■ discard after a single use 	
Isopropyl alcohol 70% v/v.....	Antiseptic	Other information	
Use for the preparation of the patient's skin prior to surgery or injection		<ul style="list-style-type: none"> ■ store between 68-77°F (20-25°C) ■ avoid freezing and excessive heat above 104°F (40°C) ■ for hospital and professional use only 	
Warnings		Inactive ingredient	
<p>For external use only</p> <p>Flammable, keep away from fire or flame</p> <p>Do not use with electrocautery procedures</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ in premature or low birth weight infants, or on children less than 2 months of age because of the potential for excessive skin irritation and increased drug absorption ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser ■ under occlusive patch <p>When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>		USP purified water	
		Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)	

50S324IS

Each Swabstick contains 0.054 fl.oz. (1.6 mL)



PDI®, The Healthcare Division of Nice-Pak Products, Inc., Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • Chlorascrub™ is under license by American Solumed Inc.

www.pdipdi.com
Made in U.S.A.

Maxi Swabstick Package Insert

REORDER NO. S27350
NDC 10819-4075-3

Chlorascrub™ Maxi Swabstick
[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

Drug Facts		Drug Facts (continued)	
Active ingredients	Purpose	Directions	
Chlorhexidine gluconate 3.15% w/v.....	Antiseptic	<ul style="list-style-type: none"> ■ maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm) ■ tear open the pouch to expose the swabstick ■ remove the swabstick from package. Do not touch the foam tip. ■ prior to surgery or injection, apply swabstick to the procedure site ■ place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute ■ turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute ■ allow the prepped area to air dry for one and one half minutes (1.5 minutes) ■ do not blot or wipe dry ■ discard after a single use 	
Use for the preparation of the patient's skin prior to surgery or injection		Other information	
Warnings		<ul style="list-style-type: none"> ■ store between 68-77°F (20-25°C) ■ avoid freezing and excessive heat above 104°F (40°C) ■ for hospital and professional use only 	
<p>For external use only Flammable, keep away from fire or flame Do not use with electrocautery procedures</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ in premature or low birth weight infants, or on children less than 2 months of age because of the potential for excessive skin irritation and increased drug absorption ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser ■ under occlusive patch <p>When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>		Inactive ingredient USP purified water	
		Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)	

50S273IS

Each Maxi Swabstick contains 0.172 fl.oz. (5.1 mL)



PDI®, The Healthcare Division of Nice-Pak Products, Inc., Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • www.pdipi.com
Chlorascrub™ is under license by American Solumed Inc. Made in U.S.A.

Swabstick Shipper Label

NET CONTENTS: 500 INDIVIDUAL SWABSTICKS
REORDER NO. S32450
NDC 10819-4075-2

Chlorascrub™ Swabstick
[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation

*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

<p>Drug Facts</p>	<p>Drug Facts (continued)</p>
<p>Active ingredients</p> <p>Chlorhexidine gluconate 3.15% w/v..... Antiseptic Isopropyl alcohol 70% v/v..... Antiseptic</p>	<p>Directions</p> <ul style="list-style-type: none"> ■ maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm) ■ tear open the pouch to expose the swabstick ■ remove the swabstick from package. Do not touch the foam tip. ■ prior to surgery or injection, apply swabstick to the procedure site ■ place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute ■ turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute ■ allow the prepped area to air dry for one and one half minutes (1.5 minutes) ■ do not blot or wipe dry ■ discard after a single use
<p>Use for the preparation of the patient's skin prior to surgery or injection</p>	
<p>Warnings</p> <p>For external use only Flammable, keep away from fire or flame Do not use with electrocautery procedures</p>	
<p>Do not use</p> <ul style="list-style-type: none"> ■ in premature or low birth weight infants, or on children less than 2 months of age because of the potential for excessive skin irritation and increased drug absorption ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser ■ under occlusive patch 	<p>Other information</p> <ul style="list-style-type: none"> ■ store between 68-77°F (20-25°C) ■ avoid freezing and excessive heat above 104°F (40°C) ■ for hospital and professional use only
<p>When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p>	<p>Inactive ingredient USP purified water</p>
<p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p>	
<p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>	<p>Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>



80S32450
YK

Each Swabstick contains 0.054 fl.oz. (1.6 mL)



PDI®, The Healthcare Division of Nice-Pak Products, Inc., Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • www.pdipdi.com
 Chlorascrub™ is under license by American Solumed Inc. Made in U.S.A.

Maxi Swabstick Shipper Label

NET CONTENTS: 300 INDIVIDUAL SWABSTICKS
 REORDER NO. S27350
 NDC 10819-4075-3

Chlorascrub™ Maxi Swabstick
 [Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
 Patient preoperative skin preparation

*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

Drug Facts	Drug Facts (continued)				
<p>Active ingredients</p> <table border="0"> <tr> <td>Chlorhexidine gluconate 3.15% w/v.....</td> <td>Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v.....</td> <td>Antiseptic</td> </tr> </table>	Chlorhexidine gluconate 3.15% w/v.....	Antiseptic	Isopropyl alcohol 70% v/v.....	Antiseptic	<p>Directions</p> <ul style="list-style-type: none"> ■ maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm) ■ tear open the pouch to expose the swabstick ■ remove the swabstick from package. Do not touch the foam tip. ■ prior to surgery or injection, apply swabstick to the procedure site ■ place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute ■ turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute ■ allow the prepped area to air dry for one and one half minutes (1.5 minutes) ■ do not blot or wipe dry ■ discard after a single use
Chlorhexidine gluconate 3.15% w/v.....	Antiseptic				
Isopropyl alcohol 70% v/v.....	Antiseptic				
<p>Use for the preparation of the patient's skin prior to surgery or injection</p>					
<p>Warnings</p> <p>For external use only</p> <ul style="list-style-type: none"> ■ Flammable, keep away from fire or flame ■ Do not use with electrocautery procedures 					
<p>Do not use</p> <ul style="list-style-type: none"> ■ in premature or low birth weight infants, or on children less than 2 months of age because of the potential for excessive skin irritation and increased drug absorption ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser ■ under occlusive patch 	<p>Other information</p> <ul style="list-style-type: none"> ■ store between 68-77°F (20-25°C) ■ avoid freezing and excessive heat above 104°F (40°C) ■ for hospital and professional use only 				
<p>When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p>					
<p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p>					
<p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>	<p>Inactive ingredient USP purified water</p>				
	<p>Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>				



8AS27350
YK

Each Maxi Swabstick contains 0.172 fl.oz. (5.1 mL)



PDI®, The Healthcare Division of Nice-Pak Products, Inc., Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • www.pdipdi.com
 Chlorascrub™ is under license by American Solumed Inc. Made in U.S.A.