

100 Tablets
Antihistamine
5 mg
Tablets
**Cetirizine
Hydrochloride**
NDC 49884-645-01

Fonts: Univers, light, plain, bold, oblique- Cond.,
 Drug Facts: 9 pt. Leading: 7.5 pt.
 Header: 8 pt. Bullets: 5 pt.
 Subheader: 6 pt. Barlines: 1.5 pt.
 Body Text: 6 pt. Hairlines: .5 pt.
 Drug Facts (continued) 9 pt.
 Horizontal Scale: 90-100% Kerning: 0

Drug Facts (continued)

If pregnant or breast-feeding:
 ■ if breast-feeding, not recommended
 ■ if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 6 years and over	1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 2 tablets in 24 hours.
Adults 65 years and over	1 tablet once a day; do not take more than 1 tablet in 24 hours.
Children under 6 years of age	ask a doctor
Consumers with liver or kidney disease	ask a doctor

Other information
 ■ store between 20° to 25°C (68° to 77°F).

Inactive ingredients
 hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, polysorbate, povidone, pregelatinized starch, and titanium dioxide.

Control No:

Exp. Date:

I12/07

Par Pharmaceutical Companies, Inc.
 Spring Valley, NY 10977



Original Prescription Strength
 NDC 49884-645-01
**Cetirizine
 Hydrochloride**
Tablets
5 mg
Antihistamine

Do not use if imprinted foil inner seal on bottle is broken or missing.

ALLERGY Indoor & Outdoor Allergies

24 Relief of
 hour Sneezing
 Runny Nose
 Itchy, Watery Eyes
 Itchy Throat or Nose

100 Tablets



Drug Facts

Active ingredient (in each tablet) Purpose
 Cetirizine Hydrochloride 5 mg Antihistamine

Uses
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ runny nose ■ sneezing ■ itchy, watery eyes
 ■ itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product
 ■ drowsiness may occur ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. ▶

100 Tablets
Antihistamine
10 mg
Tablets
Cetirizine
Hydrochloride
 NDC 49884-646-01

Fonts: Univers, light, plain, bold, oblique- Cond.,
 Drug Facts: 9 pt. Leading: 7.5 pt.
 Header: 8 pt. Bullets: 5 pt.
 Subheader: 6 pt. Barlines: 1.5 pt.
 Body Text: 6 pt. Hairlines: .5 pt.
 Drug Facts (continued) 9 pt.
 Horizontal Scale: 90-100% Kerning: 0

Drug Facts (continued)

If pregnant or breast-feeding:
 ■ if breast-feeding, not recommended
 ■ if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 6 years and over	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms
Adults 65 years and over	ask a doctor
Children under 6 years of age	ask a doctor
Consumers with liver or kidney disease	ask a doctor

Other information
 ■ store between 20° to 25°C (68° to 77°F).

Inactive ingredients
 hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, polysorbate, povidone, pregelatinized starch, and titanium dioxide.

Control No:
 Exp. Date:

I12/07

Par Pharmaceutical Companies, Inc.
 Spring Valley, NY 10977



Original Prescription Strength
 NDC 49884-646-01
Cetirizine
Hydrochloride
Tablets
10 mg
Antihistamine

Do not use if imprinted foil inner seal on bottle is broken or missing.

ALLERGY Indoor & Outdoor Allergies

24 Relief of
 hour Sneezing
 Runny Nose
 Itchy, Watery Eyes
 Itchy Throat or Nose

100 Tablets

PAR
 PHARMACEUTICAL

Drug Facts

Active ingredient (in each tablet) Purpose
 Cetirizine Hydrochloride 10 mg Antihistamine

Uses
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ runny nose ■ sneezing ■ itchy, watery eyes
 ■ itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product
 ■ drowsiness may occur ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. ▶

NDC 49884-645-01

Cetirizine Hydrochloride Tablets

5 mg

ALLERGY

Antihistamine

Do not use if imprinted foil inner seal on bottle is broken or missing.

100 Tablets



Active ingredient (in each tablet) Purpose
Cetirizine Hydrochloride 5 mg Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor before use if you have liver or kidney disease.** Your doctor should determine if you need a different dose. **Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.** **When using this product** ■ drowsiness may occur

■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness

■ be careful when driving a motor vehicle or operating machinery. **Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- If breast-feeding: not recommended
- if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions **Adults and children 6 years and over:** 1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 2 tablets in 24 hours. **Adults 65 years and over:** 1 tablet once a day, do not take more than 1 tablet in 24 hours. **Children under 6 years of age:** ask a doctor. **Consumers with liver or kidney disease:** ask a doctor. **Other information** ■ store between 20° to 25°C (68° to 77°F).

Inactive ingredients hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, polysorbate, povidone, pregelatinized starch, and titanium dioxide.



0 49884 - 645 - 01 4

Control No.:
Exp. Date:

11/2/07
Par Pharmaceutical Cos., Inc.
Spring Valley, NY 10977

Fonts: Univers, light, plain, bold, oblique- Cond.,	
Body Text:	6 pt. Bullets: 5 pt.
Horizontal Scale: 90-100%	Kerning: 0

NDC 49884-646-01

Cetirizine Hydrochloride Tablets

10 mg **ALLERGY**

Antihistamine

Do not use if imprinted foil inner seal on bottle is broken or missing.

100 Tablets



Active ingredient (in each tablet) Purpose
Cetirizine Hydrochloride 10 mg Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose ■ sneezing ■ itchy, watery eyes
- itching of the nose or throat

Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor before use if you have liver or kidney disease.** Your doctor should determine if you need a different dose. **Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives. When using this product** ■ drowsiness may occur

■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness

■ be careful when driving a motor vehicle or operating machinery. **Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- If breast-feeding: not recommended
- If pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions Adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours.

A 5 mg product may be appropriate for less severe symptoms. **Adults 65 years and over:** ask a doctor. **Children under 6 years of age:** ask a doctor. **Consumers with liver or kidney disease:** ask a doctor. **Other information** ■ store between 20° to 25°C (68° to 77°F).

Inactive ingredients hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, polysorbate, povidone, pregelatinized starch, and titanium dioxide.



0 49884-646-01 1

Control No.:
Exp. Date:

112/07
Par Pharmaceutical Cos., Inc.
Spring Valley, NY 10977

Fonts: Univers, light, plain, bold, oblique- Cond.,	
Body Text:	6 pt. Bullets: 5 pt.
Horizontal Scale: 90-100%	Kerning: 0