



Tamper Evident: do not use if foil seal under cap is missing, open or broken.



NDC 0378-3635-01

ALLERGY

### Cetirizine Hydrochloride Tablets 5 mg antihistamine

#### Indoor & Outdoor Allergies

- 24 Hour Relief of
- runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat

100 TABLETS      RM-AR3635A1

#### Drug Facts

##### Active Ingredient (in each tablet)

Cetirizine Hydrochloride 5 mg

##### Purpose

Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

##### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

OPEN FOR FULL LABELING

Front Layer

1 of 3

BLACK   200 RED   306 BLUE

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20

Drug Facts:	9 pt
Headers	7 pt
Body Text	6 pt
Drug Facts (continued)	8 pt
Leading	7 pt/6.5 pt
Bullets	5 pt
Frames	1.5 pt
Bars	1 pt
Rules	0.5 pt

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur    ■ avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor** if an allergic reaction to this product occurs.

Seek medical help right away.

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts (continued)****Directions**

adults and children  
6 years and over

1 to 2 tablets once daily depending upon  
severity of symptoms; do not take more than  
2 tablets in 24 hours.

adults 65 years  
and over

1 tablet once a day; do not take more than  
1 tablet in 24 hours

children under  
6 years of age

ask a doctor

consumers with liver  
or kidney disease

ask a doctor

Back of Front Layer 2

2 of 3

**BLACK**

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold  
Cond. #20; Trade Gothic Bold Cond. Italics #20

<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

PERMANENT GLUE

REMOVABLE GLUE

 NDC 0378-3635-01  
**ALLERGY**  
**Cetirizine Hydrochloride**  
**Tablets 5 mg**  
**antihistamine**  
**Indoor & Outdoor Allergies**  
**24 Hour Relief of** ■ runny nose  
■ sneezing ■ itchy, watery eyes  
■ itching of the nose or throat  
100 TABLETS **RM-AR3635A1**

**Drug Facts** (continued)

---

**Other information**

■ store between 20° to 25°C (68° to 77°F)

---

**Inactive ingredients**

anhydrous lactose, colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium lauryl sulfate, titanium dioxide, and triacetin

---

**Questions?** call 1-877-446-3679

---

**Mylan Pharmaceuticals Inc.**  
**Morgantown, WV 26505**

Base Layer

3 of 3

BLACK

<b>FONTS:</b> Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20	
<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts</b> (continued)	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt



Tamper Evident: do not use if foil seal under cap is missing, open or broken.

 NDC 0378-3635-05  
**ALLERGY**

**Cetirizine Hydrochloride**  
**Tablets 5 mg**  
**antihistamine**

**Indoor & Outdoor Allergies**  
**24 Hour Relief of** ■ runny nose  
 ■ sneezing ■ itchy, watery eyes  
 ■ itching of the nose or throat

500 TABLETS **RM-AR3635B1**

**Drug Facts**

Active Ingredient (in each tablet)	Purpose
Cetirizine Hydrochloride 5 mg	Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  
 ■ runny nose ■ sneezing ■ itchy, watery eyes  
 ■ itching of the nose or throat

**Warnings**  
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.  
 Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

OPEN FOR FULL LABELING

Front Layer

1 of 3

BLACK 200 RED 306 BLUE

<b>FONTS:</b> Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20	
<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts</b> (continued)	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur    ■ avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if an allergic reaction to this product occurs.  
Seek medical help right away.

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts (continued)****Directions**

adults and children  
6 years and over

1 to 2 tablets once daily depending upon  
severity of symptoms; do not take more than  
2 tablets in 24 hours.

adults 65 years  
and over

1 tablet once a day; do not take more than  
1 tablet in 24 hours

children under  
6 years of age

ask a doctor

consumers with liver  
or kidney disease

ask a doctor

Back of Front Layer 2

2 of 3

**BLACK**

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold  
Cond. #20; Trade Gothic Bold Cond. Italics #20

<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

PERMANENT GLUE

REMOVABLE GLUE

 NDC 0378-3635-05  
**ALLERGY**  
**Cetirizine Hydrochloride**  
**Tablets 5 mg**  
**antihistamine**  
**Indoor & Outdoor Allergies**  
24 Hour Relief of ■ runny nose  
■ sneezing ■ itchy, watery eyes  
■ itching of the nose or throat  
500 TABLETS **RM-AR3635B1**

*Drug Facts* (continued)

*Other information*

■ store between 20° to 25°C (68° to 77°F)

*Inactive ingredients*

anhydrous lactose, colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium lauryl sulfate, titanium dioxide, and triacetin

*Questions?* call 1-877-446-3679

**Mylan Pharmaceuticals Inc.**  
**Morgantown, WV 26505**

Base Layer

3 of 3

BLACK

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20

<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts</b> (continued)	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

NM  
0378-3637-01  
3



Tamper Evident: do not use if foil seal under cap is missing, open or broken.



NDC 0378-3637-01

ALLERGY

**Cetirizine Hydrochloride  
Tablets 10 mg  
antihistamine**

**Indoor & Outdoor Allergies**

**24 Hour Relief of** ■ runny nose  
■ sneezing ■ itchy, watery eyes  
■ itching of the nose or throat

100 TABLETS      RM-AR3637A1

**Drug Facts**

**Active Ingredient (in each tablet)**

Cetirizine Hydrochloride 10 mg

**Purpose**

Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

OPEN FOR FULL LABELING

Front Layer

1 of 3

BLACK 354 GREEN 306 BLUE

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20

<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

**Drug Facts** (continued)

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur    ■ avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if an allergic reaction to this product occurs.**  
Seek medical help right away.

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ▶

**Drug Facts** (continued)**Directions**

adults and children  
6 years and over

one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.

adults 65 years and over

ask a doctor

children under  
6 years of age

ask a doctor

consumers with liver  
or kidney disease

ask a doctor ▶

Back of Front Layer 2

2 of 3

**BLACK**

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20

<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts</b> (continued)	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt



■ PERMANENT GLUE

■ REMOVABLE GLUE

 NDC 0378-3637-01  
MYLAN® **ALLERGY**

**Cetirizine Hydrochloride  
Tablets 10 mg  
antihistamine**

**Indoor & Outdoor Allergies**  
24 Hour Relief of ■ runny nose  
■ sneezing ■ itchy, watery eyes  
■ itching of the nose or throat

100 TABLETS **RM-AR3637A1**

**Drug Facts** (continued)

**Other information**

■ store between 20° to 25°C (68° to 77°F)

**Inactive ingredients**

anhydrous lactose, colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium lauryl sulfate, titanium dioxide, and triacetin

**Questions?** call 1-877-446-3679

**Mylan Pharmaceuticals Inc.  
Morgantown, WV 26505**

Base Layer

3 of 3

**BLACK**

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20

<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts</b> (continued)	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt



Tamper Evident: do not use if foil seal under cap is missing, open or broken.



NDC 0378-3637-05

ALLERGY

**Cetirizine Hydrochloride  
Tablets 10 mg  
antihistamine**

**Indoor & Outdoor Allergies**

- 24 Hour Relief of** ■ runny nose  
 ■ sneezing ■ itchy, watery eyes  
 ■ itching of the nose or throat

500 TABLETS **RM-AR3637B1**

**Drug Facts**

**Active Ingredient (in each tablet)**

Cetirizine Hydrochloride 10 mg ..... Antihistamine

**Purpose**

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose ■ sneezing ■ itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

OPEN FOR FULL LABELING

Front Layer

1 of 3

BLACK 354 GREEN 306 BLUE

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20

<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur    ■ avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor** if an allergic reaction to this product occurs.  
Seek medical help right away.

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts (continued)****Directions**

adults and children  
6 years and over

one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.

adults 65 years and over

ask a doctor

children under  
6 years of age

ask a doctor

consumers with liver  
or kidney disease

ask a doctor

Back of Front Layer 2

2 of 3

**BLACK**

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20

<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

■ PERMANENT GLUE

■ REMOVABLE GLUE

 NDC 0378-3637-05  
MYLAN® **ALLERGY**

**Cetirizine Hydrochloride  
Tablets 10 mg  
antihistamine**

**Indoor & Outdoor Allergies**  
24 Hour Relief of ■ runny nose  
■ sneezing ■ itchy, watery eyes  
■ itching of the nose or throat

500 TABLETS      RM-AR3637B1

**Drug Facts** (continued)

**Other information**

■ store between 20° to 25°C (68° to 77°F)

**Inactive ingredients**

anhydrous lactose, colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium lauryl sulfate, titanium dioxide, and triacetin

**Questions?** call 1-877-446-3679

**Mylan Pharmaceuticals Inc.  
Morgantown, WV 26505**

Base Layer


3 of 3

BLACK

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20

**Drug Facts:** 9 pt  
**Headers** 7 pt  
**Body Text** 6 pt  
**Drug Facts** (continued) 8 pt  
**Leading** 7 pt/6.5 pt  
**Bullets** 5 pt  
**Frames** 1.5 pt  
**Bars** 1 pt  
**Rules** 0.5 pt

N  
3 0378-3635-01 9





NDC 0378-3635-01

HIVES  
Relief

24 **Relief of**  
**HOUR ITCHING**  
Due to Hives

Tampers Evident: do not use if foil seal under cap  
is missing, open or broken

100 TABLETS **RM-H3635A1**

**Drug Facts**

**Active Ingredient (in each tablet)**

Cetirizine Hydrochloride 5 mg ..... Antihistamine

**Purpose**

---

**Use** relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

**Warnings**

**Severe Allergy Warning:** Get emergency help **immediately** if you have hives along with any of the following symptoms:

- trouble swallowing   ■ swelling of tongue   ■ trouble speaking   ■ wheezing or problems breathing
- dizziness or loss of consciousness   ■ swelling in or around mouth   ■ drooling

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

**Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. ▶

OPEN FOR FULL LABELING

Front Layer

1 of 3

BLACK PANT. PURPLE 306 BLUE

<b>FONTS:</b> Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20	
<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

**Drug Facts (continued)**

If you have been prescribed an epinephrine injector, you should carry it with you at all times.

**Do not use**

- to prevent hives from any known cause such as:
  - foods
  - insect stings
  - medicines
  - latex or rubber glovesbecause this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have**

- liver or kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch ▶

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use. ▶

Back of Front Layer 2

2 of 3

BLACK

<b>Fonts:</b> Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20	
<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

■ PERMANENT GLUE

■ REMOVABLE GLUE


 NDC 0378-3635-01  
  
**Cetirizine Hydrochloride**  
**Tablets 5 mg**  
**antihistamine**  
**24** Relief of  
**ITCHING**  
**Hour** Due to Hives  
Tamper Evident: do not use if foil seal under cap is missing, open or broken  
**100 TABLETS    RM-H3635A1**

**Drug Facts** (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

adults and children 6 years and over	1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 2 tablets in 24 hours.
adults 65 years and over	1 tablet once a day; do not take more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Drug Facts** (continued)

**Other information**

- store between 20° to 25°C (68° to 77°F)

**Inactive ingredients**

anhydrous lactose, colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium lauryl sulfate, titanium dioxide, and triacetin

**Questions?** call 1-877-446-3679

**Mylan Pharmaceuticals Inc.**  
**Morgantown, WV 26505**

Base Layer

3 of 3

BLACK

<b>Fonts:</b> Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20	
<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts</b> (continued)	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

3  
0378-3635-05  
7





NDC 0378-3635-05

HIVES  
Relief

Cetirizine Hydrochloride  
Tablets 5 mg  
antihistamine

24 Relief of  
HOUR ITCHING  
Due to Hives

Tampers Evident: do not use if foil seal under cap  
is missing, open or broken

500 TABLETS    RM-H3635B1

Drug Facts

Active Ingredient (in each tablet)	Purpose
Cetirizine Hydrochloride 5 mg .....	Antihistamine

**Use** relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

**Warnings**

**Severe Allergy Warning:** Get emergency help **immediately** if you have hives along with any of the following symptoms:

- trouble swallowing   ■ swelling of tongue   ■ trouble speaking   ■ wheezing or problems breathing
- dizziness or loss of consciousness   ■ swelling in or around mouth   ■ drooling

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

**Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. ▶

OPEN FOR FULL LABELING

Front Layer

1 of 3

BLACK   PANT. PURPLE   306 BLUE

<b>FONTS:</b> Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20	
<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt



**Drug Facts (continued)**

If you have been prescribed an epinephrine injector, you should carry it with you at all times.

**Do not use**

- to prevent hives from any known cause such as:
    - foods
    - insect stings
    - medicines
    - latex or rubber gloves
- because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have**

- liver or kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch ▶

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use. ▶

Back of Front Layer 2

2 of 3

BLACK

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20

<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

■ PERMANENT GLUE

■ REMOVABLE GLUE


 NDC 0378-3635-05  
  
**Cetirizine Hydrochloride**  
**Tablets 5 mg**  
**antihistamine**  
**24** Relief of  
**ITCHING**  
**HOUR** Due to Hives  
Tamper Evident: do not use if foil seal under cap is missing, open or broken  
**500 TABLETS    RM-H3635B1**

**Drug Facts** (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

adults and children 6 years and over	1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 2 tablets in 24 hours.
adults 65 years and over	1 tablet once a day; do not take more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Drug Facts** (continued)

**Other information**

- store between 20° to 25°C (68° to 77°F)

**Inactive ingredients**

anhydrous lactose, colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium lauryl sulfate, titanium dioxide, and triacetin

**Questions?** call 1-877-446-3679

**Mylan Pharmaceuticals Inc.**  
**Morgantown, WV 26505**


Base Layer

3 of 3

BLACK

<b>FONTS:</b> Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20	
<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts</b> (continued)	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

N  
3 0378-3637-01 3





NDC 0378-3637-01

HIVES  
Relief

24 Relief of  
**ITCHING**  
Due to Hives

Tamper Evident: do not use if foil seal under cap  
is missing, open or broken

100 TABLETS    **RM-H3637A1**

**Drug Facts**

**Active Ingredient (in each tablet)**

Cetirizine Hydrochloride 10 mg	Antihistamine
--------------------------------	---------------

---

**Use** relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

---

**Warnings**

**Severe Allergy Warning:** Get emergency help **immediately** if you have hives along with any of the following symptoms:

- trouble swallowing    ■ swelling of tongue    ■ trouble speaking    ■ wheezing or problems breathing
- dizziness or loss of consciousness    ■ swelling in or around mouth    ■ drooling

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

**Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. ▶

OPEN FOR FULL LABELING

Front Layer

1 of 3

BLACK    PANT. PURPLE    306 BLUE

<b> FONTS:</b> Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20	
<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

**Drug Facts (continued)**

If you have been prescribed an epinephrine injector, you should carry it with you at all times.

**Do not use**

- to prevent hives from any known cause such as:
  - foods
  - insect stings
  - medicines
  - latex or rubber glovesbecause this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have**

- liver or kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch ▶

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use. ▶

Back of Front Layer 2

2 of 3

BLACK

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20

<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

■ PERMANENT GLUE

■ REMOVABLE GLUE


 NDC 0378-3637-01  
**MYLAN®** **HIVES Relief**

**Cetirizine Hydrochloride**  
**Tablets 10 mg**  
**antihistamine**

**24** Relief of  
**HOUR** **ITCHING**  
Due to Hives

Tamper Evident: do not use if foil seal under cap is missing, open or broken

**100 TABLETS**    **RM-H3637A1**

**Drug Facts** (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

adults and children 6 years and over	one 10 mg tablet once daily, do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
adults 65 years and over	ask a doctor
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Drug Facts** (continued)

**Other information**

- store between 20° to 25°C (68° to 77°F)

**Inactive ingredients**

anhydrous lactose, colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium lauryl sulfate, titanium dioxide, and triacetin

**Questions?** call 1-877-446-3679

**Mylan Pharmaceuticals Inc.**  
**Morgantown, WV 26505**

Base Layer

3 of 3

BLACK

<b>FONTS:</b> Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20
<b>Drug Facts:</b> 9 pt
<b>Headers</b> 7 pt
<b>Body Text</b> 6 pt
<b>Drug Facts</b> (continued) 8 pt
<b>Leading</b> 7 pt/6.5 pt
<b>Bullets</b> 5 pt
<b>Frames</b> 1.5 pt
<b>Bars</b> 1 pt
<b>Rules</b> 0.5 pt



Front Layer

BLACK PANT. PURPLE 306 BLUE

<b>FONTS:</b> Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20	
<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

**Drug Facts (continued)**

If you have been prescribed an epinephrine injector, you should carry it with you at all times.

**Do not use**

- to prevent hives from any known cause such as:
  - foods
  - insect stings
  - medicines
  - latex or rubber gloves
- because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have**

- liver or kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch ▶

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use. ▶

Back of Front Layer 2

2 of 3

BLACK

**FONTS:** Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20

<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts (continued)</b>	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt

■ PERMANENT GLUE

■ REMOVABLE GLUE


 NDC 0378-3637-05  
**MYLAN**<sup>®</sup>


**Cetirizine Hydrochloride**  
**Tablets 10 mg**  
**antihistamine**

**24** Relief of  
**HOUR** **ITCHING**  
Due to Hives

Tamper Evident: do not use if foil seal under cap is missing, open or broken

**500 TABLETS**    **RM-H3637B1**

**Drug Facts** (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

adults and children 6 years and over	one 10 mg tablet once daily, do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
adults 65 years and over	ask a doctor
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Drug Facts** (continued)

**Other information**

- store between 20° to 25°C (68° to 77°F)

**Inactive ingredients**

anhydrous lactose, colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium lauryl sulfate, titanium dioxide, and triacetin

**Questions?** call 1-877-446-3679

**Mylan Pharmaceuticals Inc.**  
**Morgantown, WV 26505**

Base Layer

3 of 3

BLACK

<b>Fonts:</b> Trade Gothic Cond. #18; Trade Gothic Bold Cond. #20; Trade Gothic Bold Cond. Italics #20	
<b>Drug Facts:</b>	9 pt
<b>Headers</b>	7 pt
<b>Body Text</b>	6 pt
<b>Drug Facts</b> (continued)	8 pt
<b>Leading</b>	7 pt/6.5 pt
<b>Bullets</b>	5 pt
<b>Frames</b>	1.5 pt
<b>Bars</b>	1 pt
<b>Rules</b>	0.5 pt