

**Pharmacist Information:**

Dispense in a tight, light-resistant container as described in the USP.

Keep out of the reach of children.

Store at 20° to 25°C (68° to 77°F)  
[See USP Controlled Room Temperature].

Protect from moisture.

**Each Tablet Contains:**

Carvedilol.....12.5 mg

C.S. No. 5413L02

Iss. 8/07

**NDC 57664-245-88**

# Carvedilol Tablets

**12.5 mg**

**100 TABLETS**

Pharmacist: Dispense With Information For Patients.  
Rx Only



**USUAL DOSAGE:**

See package outsert for complete product information.



**Pharmacist Information:**

Dispense in a tight, light-resistant container as described in the USP.

Keep out of the reach of children.

Store at 20° to 25°C (68° to 77°F)  
[See USP Controlled Room Temperature].

Protect from moisture.

**Each Tablet Contains:**

Carvedilol.....12.5 mg

C.S. No. 5414L02  
Iss. 8/07

NDC 57664-245-13

# Carvedilol Tablets

**12.5 mg**


500 TABLETS

Pharmacist: Dispense With Information For Patients.  
Rx Only



**USUAL DOSAGE:**

See package outsert for complete product information.



**Pharmacist Information:**

Dispense in a tight, light-resistant container as described in the USP.

Keep out of the reach of children.

Store at 20° to 25°C (68° to 77°F)  
[See USP Controlled Room Temperature].

Protect from moisture.

**Each Tablet Contains:**

Carvedilol.....12.5 mg

C.S. No. 5415L02  
Iss. 8/07

**NDC 57664-245-18**

# Carvedilol Tablets

**12.5 mg**

**1000 TABLETS**

**Pharmacist: Dispense With Information For Patients.  
Rx Only**



**USUAL DOSAGE:**

See package outsert for complete product information.



**Pharmacist Information:**

Dispense in a tight, light-resistant container as described in the USP.

Keep out of the reach of children.

Store at 20° to 25°C (68° to 77°F)  
[See USP Controlled Room Temperature].

Protect from moisture.

**Each Tablet Contains:**

Carvedilol.....25 mg

C.S. No. 5416L02

Iss. 8/07

**NDC 57664-247-88**

# Carvedilol Tablets

**25 mg**

**100 TABLETS**

Pharmacist: Dispense With Information For Patients.  
Rx Only



**USUAL DOSAGE:**

See package outsert for complete product information.



**Pharmacist Information:**

Dispense in a tight, light-resistant container as described in the USP.

Keep out of the reach of children.

Store at 20° to 25°C (68° to 77°F)  
[See USP Controlled Room Temperature].

Protect from moisture.

**Each Tablet Contains:**

Carvedilol.....25 mg

C.S. No. 5417L02  
Iss. 8/07

**NDC 57664-247-13**

# Carvedilol Tablets

**25 mg**

**500 TABLETS**

**Pharmacist: Dispense With Information For Patients.  
Rx Only**



**USUAL DOSAGE:**

See package insert for complete product information.



**Pharmacist Information:**

Dispense in a tight, light-resistant container as described in the USP.

Keep out of the reach of children.

Store at 20° to 25°C (68° to 77°F)  
[See USP Controlled Room Temperature].

Protect from moisture.

**Each Tablet Contains:**

Carvedilol.....25mg

C.S. No. 5418L02

Iss. 8/07

**NDC 57664-247-18**

# Carvedilol Tablets

**25 mg**

**1000 TABLETS**

**Pharmacist: Dispense With Information For Patients.  
Rx Only**



**USUAL DOSAGE:**

See package outsert for complete product information.



**Pharmacist Information:**

Dispense in a tight, light-resistant container as described in the USP.

Keep out of the reach of children.

Store at 20° to 25°C (68° to 77°F)  
[See USP Controlled Room Temperature].

Protect from moisture.

**Each Tablet Contains:**

Carvedilol.....6.25 mg

C.S. No. 5410L02

Iss. 8/07

**NDC 57664-244-88**

# Carvedilol Tablets

**6.25 mg**

**100 TABLETS**

Pharmacist: Dispense With Information For Patients.  
Rx Only



**USUAL DOSAGE:**

See package outsert for complete product information.



**Pharmacist Information:**

Dispense in a tight, light-resistant container as described in the USP.

Keep out of the reach of children.

Store at 20° to 25°C (68° to 77°F)  
[See USP Controlled Room Temperature].

Protect from moisture.

**Each Tablet Contains:**

Carvedilol.....6.25 mg

C.S. No. 5411L02

Iss. 8/07

NDC 57664-244-13

# Carvedilol Tablets

**6.25 mg**

500 TABLETS

Pharmacist: Dispense With Information For Patients.  
Rx Only



**USUAL DOSAGE:**

See package outsert for complete product information.





**Pharmacist Information:**

Dispense in a tight, light-resistant container as described in the USP.

Keep out of the reach of children.

Store at 20° to 25°C (68° to 77°F)  
[See USP Controlled Room Temperature].

Protect from moisture.

**Each Tablet Contains:**

Carvedilol.....6.25 mg

C.S. No. 5412L02  
Iss. 8/07

**NDC 57664-244-18**

# Carvedilol Tablets

**6.25 mg**

**1000 TABLETS**

**Pharmacist: Dispense With Information For Patients.  
Rx Only**



**USUAL DOSAGE:**

See package insert for complete product information.



## INFORMATION FOR PATIENTS TAKING CARVEDILOL TABLETS

### Rx only

Read the patient information that comes with CARVEDILOL before you start taking it and each time you get a refill. There may be new information. This information does not take the place of talking with your doctor about your medical condition or your treatment. If you have any questions about CARVEDILOL, ask your doctor or pharmacist.

### WHAT IS CARVEDILOL?

CARVEDILOL is a prescription medicine that belongs to a group of medicines called “beta-blockers”. CARVEDILOL is used, often with other medicines, for the following conditions:

- To treat patients with high blood pressure (hypertension)
- To treat patients who had a heart attack that worsened how well the heart pumps

CARVEDILOL is not approved for use in children under 18 years of age.

### WHO SHOULD NOT TAKE CARVEDILOL?

Do not take CARVEDILOL if you:

- Have severe heart failure and are hospitalized in the intensive care unit or require certain intravenous medications that help support circulation (inotropic medications)
- Are prone to asthma or other breathing problems
- Have a slow heartbeat or a heart that skips a beat (irregular heartbeat)
- Have liver problems
- Are allergic to any of the ingredients in carvedilol tablet. The active ingredient is carvedilol. See the end of this leaflet for a list of all the ingredients in carvedilol tablets.

### WHAT SHOULD I TELL MY DOCTOR BEFORE TAKING CARVEDILOL?

Tell your doctor about all of your medical conditions, including if you:

- Have asthma or other lung problems (such as bronchitis or emphysema)
- Have problems with blood flow in your feet and legs (peripheral vascular disease). CARVEDILOL can make some of your symptoms worse.
- Have diabetes
- Have thyroid problems
- Have a condition called pheochromocytoma
- Have had severe allergic reactions
- Are pregnant or trying to become pregnant. It is not known if CARVEDILOL is safe for your unborn baby. You and your doctor should talk about the best way to control your high blood pressure during pregnancy.
- Are breastfeeding. It is not known if CARVEDILOL passes into your breast milk. You should not breastfeed while using CARVEDILOL.
- Are scheduled for surgery and will be given anesthetic agents
- Are taking prescription or non-prescription medicines, vitamins, and herbal supplements. CARVEDILOL and certain other medicines can affect each other and cause serious side effects. CARVEDILOL may affect the way other medicines work. Also, other medicines may affect how well CARVEDILOL works.

Keep a list of all the medicines you take. Show this list to your doctor and pharmacist before you start a new medicine.

### HOW SHOULD I TAKE CARVEDILOL?

**It is important for you to take your medicine every day as directed by your doctor. If you stop taking CARVEDILOL suddenly, you could have chest pain and/or a heart attack. If your doctor decides that you should stop taking CARVEDILOL, your doctor may slowly lower your dose over a period of time before stopping it completely.**

- Take CARVEDILOL exactly as prescribed. Your doctor will tell you how many tablets to take and how often. In order to minimize possible side effects, your doctor might begin with a low dose and then slowly increase the dose.
- **Do not stop taking CARVEDILOL and do not change the amount of CARVEDILOL you take without**

#### **talking to your doctor.**

- Tell your doctor if you gain weight or have trouble breathing while taking CARVEDILOL.
- Take CARVEDILOL with food.
- If you miss a dose of CARVEDILOL, take your dose as soon as you remember, unless it is time to take your next dose. Take your next dose at the usual time. Do not take 2 doses at the same time.
- If you take too much CARVEDILOL, call your doctor or poison control center right away.

#### **WHAT SHOULD I AVOID WHILE TAKING CARVEDILOL?**

CARVEDILOL can cause you to feel dizzy, tired, or faint. Do not drive a car, use machinery, or do anything that needs you to be alert if you have these symptoms.

#### **WHAT ARE POSSIBLE SIDE EFFECTS OF CARVEDILOL?**

- **Low blood pressure (which may cause dizziness or fainting when you stand up).** If these happen, sit or lie down right away and tell your doctor.
- **Tiredness.** If you feel tired or dizzy you should not drive, use machinery, or do anything that needs you to be alert.
- **Slow heart beat**
- **Changes in your blood sugar. If you have diabetes, tell your doctor if you have any changes in your blood sugar levels.**
- CARVEDILOL may hide some of the symptoms of low blood sugar, especially a fast heartbeat.
- CARVEDILOL may mask the symptoms of hyperthyroidism (overactive thyroid).
- **Worsening of severe allergic reactions.**

Other side effects of CARVEDILOL include shortness of breath, weight gain, diarrhea, and fewer tears or dry eyes that become bothersome if you wear contact lenses.

Call your doctor if you have any side effects that bother you or don't go away.

#### **How should I store CARVEDILOL?**

- Store carvedilol tablet at 20° to 25°C (68° to 77° F). [See USP Controlled Room Temperature]. Keep the tablets dry.
- Safely, throw away CARVEDILOL that is out of date or no longer needed.
- Keep CARVEDILOL and all medicines out of the reach of children.

#### **GENERAL INFORMATION ABOUT CARVEDILOL**

Medicines are sometimes prescribed for conditions other than those described in patient information leaflets. Do not use CARVEDILOL for a condition for which it was not prescribed. Do not give CARVEDILOL to other people, even if they have the same symptoms you have. It may harm them.

This leaflet summarizes the most important information about CARVEDILOL. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about CARVEDILOL that is written for healthcare professionals. You can also find out more about CARVEDILOL by visiting the website [www.caraco.com](http://www.caraco.com) or calling 1-800-818-4555. This call is free.

#### **WHAT ARE THE INGREDIENTS IN CARVEDILOL TABLET?**

Active Ingredient: Carvedilol

Inactive Ingredients: mannitol, povidone, colloidal silicon dioxide, sucrose, crospovidone, talc, magnesium stearate, and opadry white

Carvedilol tablets come in the following strengths: 6.25 mg, 12.5 mg, 25 mg





