

Original Prescription Strength

Cetirizine HCl
Tablets, 5 mg
Antihistamine
Allergy

Indoor & Outdoor Allergies
24 Hour Relief of
Sneezing
Runny Nose
Itchy, Watery Eyes
Itchy Throat or Nose

30 TABLETS
5 mg EACH

TAMPER EVIDENT; DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.

Active ingredient
(in each tablet)

Cetirizine HCl 5 mg

Purpose
Antihistamine

Use Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor** before use if you are taking tranquilizers or sedatives, if you have been drinking alcohol, or if you have ever had seizures. **Ask a doctor** if you are pregnant or taking a motor vehicle or operating machinery. **Stop use** if you experience increased drowsiness or blurred vision, or if you are taking any other drugs. **Seek medical help** right away if **pregnant or breast-feeding**; ■ if breast-feeding; not recommended ■ if pregnant or breastfeeding. ■ If you experience any of the above symptoms, contact a Poison Control Center right away.

Directions **Adults and children 6 years and over:** 1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 5 tablets in 24 hours. **Children 6 years and over:** 1 tablet once a day; do not take more than 1 tablet in 24 hours. **Children under 6 years of age:** ask a doctor. **Consumers with liver or kidney disease:** ask a doctor.

Other information ■ store between 20° to 25° C (68° to 77° F)

Inactive ingredients See carton for complete list of ingredients

Questions? call 1-800-406-7984

XXXX
Distributed by:
Chm Laboratories Inc.
14 Terminal Road
New Brunswick, NJ 08901
Batch No. Expiration Date:
XXXXXXXXXX

Non-Varnish Area

Standard Format
Helvetica Condensed
Variable horizontal scale
adheres to 39 characters per inch

Active ingredient - 5.63 pt. Helv. Bold Oblique,
Uses - 5.1 pt. Helvetica Cnd. Bold Italic
Headings - 5.1 pt. Helv. Cnd. Bld. Italic
Sub Heads - 4.08 pt. Helv. Cnd. Bold,
Text - 4.08 pt. Helv. Cnd., min. leading 4.21,
horiz 106.86%

Square Bullets - 3.4 pt. Zapf Ding.,
Border 1 pt. rule, 1/16 gap all around
Barline 1pt. rule _____
Hairline .5 pt. rule _____

Original Prescription Strength

Cetirizine HCl Tablets, 5 mg

Antihistamine Allergy

Indoor & Outdoor Allergies

24 Hour Relief of

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

45 TABLETS
5 mg EACH

NEVER OPEN OR USE IF IMPAIRED SEAL IS BROKEN OR MISSING FROM BOTTLE.

Active Ingredient
(in each Tablet)
Cetirizine HCl 5 mg.....Antihistamine

Uses ■ temporarily relieves symptoms of the following allergic conditions: ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings ■ **Do not use** if you have ever had an allergic reaction to any medicine containing hydroxyzine. ■ **Ask a doctor before use** if you have liver or kidney disease. Your doctor should determine if you need a different dose. ■ **Ask a doctor or pharmacist** if you are taking any other medicine, especially sedatives. ■ **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase the drowsiness caused by this product ■ **Do not drive a motor vehicle or operate machinery. Stop uses and ask a doctor** if an allergic reaction to this product occurs. ■ **Seek medical help right away. If pregnant or planning to get pregnant, tell the health-care provider you are taking this medicine before use. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions: Adults and children 6 years and over: 1 to 2 tablets, 1 to 2 times a day. Children 2 to 5 years: do not take more than 2 tablets in 24 hours. Adults 65 years and over: 1 tablet once a day; do not take more than 1 tablet in 24 hours. ■ **Keep out of reach of children. Ask a doctor. Consumers with liver or kidney disease:** ask a doctor

Other Information ■ store between 20° to 25° C (68° to 77° F)

Inactive Ingredients See carton for complete list of ingredients

Questions? call 1-800-406-7384

XXXX

Distributed by:
Lannett Laboratories, Inc.
14000 Lannett Drive
New Brunswick, NJ 08901

Batch No. _____

Expiration Date: _____

Non Varnish Area

Standard Format
 Helvetica Condensed
 Variable horizontal scale
 adheres to 39 characters per inch

Active ingredient - 5.63 pt. Helv. Bold Oblique,
 Uses - 5.1 pt. Helvetica Cnd. Bold Italic
 Headings - 5.1 pt. Helv. Cnd. Bld. Italic
 Sub Heads - 4.08 pt. Helv. Cnd. Bold,
 Text - 4.08 pt. Helv. Cnd., min. leading 4.21,
 horiz 106.86%

Square Bullets - 3.4 pt. Zapf Ding.,
 Border 1 pt. rule, 1/16 gap all around
 Barline 1pt. rule _____
 Hairline .5 pt. rule _____

Distributed by:
Ohm Laboratories Inc.
14 Terminal Road
New Brunswick, NJ 08901



Original Prescription Strength

Cetirizine HCl Tablets, 5 mg Antihistamine

Allergy Indoor & Outdoor Allergies

**24 Hour Relief of
Sneezing
Runny Nose
Itchy, Watery Eyes
Itchy Throat or Nose**

**75 Tablets
5 mg EACH
(contains one bottle of 45 tablets
and one bottle of 30 tablets)**

Batch No.

Expiration Date:

Non Varnish Area

Cetirizine HCl Tablets, 5 mg

Antihistamine Allergy

Indoor & Outdoor Allergies

Drug Facts
Active ingredient (in each tablet):
Cetirizine HCl 5 mg.....Antihistamine
Purpose
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ itchy, watery eyes ■ itchy of the nose or throat ■ runny nose ■ sneezing ■ sneezing

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

Drug Facts (continued)
When using this product
■ drowsiness may occur
■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery
■ avoid alcoholic drinks
Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding: ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
adults and children 6 years and over
1 to 2 tablets once daily depending upon severity of symptoms;
adults 65 years and over
1 tablet once a day; do not take more than 1 tablet in 24 hours
children under 6 years of age
ask a doctor
consumers with liver or kidney disease
ask a doctor

Other Information
■ TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.
■ store between 20° to 25° C (68° to 77° F)

Inactive ingredients corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, talc, titanium dioxide

Questions? call 1-800-406-7984

Keep the carton. It contains important information.
See end panel for expiration date.

XXXX

Standard Format
Vertical Orientation
Vertical Orientation
Vertical Orientation
adheres to 39 characters per inch

Drug Facts - 11 pt. Helv. Bold Italic,
Drug Facts Continued - 9 pt. Helv. Bold Italic,
Headings - 8pt. Helv. Bold Italic,
Sub-heads - 6pt. Helv. Bold,
Text - 6pt. Helv. Bold, min. leading 6.5
Square Bullets - 5 pt. Zap Ding,
Border 2.5 pt. rule, 1/8 gap all around
Baseline 2.5pt. rule
Hairline .5 pt. rule

Original Prescription Strength

Cetirizine HCl Tablets, 5 mg

Antihistamine

Hives Relief

24 Hour Relief of Itching Due to Hives

30 TABLETS
5 mg EACH

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTOM.

Active Ingredient
(in each tablet)

Cetirizine HCl 5 mg

Purpose
Antihistamine

Uses relieves itching due to hives (urticaria). This product

is used to prevent hives or an allergic skin reaction from occurring.

Warnings
Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms: ■ trouble swallowing ■ swelling of your face, lips, tongue or throat ■ wheezing or trouble breathing ■ trouble speaking ■ drooling ■ wheezing or trouble breathing. These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated immediately. Stop use of this product and call your doctor immediately. ■ **Do not use if you are taking any of the following medicines:** ■ alcohol, sedatives, and tranquilizers may increase the effect of cetirizine HCl. ■ **Do not use if you have had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before using this product if you are taking any of the following medicines:** ■ alcohol, sedatives, and tranquilizers may increase the effect of cetirizine HCl. ■ **Do not use if you have had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before using this product if you are taking any of the following medicines:** ■ alcohol, sedatives, and tranquilizers may increase the effect of cetirizine HCl. ■ **Do not use if you have had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before using this product if you are taking any of the following medicines:** ■ alcohol, sedatives, and tranquilizers may increase the effect of cetirizine HCl. ■ **Do not use if you have had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. Ask a doctor before using this product if you are taking any of the following medicines:** ■ alcohol, sedatives, and tranquilizers may increase the effect of cetirizine HCl.

Directions Adults and children 6 years and over: 1 to 2 tablets once daily, depending upon severity of symptoms; do not take more than 10 tablets in 24 hours. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor.

Other Information ■ store between 20° to 25° C (68° to 77° F)

Inactive Ingredients See carton for complete list of ingredients

Questions? call 1-800-405-7894

XXXX

Distributed by:
Chm Laboratories Inc.

47 Terminal Park, NJ 08801

New Brunswick, NJ 08801

Batch No.

Expiration Date:

XXXXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

XXXXXX

Standard Format
Helvetica Condensed
Variable horizontal scale
adheres to 39 characters per inch

Active ingredient - 5.63 pt. Helv. Bold Oblique,

Uses - 5.1 pt. Helvetica Cnd. Bold Italic

Headings - 5.1 pt. Helv. Cnd. Bld. Italic

Sub Heads - 4.08 pt. Helv. Cnd. Bold,

Text - 4.08 pt. Helv. Cnd., min. leading 4.21, horiz 106.86%

Square Bullets - 3.4 pt. Zapf Ding.,

Border 1 pt. rule, 1/16 gap all around

Barline 1pt. rule _____

Hairline .5 pt. rule _____



Original Prescription Strength

Cetirizine HCl Tablets, 5 mg

Antihistamine

Hives Relief

24 Hour Relief of Itching Due to Hives

45 TABLETS
5 mg EACH

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.

Active Ingredient (in each tablet) Cetirizine HCl, 5 mg

Purpose Antihistamine

Uses relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from returning.

Warnings Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms: ■ trouble swallowing ■ swelling of the face, lips, tongue, or throat ■ difficulty breathing or normal mouth ■ trouble speaking ■ drooling ■ wheezing or problems breathing. These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. **Not a Substitute for Epinephrine.** If you do not have an epinephrine injector, you could occur with your hives; never use this product as a substitute for the epinephrine injector. If you have been told to use an epinephrine injector, use it as directed with you at all times. **Do not use ■ to prevent hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from returning. Hives can sometimes be the only way to prevent them.** Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor. Do not use this product if you are not able to help you find a cause. ■ If you have ever had an allergic reaction to this product or any of its ingredients, or to an antihistamine containing hydroxyzine. **Ask a doctor before use if you are pregnant, breastfeeding, or taking other medicines.** You should determine if you need a different dose. ■ Hives that are an unusual color, look bruised or blistered ■ Hives that do not itch. **Ask a doctor or pharmacist before use if you are taking other medicines.** ■ Do not use this product if the product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness. ■ Do not use this product if you are operating machinery. **Stop use and ask a doctor if ■ an allergic reaction to this product occurs. Seek medical help right away. ■ Symptoms do not improve after 3 days of use. ■ Symptoms do not improve after 7 days of use. ■ Symptoms do not improve after 14 days of use. ■ If pregnant or breast-feeding, ■ if breast-feeding, not recommended ■ if pregnant, ask a health professional before use. ■ Keep out of reach of children. In case of overdose, call your doctor for medical advice or contact a Poison Control Center right away.**

Directions Adults and children 6 years and over: 1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 1 tablet once daily. **Children 6 years and over:** 1 tablet in 24 hours. **Children under 6 years of age:** ask a doctor. **Consumers with liver or kidney disease:** ask a doctor.

Other information ■ Store between 20° to 25° C (68° to 77° F).

Inactive ingredients See carton for complete list of ingredients.

Questions? call 1-800-406-7994

XXXX
Distributed by:
Chim Laboratories Inc.
14 Terminal Road
New Brunswick, NJ 08901
Batch No. _____ Expiration Date: _____



Standard Format
 Helvetica Condensed
 Variable horizontal scale
 adheres to 39 characters per inch

Active ingredient - 5.63 pt. Helv. Bold Oblique,

Uses - 5.1 pt. Helvetica Cnd. Bold Italic

Headings - 5.1 pt. Helv. Cnd. Bld. Italic

Sub Heads - 4.08 pt. Helv. Cnd. Bold,

Text - 4.08 pt. Helv. Cnd., min. leading 4.21,
 horiz 106.86%

Square Bullets - 3.4 pt. Zapf Ding.,

Border 1 pt. rule, 1/16 gap all around

Barline 1pt. rule _____

Hairline .5 pt. rule _____



Distributed by:
 Ohm Laboratories Inc.
 14 Terminal Road
 New Brunswick, NJ 08901

Original Prescription Strength

Cetirizine HCl

Tablets, 5 mg

Antihistamine

Hives Relief

24 Hour Relief of Itching Due to Hives

75 Tablets
 5 mg EACH
 (contains one bottle of 45 tablets and one bottle of 30 tablets)

Batch No.

Expiration Date:

Non Varnish Area

Drug Facts (continued)
 Inactive ingredients: corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, talc, titanium dioxide

Keep the carton. It contains important information.
 See end panel for expiration date.

Drug Facts (continued)
 Active ingredient (in each tablet): Cetirizine HCl 5 mg, Antihistamine

Uses relieves itching due to hives (urticaria). This product will not prevent skin reaction from occurring.

Warnings
 Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms:
 ■ trouble swallowing ■ dizziness or loss of consciousness ■ swelling of tongue ■ swelling in or around mouth
 ■ trouble speaking ■ wheezing or problems breathing ■ drooling
 These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

Drug Facts (continued)
 Not a substitute for epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.
 Do not use ■ to prevent hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
 Ask a doctor before use if you have ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch ■ liver or kidney disease. Your doctor should determine if you need a different dose.
 Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery
 Stop use and ask a doctor if ■ symptoms do not improve after 3 days of treatment ■ the hives have lasted more than 6 weeks ■ an allergic reaction to this product occurs. Seek medical help right away.
 If pregnant or breast-feeding: ■ If breast-feeding, not recommended ■ If pregnant, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
 adults and children 6 years and over 1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 2 tablets in 24 hours
 children under 6 years of age ask a doctor
 ask a doctor
 ask a doctor
 ask a doctor
Other information
 ■ store between 20° to 25° C (68° to 77° F)
 ■ TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.

Standard Format
 10 Characters
 Vertical Orientation
 Adheres to 39 characters per inch

Drug Facts - 11 pt. Hxlv. Bold Italic,
 Drug Facts Continued - 9 pt. Hxlv. Bold Italic,
 Headings - 8pt. Hxlv. Bld. Italic,
 Sub heads - 6pt. Hxlv. Bld.
 Text - 6pt. Hxlv. Bld. min. leading 6.5
 Square Bullets - 5 pt. Zap Ding.
 Border 2.5 pt. rule, 1/8 gap all around
 Baseline 2.5pt. rule
 Hairline .5 pt. rule

Original Prescription Strength

Cetirizine HCl

Tablets, 10 mg

Antihistamine

Allergy

Indoor & Outdoor Allergies

24 Hour Relief of

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

7 TABLETS
10 mg EACH

Expiration Date: _____

Batch No. _____

Non Varnish Area

Drug Facts (continued)

Questions? call 1-800-406-7984

Keep the carton. It contains important information.

See end panel for expiration date.

XXXX

Drug Facts

Active ingredient (in each tablet)

Cetirizine HCl 10 mg.....Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Drug Facts (continued)

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding: If breast-feeding: not recommended. If pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 6 years and over

one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.

ask a doctor

children under 6 years of age

ask a doctor

ask a doctor

ask a doctor

ask a doctor

ask a doctor

Other information

TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.

store between 20° to 25° C (68° to 77° F)

Inactive ingredients corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, talc, titanium dioxide

Standard Format

Vertical text on horizontal scale adheres to 39 characters per inch

Drug Facts - 11 pt. Heli; Bold Italic;

Drug Facts Continued - 9 pt. Heli; Bold Italic;

Headings - 8pt. Heli; Bd; Italic;

Sub-heads - 6pt. Heli; Bd;

Text - 6pt. Heli; Bd; min. leading 6.5

Square Bullets - 5 pt. Zap Ding.

Border 2.5 pt. rule, 1/8 gap all around

Baseline 2.5pt. rule

Headline .5 pt. rule

125 C



Distributed by:
Ohm Laboratories Inc.
14 Terminal Road,
New Brunswick, NJ 08901

XXXX
Distributed by:
Omn Laboratories Inc.
14 Terminal Road,
New Brunswick, NJ 08901



Original Prescription Strength

Cetirizine HCl Tablets, 10 mg Antihistamine Hives Relief

24 Hour Relief of
itching Due to Hives

7 TABLETS
10 mg EACH

Keep the carton. It contains important information.
See end panel for expiration date.

Batch No. _____
Expiration Date: _____

Non Varnish Area

Drug Facts (continued)
Other information ■ store between 20° C to 25° C (68° F to 77° F)
■ TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.
Inactive ingredients corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, talc, titanium dioxide
? call 1-800-406-7984

Drug Facts
Active ingredient (in each tablet) Cetirizine HCl 10 mg
Antihistamine
Uses relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings (continued)
Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms:
■ trouble swallowing ■ dizziness or loss of consciousness ■ swelling of tongue ■ difficulty breathing
■ trouble speaking ■ wheezing or problems breathing ■ drooling
■ swelling in or around mouth
Not a Substitute for Epinephrine: If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergic symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.
Do not use ■ to prevent hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
■ If you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
Ask a doctor before use if you have ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch ■ liver or kidney disease. Your doctor should determine if you need a different dose.
Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives
■ drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery
When using this product ■ symptoms do not improve after 3 days of treatment ■ the hives have lasted more than 6 weeks ■ an allergic reaction to this product occurs. Seek medical help right away.
Stop use and ask a doctor if ■ If breast-feeding: not recommended ■ If pregnant: ask a health professional before use ■ If pregnant or breast-feeding:
Directions one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours.
adults and children 6 years and over
A 5 mg product may be appropriate for less severe symptoms.
adults 65 years and over
ask a doctor
children under 6 years of age
ask a doctor
consumers with liver or kidney disease
ask a doctor



Standard Format
Helvetica Medium Condensed
Variable horizontal scale
adheres to 39 characters per inch
Drug Facts - 11 pt. Helv. Bold Cond. Italic, cond 90%
Drug Facts (continued)
Drug Facts - 9pt Helv. Bold Cond. Italic, cond 90%
(continued) - 7pt. Helvetica, cond 85%
Headings - 6pt. Helv. Bld. Cond. Italic, cond 90%
Sub-Heads - 6pts. Helv. Bold, Cond 90%
Text 6pt. Helv., Med Cond. 90%, min. leading 6.5
Square bullets - 5pt. Zapf Ding., cond 85%
Border 2.5 pt. rule, 1/8 gap all around
Barline 2.0pt. rule
Headline .5 pt. rule



XXXXXX

Distributed by:
Olm Laboratories Inc.
14 Terminal Road,
New Brunswick, NJ 08901



010000001000000000

Original Prescription Strength

Cetirizine HCl Tablets, 5 mg Antihistamine

Allergy
Indoor & Outdoor Allergies

24 Hour Relief of
Sneezing
Runny Nose
Itchy, Watery Eyes
Itchy Throat or Nose

7 TABLETS
5 mg EACH

Expiration Date:

Batch No.

Non Varnish Area

Drug Facts (continued)
Questions? call 1-800-406-7984
Keep the carton. It contains important information.
See end panel for expiration date.

Drug Facts
Active ingredient (in each tablet)
Cetirizine HCl 5 mg.....Antihistamine
Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Drug Facts (continued)
Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.
When using this product:
■ drowsiness may occur ■ avoid alcoholic drinks
■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery.
Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding: ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
adults and children 6 years and over
1 to 2 tablets once daily depending upon severity of symptoms;
do not take more than 2 tablets in 24 hours
adults 65 years and over
1 tablet once a day; do not take more than 1 tablet in 24 hours
children under 6 years of age
ask a doctor
consumers with liver or kidney disease
ask a doctor
Other Information
■ TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.
■ store between 20° to 25° C (68° to 77° F)
Inactive ingredients corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, talc, titanium dioxide

Standard Format
Helvetica Condensed
Variable horizontal scale
adheres to 35 characters per inch
Drug Facts - 11 pt. Helv. Bold Italic,
Drug Facts Continued - 9 pt. Helv. Bold Italic,
Headings - 8pt. Helv. Bld. Italic,
Sub-Heads - 6pts. Helv. Bld.
Text 6pt. Helv. Bld. min. heading 6.5
Square Bullets - 5 pt. Zap Ding.
Border 2.5 pt. rule, 1/8 gap all around
Baseline 2.5pt. rule
Hairline .5 pt. rule

499 C

XXXX
Distributed by:
Omn Laboratories Inc.
14 Terminal Road,
New Brunswick, NJ 08901



Original Prescription Strength

Cetirizine HCl Tablets, 5 mg Antihistamine Hives Relief

24 Hour Relief of itching Due to Hives

7 TABLETS
5 mg EACH

Keep the carton. It contains important information.
See end panel for expiration date.

Batch No. _____
Expiration Date: _____

Non Varnish Area

Drug Facts (continued)
Other information ■ store between 20° to 25° C (68° to 77° F)
■ TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS BENT, BROKEN OR SHOW ANY SIGNS OF TAMPERING.
Inactive ingredients corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, talc, titanium dioxide
? call 1-800-406-7984

Drug Facts
Active ingredient (in each tablet) Cetirizine HCl 5 mg
Antihistamine
Uses relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings (continued)
Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms:
■ trouble swallowing ■ dizziness or loss of consciousness ■ swelling of tongue ■ difficulty breathing
■ trouble speaking ■ wheezing or problems breathing ■ drooling
immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.
If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergic symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.
Do not use ■ to prevent hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves ■ because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
Ask a doctor before use if you have ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch ■ liver or kidney disease. Your doctor should determine if you need a different dose.
When using this product ■ drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ symptoms do not improve after 3 days of treatment ■ the hives have lasted more than 6 weeks ■ an allergic reaction to this product occurs. Seek medical help right away.
Stop use and ask a doctor if ■ If pregnant or breast-feeding: not recommended ■ If pregnant, ask a health professional before use ■ If breast-feeding: not recommended ■ If pregnant, ask a health professional before use.
Directions adults and children 6 years and over ■ 1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 2 tablets in 24 hours
adults 65 years and over ■ 1 tablet once a day; do not take more than 1 tablet in 24 hours
ask a doctor
children under 6 years of age ■ ask a doctor
consumers with liver or kidney disease ■ ask a doctor

Standard Format
Helvetica Medium Condensed
Variable horizontal scale
adheres to 39 characters per inch
Drug Facts - 11 pt. Helv. Bold Cond. Italic, cond 90%
Drug Facts (continued)
Drug Facts - 9pt Helv. Bold Cond. Italic, cond 90%
(continued) - 7pt. Helvetica, cond 85%
Headings - 6pt. Helv. Bld. Cond. Italic, cond 90%
Sub-Heads - 6pts. Helv. Bold, Cond 90%
Text 6pt. Helv., Med Cond. 90%, min. leading 6.5
Square bullets - 5pt. Zapf Ding., cond 85%
Border 2.5 pt. rule, 1/8 gap all around
Barline 2.0pt. rule
Headline .5 pt. rule

Original Prescription Strength

Cetirizine HCl Tablets, 10 mg

Antihistamine Allergy

Indoor & Outdoor Allergies

24 Hour Relief of

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

30 TABLETS
10 mg EACH

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.

**Active ingredient
(in each tablet)**
Cetirizine HCl 10 mg.....Antihistamine

Purpose
Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat.

Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor before use if you are taking tranquilizers or sedatives.** Alcohol, sedatives, and tranquilizers may increase drowsiness. ■ be careful when driving a motor vehicle or operating machinery. **Stop use if you experience dizziness, headache, or other adverse effects. Seek medical help right away. If pregnant or breast-feeding:** ■ If breast-feeding, not recommended. ■ If pregnant, ask a health professional before use. **Keep out of reach of children.** For more information, contact a Poison Control Center right away.

Directions Adults and children 6 years and over: one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. Ask a health professional for help if symptoms do not improve or get worse or if you have less severe symptoms. **Adults 65 years and over:** ask a doctor. **Children under 6 years of age:** ask a doctor. **Consumers with liver or kidney disease:** ask a doctor.

Other information ■ store between 20° to 25° C (68° to 77° F). ■ see USP Controlled Room Temperature. ■ see USP Full Protection from Light.

Inactive ingredients See caution for complete list of ingredients.

Questions? call 1-800-406-7884

XXXX
Distributed by:
Ohm Laboratories, Inc.
14 Terminal Road
New Brunswick, NJ 08901

Batch No. XXXXXX
Expiration Date: XXXXXX

Non Varnish Area

Standard Format
Helvetica Condensed
Variable horizontal scale
adheres to 39 characters per inch

Active ingredient - 5.63 pt. Helv. Bold Oblique,
Uses - 5.1 pt. Helvetica Cnd. Bold Italic
Headings - 5.1 pt. Helv. Cnd. Bld. Italic
Sub Heads - 4.08 pt. Helv. Cnd. Bold,
Text - 4.08 pt. Helv. Cnd., min. leading 4.21,
horiz 106.86%

Square Bullets - 3.4 pt. Zapf Ding.,
Border 1 pt. rule, 1/16 gap all around
Barline 1pt. rule _____
Hairline .5 pt. rule _____

Original Prescription Strength

Cetirizine HCl Tablets, 10 mg

Antihistamine Allergy

Indoor & Outdoor Allergies

24 Hour Relief of

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

45 TABLETS
10 mg EACH

TAMPER EVIDENT. DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.

Active Ingredient
(in each Tablet)

Cetirizine HCl 10 mg

Purpose
Cetirizine HCl 10 mg relieves these symptoms of allergic rhinitis or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat.

Warnings: Do not use if you have ever had an allergic reaction to cetirizine HCl or any other antihistamine containing hydroxyzine. **Ask a doctor before use** if you have liver or kidney disease. Your doctor should use caution if you are taking other antihistamines or sedatives. **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and other drugs may increase drowsiness. **Stop use and ask a doctor** if an allergic reaction to this product occurs. **Do not drive or operate machinery** if you feel drowsy. **Do not use** if pregnant, nursing, or trying to become pregnant; ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.


Directions: Adults and children 12 years of age and over: One 10 mg Tablet once daily, do not take more than one 10 mg Tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. **Adults 65 years and over:** ask a health professional before use. **Children:** ask a doctor. **Keep out of reach of children.** **Other Information** ■ store between 20° to 25° C (68° to 77° F).

Inactive Ingredients See carton for complete list of ingredients.

Questions? call 1-800-406-7984

XXXX

Distributed by:
Oryz Laboratories Inc.
1400 Old Bridge Road
New Brunswick, NJ 08901



XXXXXXX
XXXXXXX

Batch No. _____

Expiration Date: _____

Non Varnish Area

Standard Format
Helvetica Condensed
Variable horizontal scale
adheres to 39 characters per inch

Active ingredient - 5.63 pt. Helv. Bold Oblique,

Uses - 5.1 pt. Helvetica Cnd. Bold Italic

Headings - 5.1 pt. Helv. Cnd. Bld. Italic

Sub Heads - 4.08 pt. Helv. Cnd. Bold,

Text - 4.08 pt. Helv. Cnd., min. leading 4.21,
horiz 106.86%

Square Bullets - 3.4 pt. Zapf Ding.,

Border 1 pt. rule, 1/16 gap all around

Barline 1pt. rule _____

Hairline .5 pt. rule _____

Distributed by:
 Ohm Laboratories Inc.
 14 Terminal Road
 New Brunswick, NJ 08901



Original Prescription Strength

Cetirizine HCl Tablets, 10 mg Antihistamine

Allergy
 Indoor & Outdoor Allergies

24 Hour Relief of
 Sneezing
 Runny Nose
 Itchy, Watery Eyes
 Itchy Throat or Nose

75 Tablets
10 mg EACH
 (contains one bottle of 45 tablets
 and one bottle of 30 tablets)

Batch No. _____ Expiration Date: _____

Non Varnish Area

Cetirizine HCl Tablets, 10 mg
Antihistamine
Allergy
 Indoor & Outdoor Allergies

Drug Facts
Active ingredient (in each tablet)
 Cetirizine HCl 10 mg.....Antihistamine

Uses
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

Drug Facts (continued)
 When using this product
 ■ drowsiness may occur
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding: ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 adults and children 6 years and over
 one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.

ask a doctor	adults 65 years and over
ask a doctor	children under 6 years of age
ask a doctor	consumers with liver or kidney disease

Other information
 ■ TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.
 ■ store between 20° to 25° C (68° to 77° F)

Inactive ingredients corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, talc, titanium dioxide

Questions? call 1-800-406-7984

Keep the carton. It contains important information.
 See end panel for expiration date.



Standard Format
 Tablets - 10 mg scale
 adheres to 39 characters per inch

Drug Facts - 11 pt. HeliV Bold Italic,
 Drug Facts Continued - 9 pt. HeliV Bold Italic,
 Headings - 8pt. HeliV Bld. Italic,
 Sub heads - 6pt. HeliV Bld.
 Text 6pt. HeliV Bld. min. leading 6.5
 Square bullets - 5 pt. Zap Ding.
 Border 2.5 pt. rule, 1/8 gap all around
 Baseline 2.5pt. rule
 Hairline .5 pt. rule

Original Prescription Strength

Cetirizine HCl Tablets, 10 mg Antihistamine Hives Relief

24 Hour Relief of
Itching Due to Hives

30 TABLETS
10 mg EACH

**TAMPER EVIDENT: DO NOT USE IF IMPROPER
SEAL IS BROKEN OR MISSING FROM BOTTLE.**

Active ingredient (in each tablet)

Cetirizine HCl, 10 mg.....Antihistamine
Other inactive ingredients in this product
will not prevent hives or an allergic skin reaction from
occurring.

Purpose

Relief of itching due to hives or an allergic skin reaction from occurring.

Warnings/Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following: trouble breathing, swelling of the face, lips, tongue or throat, dizziness or loss of consciousness, swelling in or around mouth, trouble speaking, drooling, wheezing or around mouth, trouble breathing, wheezing, or a problem swallowing. These symptoms may be the signs of a severe allergic reaction. Do not use this product if you are not treated by a health professional immediately.

Epinephrine: If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. You should carry it with you at all times. Do not use to prevent hives from any other cause. If you have ever had an allergic reaction to this product or any of its ingredients or to an epinephrine injector, do not use this product. Do not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be prevented by using antihistamines. Your doctor may be able to help you find a cause. If you have ever had an allergic reaction to this product or any of its ingredients or to an epinephrine injector, do not use this product. Your doctor should determine if you need a different dose. Hives that do not itch. Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives. When using this product, drowsiness may occur. Avoid alcoholic drinks and grapefruit juice. Do not drink alcohol while driving or operating machinery. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help if you experience any of the following symptoms: If pregnant or breast-feeding: If breast-feeding, not known if this drug passes into breast milk. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions: Adults and children 6 years and over: one (1) tablet twice a day with water. For best results, take 1 hour before bedtime. A 5 mg product may be appropriate for less severe symptoms. Adults 65 years and over: ask a doctor. Children under 6 years: ask a doctor. Consumers with liver or kidney disease: ask a doctor.

Other information: Store between 20° to 25° C (68° to 77° F).

Inactive ingredients: See carton for complete list of ingredients.

Questions? call 1-800-406-7964

XXXX

Distributed by
Omnilabs Laboratories, Inc.
14 Terminal Road
New Brunswick, NJ 08901

Expiration Date: XXXXXX

Batch No. XXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX

XXXXXXXXXX



Standard Format
 Helvetica Condensed
 Variable horizontal scale
 adheres to 39 characters per inch

Active ingredient - 5.63 pt. Helv. Bold Oblique,
 Uses - 5.1 pt. Helvetica Cnd. Bold Italic
 Headings - 5.1 pt. Helv. Cnd. Bld. Italic
 Sub Heads - 4.08 pt. Helv. Cnd. Bold,
 Text - 4.08 pt. Helv. Cnd., min. leading 4.21,
 horiz 106.86%

Square Bullets - 3.4 pt. Zapf Ding.,
 Border 1 pt. rule, 1/16 gap all around
 Barline 1pt. rule _____
 Hairline .5 pt. rule _____



Distributed by:
 Ohm Laboratories Inc.
 14 Terminal Road
 New Brunswick, NJ 08901

Original Prescription Strength

Cetirizine HCl

Tablets, 10 mg

Antihistamine

Hives Relief

24 Hour Relief of Itching Due to Hives

75 Tablets
10 mg EACH
(contains one bottle of 45 tablets and one bottle of 30 tablets)

Batch No. _____ Expiration Date: _____

Non Varnish Area

Drug Facts (continued)
 Inactive ingredients corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, talc, titanium dioxide

Questions? call 1-800-406-7984

Keep the carton. It contains important information.
 See end panel for expiration date.

Drug Facts (in each tablet)
 Cetirizine HCl 10 mg, Antihistamine

Uses relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings
 Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms:
 ■ trouble swallowing ■ dizziness or loss of consciousness ■ swelling of tongue ■ swelling in or around mouth ■ trouble speaking ■ wheezing or problems breathing ■ drooling

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

Drug Facts (continued)
 Not a substitute for epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

Do not use ■ to prevent hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves ■ because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.

■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. ■ hives that do not itch ■ hives that are an unusual color, look bruised or blistered ■ liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives. ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ symptoms do not improve after 3 days of treatment ■ the hives have lasted more than 6 weeks ■ an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding: ■ If breast-feeding, not recommended ■ If pregnant, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 adults and children 6 years and over one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours.
 adults 65 years and over ask a doctor
 children under 6 years of age ask a doctor
 ask a doctor
 ask a doctor
 ask a doctor
 ask a doctor

Other information
 ■ store between 20° to 25° C (68° to 77° F)
 ■ TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.

Standard Format
 Vertical Orientation
 Vertical Orientation
 Vertical Orientation
 adheres to 39 characters per inch

Drug Facts - 11 pt. Hxlv, Bold Italic,
 Drug Facts Continued - 9 pt. Hxlv, Bold Italic,
 Headings - 8pt. Hxlv, Bld, Italic,
 Sub heads - 6pts. Hxlv, Bld,
 Text - 6pt. Hxlv, Bld, min. leading 6.5
 Square Bullets - 5 pt. Zap Ding,
 Border 2.5 pt. rule, 1/8 gap all around
 Baseline 2.5pt. rule
 Hairline .5 pt. rule