

NDC 0228-1196-55



**Cetirizine**  
Hydrochloride  
Tablets **5 mg**  
Antihistamine  
Allergy

Indoor & Outdoor Allergies

100 tablets - 5 mg each

Do not use if imprinted foil inner seal on bottle is broken or missing.

**Active ingredient (in each tablet)**  
Cetirizine HCl 5 mg

**Purpose**  
Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

**Warnings Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Rev. 11/07  
Mfg by: Saurus Chemical & Drugs Limited,  
Unit IV, R.S. No. 32, 33, & 34, PIMS Road, Periyakalpet,  
Pondicherry - 605014, India  
For: Actavis Elizabeth LLC, 200 Elmora Avenue  
Elizabeth, NJ 07207 USA

**PEEL HERE** ▶

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives.

**When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor if an allergic reaction** to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding:** ■ if breast-feeding not recommended ■ if pregnant, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ **adults and children 6 years and over:** 1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 2 tablets in 24 hours. ■ **adults 65 years and over:** 1 tablet once a day; do not take more than 1 tablet in 24 hours. ■ **children under 6 years of age:** ask a doctor

■ **consumers with liver or kidney disease:** ask a doctor

**Other information** ■ store between 20° to 25° C (68° to 77° F)

■ do not use if imprinted foil inner seal on bottle is broken or missing.

**Inactive ingredients** colloidal silicon dioxide, croscopolone, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, triacetin and titanium dioxide.

**Questions?** call 1-800-432-8534

Rev. 11/07  
Mfg by: Saurus Chemical & Drugs Limited,  
Unit IV, R.S. No. 32, 33, & 34, PIMS Road, Periyakalpet,  
Pondicherry - 605014, India  
For: Actavis Elizabeth LLC, 200 Elmora Avenue  
Elizabeth, NJ 07207 USA

NDC 0228-1198-55



**Cetirizine**  
Hydrochloride  
Tablet **10 mg**  
Antihistamine  
**Allergy**  
Indoor & Outdoor Allergies  
100 tablets - 10 mg each

Do not use if imprinted foil inner seal on bottle is broken or missing.

**Active ingredient (in each tablet)**  
Cetirizine HCl 10 mg . . . . . Antihistamine

**Purpose**  
**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

**Warnings** Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Rev. 11/07

Mfg by: Shasun Chemical & Drugs Limited,  
Unit 11, S.S. No. 32, 33, & 34, PMS Road, Periyakalpet,  
Pondicherry - 605014, India  
For: Actavis Elizabeth LLC, 200 Elmora Avenue  
Elizabeth, NJ 07201 USA

**PEEL HERE** ▶

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives.

**When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding:** ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ **adults and children 6 years and over:** one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.

■ **adults 65 years and over:** ask a doctor ■ **children under 6 years of age:** ask a doctor

■ **consumers with liver or kidney disease:** ask a doctor

**Other information** ■ store between 20° to 25° C (68° to 77° F)

■ do not use if imprinted foil inner seal on bottle is broken or missing.

**Inactive ingredients** colloidal silicon dioxide, croscopvidone, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, triacetin and titanium dioxide.

**Questions?** call 1-800-432-8534

Rev. 11/07

Mfg by: Shasun Chemical & Drugs Limited,  
Unit 11, S.S. No. 32, 33, & 34, PMS Road, Periyakalpet,  
Pondicherry - 605014, India  
For: Actavis Elizabeth LLC, 200 Elmora Avenue  
Elizabeth, NJ 07201 USA

Indoor & Outdoor Allergies  
**Allergy**  
 Antihistamine  
 Tablets **5 mg**  
**Cetirizine**  
 Hydrochloride  
 actavis

**Drug Facts** (continued)

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if an allergic reaction to this product occurs.** Seek medical help right away.

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

\*This product is not manufactured or distributed by Johnson & Johnson, distributor of Zyrtec®.  
 Rev. 11/07  
 actavis  
 Mfd by: Shasun Chemical & Drugs Limited, Unit II, 65, No: 32, 33, & 34, PIMS Road, Periyakalpet, Pondicherry - 605014, India  
 For: Actavis Elizabeth LLC, 200 Elmora Avenue Elizabeth, NJ 07207 USA

**Drug Facts** (continued)

**Directions**

adults and children 6 years and over	1 to 2 tablets once daily depending upon severity of symptoms; do not take more than 2 tablets in 24 hours
adults 65 years and over	1 tablet once a day; do not take more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Other information**

- Store between 20° to 25°C (68° to 77°F)
- do not use if imprinted foil inner seal on bottle is broken or missing.

**Inactive ingredients** colloidal silicon dioxide, crospovidone, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, triacetin and titanium dioxide.

**NEW** Original Prescription Strength  
 NDC 0228-1196-55  
 actavis  
 Compare to the active ingredient in Zyrtec® Allergy\*  
**Cetirizine**  
**5 mg**  
**Allergy**  
 Indoor & Outdoor Allergies  
 • Sneezing • Runny Nose  
 • Itchy, Watery Eyes  
 • Itchy Throat or Nose  
 100 tablets - 5 mg each

**Drug Facts**

<b>Active ingredient (in each tablet)</b> Cetirizine HCl 5 mg	<b>Purpose</b> Antihistamine
--	---------------------------------

**Uses**  
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**  
**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives.



Indoor & Outdoor Allergies  
**Allergy**  
 Antihistamine  
 Tablets **10 mg**  
**Cetirizine**  
 Hydrochloride  
 actavis

**Drug Facts** (continued)

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if an allergic reaction to this product occurs.** Seek medical help right away.

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

\*This product is not manufactured or distributed by Johnson & Johnson, distributor of Zyrtec®.

actavis  
 Mfd by: Shasun Chemical & Drugs Limited,  
 Unit II, R.S. No: 32, 33, & 34, PIMS Road, Periyakalpet,  
 Pondicherry - 605014, India  
 For: Actavis Elizabeth LLC, 200 Elmora Avenue  
 Elizabeth, NJ 07207 USA

Rev. 11/07

**Drug Facts** (continued)

**Directions**

adults and children 6 years and over	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
adults 65 years and over	ask a doctor
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Other information**

- Store between 20° to 25°C (68° to 77°F)
- do not use if imprinted foil inner seal on bottle is broken or missing.

**Inactive ingredients**

colloidal silicon dioxide, crospovidone, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, triacetin and titanium dioxide.

**NEW** Original Prescription Strength

NDC 0228-1198-55



Compare to the active ingredient in Zyrtec® Allergy\*

**Cetirizine**

**10 mg**

**Allergy**

Indoor & Outdoor Allergies

- Sneezing • Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

100 tablets - 10 mg each

**Drug Facts**

**Active ingredient Purpose (in each tablet)**  
 Cetirizine HCl 10 mg ..... Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies

- runny nose ■ sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives.



3 0228-1198-55 2

Indoor & Outdoor Allergies

**Allergy**

Antihistamine

Tablets **10 mg**

**Cetirizine**

actavis

**NEW** Original Prescription Strength

NDC 0228-1198-57

Compare to the active ingredient in Zyrtec® Allergy\*

actavis

**Cetirizine**


**10 mg**

**Allergy**

Indoor & Outdoor Allergies

- Sneezing • Runny Nose
- Itchy, Watery Eyes • Itchy Throat or Nose

200 tablets 10 mg each – Contains 2 bottles of 100 tablets each



3 0228-1198-57 6

**Drug Facts**

**Active ingredient (in each tablet)**  
Cetirizine HCl 10 mg ..... Antihistamine

**Uses**  
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  
• runny nose • sneezing  
• itchy, watery eyes  
• itching of the nose or throat

**Warnings**  
**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Asks doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**Asks doctor or pharmacist before use if you are** taking tranquilizers or sedatives.

**Drug Facts (continued)**

**When using this product**  
• drowsiness may occur  
• avoid alcoholic drinks  
• alcohol, sedatives, and tranquilizers may increase drowsiness  
• be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if an allergic reaction to this product occurs.** Seek medical help right away.  
**If pregnant or breast-feeding:**  
• if breast-feeding: not recommended  
• if pregnant: ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

adults and children 6 years and over	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
adults 65 years and over	ask a doctor
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Drug Facts (continued)**

**Other information**

- Store between 20° to 25°C (68° to 77°F)
- do not use if printed foil inner seal on bottle is broken or missing.

**Inactive ingredients** colloidal silicon dioxide, crospovidone, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, triacetin and titanium dioxide.

\*This product is not manufactured or distributed by Johnson & Johnson, distributor of Zyrtec®.  
Rev. 11/07



Mid by Shasun Chemical & Drugs Limited,  
Unit 11, P.S. No. 32, 33 & 34, PIMS Road, Periyakalpet,  
Pondicherry – 605014, India  
PMS Health Care, LLC, 200 Elmora Avenue  
Elizabeth, NJ 07207 USA