


TABLETS
NOROXIN®
(NORFLOXACIN)

To reduce the development of drug-resistant bacteria and maintain the effectiveness of NOROXIN† and other antibacterial drugs, NOROXIN should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

DESCRIPTION

NOROXIN (Norfloxacin) is a synthetic, broad-spectrum antibacterial agent for oral administration. Norfloxacin, a fluoroquinolone, is 1-ethyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid. Its empirical formula is $C_{16}H_{18}FN_3O_3$ and the structural formula is:



Norfloxacin is a white to pale yellow crystalline powder with a molecular weight of 319.34 and a melting point of about 221°C. It is freely soluble in glacial acetic acid, and very slightly soluble in ethanol, methanol and water.

NOROXIN is available in 400-mg tablets. Each tablet contains the following inactive ingredients: cellulose, croscarmellose sodium, hydroxypropyl cellulose, hydroxypropyl methylcellulose, magnesium stearate, and titanium dioxide.

Norfloxacin, a fluoroquinolone, differs from non-fluorinated quinolones by having a fluorine atom at the 6 position and a piperazine moiety at the 7 position.

CLINICAL PHARMACOLOGY

In fasting healthy volunteers, at least 30-40% of an oral dose of NOROXIN is absorbed. Absorption is rapid following single doses of 200 mg, 400 mg and 800 mg. At the respective doses, mean peak serum and plasma concentrations of 0.8, 1.5 and 2.4 $\mu\text{g/mL}$ are attained approximately one hour after dosing. The presence of food and/or dairy products may decrease absorption. The effective half-life of norfloxacin in serum and plasma is 3-4 hours. Steady-state concentrations of norfloxacin will be attained within two days of dosing.

In healthy elderly volunteers (65-75 years of age with normal renal function for their age), norfloxacin is eliminated more slowly because of their slightly decreased renal function. Following a single 400-mg dose of norfloxacin, the mean (\pm SD) AUC and C_{max} of 9.8 (2.83) $\mu\text{g}\cdot\text{hr/mL}$ and 2.02 (0.77) $\mu\text{g/mL}$, respectively, were observed in healthy elderly volunteers. The extent of systemic exposure was slightly higher than that seen in younger adults (AUC 6.4 $\mu\text{g}\cdot\text{hr/mL}$ and C_{max} 1.5 $\mu\text{g/mL}$). Drug absorption appears unaffected. However, the effective half-life of norfloxacin in these elderly subjects is 4 hours.

There is no information on accumulation of norfloxacin with repeated administration in elderly patients. However, no dosage adjustment is required based on age alone. In elderly patients with reduced renal function, the dosage should be adjusted as for other patients with renal impairment (see DOSAGE AND ADMINISTRATION, *Renal Impairment*).

The disposition of norfloxacin in patients with creatinine clearance rates greater than 30 mL/min/1.73 m² is similar to that in healthy volunteers. In patients with creatinine clearance rates equal to or less than 30 mL/min/1.73 m², the renal elimination of norfloxacin decreases so that the

† Registered trademark of MERCK & CO., Inc.
COPYRIGHT © 1986, 1989, 1999, 2001 MERCK & CO., Inc.
All rights reserved

effective serum half-life is 6.5 hours. In these patients, alteration of dosage is necessary (see DOSAGE AND ADMINISTRATION). Drug absorption appears unaffected by decreasing renal function.

Norfloxacin is eliminated through metabolism, biliary excretion, and renal excretion. After a single 400-mg dose of NOROXIN, mean antimicrobial activities equivalent to 278, 773, and 82 µg of norfloxacin/g of feces were obtained at 12, 24, and 48 hours, respectively. Renal excretion occurs by both glomerular filtration and tubular secretion as evidenced by the high rate of renal clearance (approximately 275 mL/min). Within 24 hours of drug administration, 26 to 32% of the administered dose is recovered in the urine as norfloxacin with an additional 5-8% being recovered in the urine as six active metabolites of lesser antimicrobial potency. Only a small percentage (less than 1%) of the dose is recovered thereafter. Fecal recovery accounts for another 30% of the administered dose. In elderly subjects (average creatinine clearance 91 mL/min/1.73 m²) approximately 22% of the administered dose was recovered in urine and renal clearance averaged 154 mL/min.

Two to three hours after a single 400-mg dose, urinary concentrations of 200 µg/mL or more are attained in the urine. In healthy volunteers, mean urinary concentrations of norfloxacin remain above 30 µg/mL for at least 12 hours following a 400-mg dose. The urinary pH may affect the solubility of norfloxacin. Norfloxacin is least soluble at urinary pH of 7.5 with greater solubility occurring at pHs above and below this value. The serum protein binding of norfloxacin is between 10 and 15%.

The following are mean concentrations of norfloxacin in various fluids and tissues measured 1 to 4 hours post-dose after two 400-mg doses, unless otherwise indicated:

| | |
|------------------|---------------------------------------|
| Renal Parenchyma | 7.3 µg/g |
| Prostate | 2.5 µg/g |
| Seminal Fluid | 2.7 µg/mL |
| Testicle | 1.6 µg/g |
| Uterus/Cervix | 3.0 µg/g |
| Vagina | 4.3 µg/g |
| Fallopian Tube | 1.9 µg/g |
| Bile | 6.9 µg/mL (after two 200-mg doses) |

Microbiology

Norfloxacin has *in vitro* activity against a broad range of gram-positive and gram-negative aerobic bacteria. The fluorine atom at the 6 position provides increased potency against gram-negative organisms, and the piperazine moiety at the 7 position is responsible for antipseudomonal activity.

Norfloxacin inhibits bacterial deoxyribonucleic acid synthesis and is bactericidal. At the molecular level, three specific events are attributed to norfloxacin in *E. coli* cells:

- 1) inhibition of the ATP-dependent DNA supercoiling reaction catalyzed by DNA gyrase,
- 2) inhibition of the relaxation of supercoiled DNA,
- 3) promotion of double-stranded DNA breakage.

Resistance to norfloxacin due to spontaneous mutation *in vitro* is a rare occurrence (range: 10⁻⁹ to 10⁻¹² cells). Resistant organisms have emerged during therapy with norfloxacin in less than 1% of patients treated. Organisms in which development of resistance is greatest are the following:

Pseudomonas aeruginosa

Klebsiella pneumoniae

Acinetobacter spp.

Enterococcus spp.

For this reason, when there is a lack of satisfactory clinical response, repeat culture and susceptibility testing should be done. Nalidixic acid-resistant organisms are generally susceptible to norfloxacin *in vitro*; however, these organisms may have higher minimum inhibitory concentrations (MICs) to norfloxacin than nalidixic acid-susceptible strains. There is generally no cross-resistance between norfloxacin and other classes of antibacterial agents. Therefore, norfloxacin may demonstrate activity against indicated organisms resistant to some other antimicrobial agents including the aminoglycosides, penicillins, cephalosporins, tetracyclines, macrolides, and sulfonamides, including combinations of sulfamethoxazole and trimethoprim. Antagonism has been demonstrated *in vitro* between norfloxacin and nitrofurantoin.

Norfloxacin has been shown to be active against most strains of the following microorganisms both *in vitro* and in clinical infections as described in the **INDICATIONS AND USAGE** section.

Gram-positive aerobes:

Enterococcus faecalis

Staphylococcus aureus

Staphylococcus epidermidis
Staphylococcus saprophyticus
Streptococcus agalactiae

Gram-negative aerobes:

Citrobacter freundii
Enterobacter aerogenes
Enterobacter cloacae
Escherichia coli
Klebsiella pneumoniae
Neisseria gonorrhoeae
Proteus mirabilis
Proteus vulgaris
Pseudomonas aeruginosa
Serratia marcescens

The following *in vitro* data are available, **but their clinical significance is unknown.**

Norfloxacin exhibits *in vitro* MICs of ≤ 4 $\mu\text{g/mL}$ against most ($\geq 90\%$) strains of the following microorganisms; however, the safety and effectiveness of norfloxacin in treating clinical infections due to these microorganisms have not been established in adequate and well-controlled clinical trials.

Gram-negative aerobes:

Citrobacter diversus
Edwardsiella tarda
Enterobacter agglomerans
Haemophilus ducreyi
Klebsiella oxytoca
Morganella morganii
Providencia alcalifaciens
Providencia rettgeri
Providencia stuartii
Pseudomonas fluorescens
Pseudomonas stutzeri

Other:

Ureaplasma urealyticum

NOROXIN is not generally active against obligate anaerobes.

Norfloxacin has not been shown to be active against *Treponema pallidum* (see WARNINGS).

Susceptibility Tests**Dilution Techniques:**

Quantitative methods are used to determine antimicrobial MICs. These MICs provide estimates of the susceptibility of bacteria to antimicrobial compounds. The MICs should be determined using a standardized procedure. Standardized procedures are based on a dilution method¹ (broth, agar, or microdilution) or equivalent with standardized inoculum concentrations and standardized concentrations of norfloxacin powder. The MIC values should be interpreted according to the following criteria*:

| MIC ($\mu\text{g/mL}$) | Interpretation |
|--------------------------|------------------|
| ≤ 4 | Susceptible (S) |
| 8 | Intermediate (I) |
| ≥ 16 | Resistant (R) |

A report of "Susceptible" indicates that the pathogen is likely to be inhibited if the antimicrobial compound in the blood reaches the concentrations usually achievable. A report of "Intermediate" indicates that the result should be considered equivocal, and, if the microorganism is not fully susceptible to alternative, clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where high dosage of drug can be used. This category also provides a buffer zone which prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of "Resistant" indicates that

* These interpretative criteria apply only to isolates from urinary tract infections. There are no established norfloxacin interpretive criteria for *Neisseria gonorrhoeae* or organisms isolated from other infection sites.

the pathogen is not likely to be inhibited if the antimicrobial compound in the blood reaches the concentrations usually achievable; other therapy should be selected.

Standardized susceptibility test procedures require the use of laboratory control microorganisms to control the technical aspects of the laboratory procedures. Standard norfloxacin powder should provide the following MIC values:

| <u>Organism</u> | <u>MIC range (µg/mL)</u> |
|---------------------------------|--------------------------|
| <i>E. coli</i> ATCC 25922 | 0.03-0.12 |
| <i>E. faecalis</i> ATCC 29212 | 2-8 |
| <i>P. aeruginosa</i> ATCC 27853 | 1-4 |
| <i>S. aureus</i> ATCC 29213 | 0.5-2 |

Diffusion Techniques:

Quantitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. One such standardized procedure² requires the use of standardized inoculum concentrations. This procedure uses paper disks impregnated with 10-µg norfloxacin to test the susceptibility of microorganisms to norfloxacin. Reports from the laboratory providing results of the standard single-disk susceptibility test with a 10-µg norfloxacin disk should be interpreted according to the following criteria*:

| <u>Zone diameter (mm)</u> | <u>Interpretation</u> |
|---------------------------|-----------------------|
| ≥17 | Susceptible (S) |
| 13-16 | Intermediate (I) |
| ≤12 | Resistant (R) |

Interpretation should be as stated above for results using dilution techniques. Interpretation involves correlation of the diameter obtained in the disk test with the MIC for norfloxacin.

As with standard dilution techniques, diffusion methods require the use of laboratory control microorganisms that are used to control the technical aspects of the laboratory procedures. For the diffusion techniques, the 10-µg norfloxacin disk should provide the following zone diameters in these laboratory test quality control strains:

| <u>Organism</u> | <u>Zone Diameter (mm)</u> |
|---------------------------------|---------------------------|
| <i>E. coli</i> ATCC 25922 | 28-35 |
| <i>P. aeruginosa</i> ATCC 27853 | 22-29 |
| <i>S. aureus</i> ATCC 25923 | 17-28 |

INDICATIONS AND USAGE

NOROXIN is indicated for the treatment of adults with the following infections caused by susceptible strains of the designated microorganisms:

Urinary tract infections:

Uncomplicated urinary tract infections (including cystitis) due to *Enterococcus faecalis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, *Citrobacter freundii*^{**}, *Enterobacter aerogenes*^{**}, *Enterobacter cloacae*^{**}, *Proteus vulgaris*^{**}, *Staphylococcus aureus*^{**}, or *Streptococcus agalactiae*^{**}.

Complicated urinary tract infections due to *Enterococcus faecalis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, or *Serratia marcescens*^{**}.

Sexually transmitted diseases (see WARNINGS):

Uncomplicated urethral and cervical gonorrhea due to *Neisseria gonorrhoeae*.

Prostatitis:

Prostatitis due to *Escherichia coli*.

(See DOSAGE AND ADMINISTRATION for appropriate dosing instructions.)

Penicillinase production should have no effect on norfloxacin activity.

Appropriate culture and susceptibility tests should be performed before treatment in order to isolate and identify organisms causing the infection and to determine their susceptibility to norfloxacin. Therapy with norfloxacin may be initiated before results of these tests are known; once results become available, appropriate therapy should be given. Repeat culture and susceptibility testing performed periodically during therapy will provide information not only on the therapeutic effect of the antimicrobial agents but also on the possible emergence of bacterial resistance.

^{**} Efficacy for this organism in this organ system was studied in fewer than 10 infections.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of NOROXIN and other antibacterial drugs, NOROXIN should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

CONTRAINDICATIONS

NOROXIN (norfloxacin) is contraindicated in persons with a history of hypersensitivity, tendinitis, or tendon rupture associated with the use of norfloxacin or any member of the quinolone group of antimicrobial agents.

WARNINGS

Safety in Children, Adolescents, Nursing mothers, and during Pregnancy: THE SAFETY AND EFFICACY OF ORAL NORFLOXACIN IN PEDIATRIC PATIENTS, ADOLESCENTS (UNDER THE AGE OF 18), PREGNANT WOMEN, AND NURSING MOTHERS HAVE NOT BEEN ESTABLISHED. (See PRECAUTIONS, Pediatric Use, Pregnancy, and Nursing Mothers subsections.) The oral administration of single doses of norfloxacin, 6 times*** the recommended human clinical dose (on a mg/kg basis), caused lameness in immature dogs. Histologic examination of the weight-bearing joints of these dogs revealed permanent lesions of the cartilage. Other quinolones also produced erosions of the cartilage in weight-bearing joints and other signs of arthropathy in immature animals of various species (see ANIMAL PHARMACOLOGY).

Seizures: Convulsions have been reported in patients receiving norfloxacin. Convulsions, increased intracranial pressure, and toxic psychoses have been reported in patients receiving drugs in this class. Quinolones may also cause central nervous system (CNS) stimulation which may lead to tremors, restlessness, lightheadedness, confusion, and hallucinations. If these reactions occur in patients receiving norfloxacin, the drug should be discontinued and appropriate measures instituted.

The effects of norfloxacin on brain function or on the electrical activity of the brain have not been tested. Therefore, until more information becomes available, norfloxacin, like all other quinolones, should be used with caution in patients with known or suspected CNS disorders, such as severe cerebral arteriosclerosis, epilepsy, and other factors which predispose to seizures (see ADVERSE REACTIONS).

Hypersensitivity Reactions: Serious and occasionally fatal hypersensitivity (anaphylactic) reactions, some following the first dose, have been reported in patients receiving quinolone therapy, including NOROXIN. Some reactions were accompanied by cardiovascular collapse, loss of consciousness, tingling, pharyngeal or facial edema, dyspnea, urticaria and itching. Only a few patients had a history of hypersensitivity reactions. If an allergic reaction to norfloxacin occurs, discontinue the drug. Serious acute hypersensitivity reactions require immediate emergency treatment with epinephrine. Oxygen, intravenous fluids, antihistamines, corticosteroids, pressor amines, and airway management, including intubation, should be administered as indicated.

Other serious and sometimes fatal events, some due to hypersensitivity, and some due to uncertain etiology, have been reported rarely in patients receiving therapy with quinolones, including NOROXIN. These events may be severe and generally occur following the administration of multiple doses. Clinical manifestations may include one or more of the following:

- fever, rash or severe dermatologic reactions (e.g., toxic epidermal necrolysis, Stevens-Johnson syndrome);
- vasculitis; arthralgia; myalgia; serum sickness;
- allergic pneumonitis;
- interstitial nephritis; acute renal insufficiency or failure;
- hepatitis; jaundice; acute hepatic necrosis or failure;
- anemia, including hemolytic and aplastic; thrombocytopenia, including thrombotic thrombocytopenic purpura; leukopenia; agranulocytosis; pancytopenia; and/or other hematologic abnormalities.

*** Based on a patient weight of 50 kg.

The drug should be discontinued immediately at the first appearance of a skin rash, jaundice, or any other sign of hypersensitivity, and supportive measures should be instituted (see PRECAUTIONS, *Information for Patients* and ADVERSE REACTIONS).

Clostridium difficile associated diarrhea: *Clostridium difficile* associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including NOROXIN and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD.

Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

Peripheral neuropathy: Rare cases of sensory or sensorimotor axonal polyneuropathy affecting small and/or large axons resulting in paresthesias, hypoesthesias, dysesthesias and weakness have been reported in patients receiving quinolones, including norfloxacin. Norfloxacin should be discontinued if the patient experiences symptoms of neuropathy including pain, burning, tingling, numbness, and/or weakness, or is found to have deficits in light touch, pain, temperature, position sense, vibratory sensation, and/or motor strength in order to prevent the development of an irreversible condition.

Tendon effects: Ruptures of the shoulder, hand, Achilles tendons or other tendons that required surgical repair or resulted in prolonged disability have been reported in patients receiving quinolones, including norfloxacin. Post-marketing surveillance reports indicate that this risk may be increased in patients receiving concomitant corticosteroids, especially in the elderly. Norfloxacin should be discontinued if the patient experiences pain, inflammation, or rupture of a tendon. Patients should rest and refrain from exercise until the diagnosis of tendinitis or tendon rupture has been excluded. Tendon rupture can occur during or after therapy with quinolones, including norfloxacin.

Syphilis treatment: Norfloxacin has **not** been shown to be effective in the treatment of syphilis. Antimicrobial agents used in high doses for short periods of time to treat gonorrhea may mask or delay the symptoms of incubating syphilis. All patients with gonorrhea should have a serologic test for syphilis at the time of diagnosis. Patients treated with norfloxacin should have a follow-up serologic test for syphilis after three months.

PRECAUTIONS

General

Needle-shaped crystals were found in the urine of some volunteers who received either placebo, 800 mg norfloxacin, or 1600 mg norfloxacin (at or twice the recommended daily dose, respectively) while participating in a double-blind, crossover study comparing single doses of norfloxacin with placebo. While crystalluria is not expected to occur under usual conditions with a dosage regimen of 400 mg b.i.d., as a precaution, the daily recommended dosage should not be exceeded and the patient should drink sufficient fluids to ensure a proper state of hydration and adequate urinary output.

Alteration in dosage regimen is necessary for patients with impaired renal function (see DOSAGE AND ADMINISTRATION).

Moderate to severe photosensitivity/phototoxicity reactions, the latter of which may manifest as exaggerated sunburn reactions (e.g., burning, erythema, exudation, vesicles, blistering, edema) involving areas exposed to light (typically the face, "V" area of the neck, extensor surfaces of the forearms, dorsa of the hands), can be associated with the use of quinolone antibiotics after sun or UV light exposure. Therefore, excessive exposure to these sources of light should be avoided. Drug therapy should be discontinued if phototoxicity occurs (see ADVERSE REACTIONS, *Post-Marketing*).

Rarely, hemolytic reactions have been reported in patients with latent or actual defects in glucose-6-phosphate dehydrogenase activity who take quinolone antibacterial agents, including norfloxacin (see ADVERSE REACTIONS).

Quinolones, including norfloxacin, may exacerbate the signs of myasthenia gravis and lead to life-threatening weakness of the respiratory muscles. Caution should be exercised when using quinolones, including NOROXIN, in patients with myasthenia gravis (see ADVERSE REACTIONS).

Prescribing NOROXIN in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

Information for Patients

Patients should be advised:

- that norfloxacin may cause changes in the electrocardiogram (QTc interval prolongation).
- that norfloxacin should be avoided in patients receiving class IA (e.g., quinidine, procainamide) or class III (e.g., amiodarone, sotalol) antiarrhythmic agents.
- that norfloxacin should be used with caution in subjects receiving drugs that affect the QTc interval such as cisapride, erythromycin, antipsychotics, and tricyclic antidepressants.
- to inform their physicians of any personal or family history of QTc prolongation or proarrhythmic conditions such as hypokalemia, bradycardia or recent myocardial ischemia.
- that peripheral neuropathies have been associated with norfloxacin use. If symptoms of peripheral neuropathy including pain, burning, tingling, numbness, and/or weakness develop, they should discontinue treatment and contact their physicians.
- to drink fluids liberally.
- that norfloxacin should be taken at least one hour before or at least two hours after a meal or ingestion of milk and/or other dairy products.
- that multivitamins or other products containing iron or zinc, antacids or Videx®[†] (Didanosine), chewable/buffered tablets or the pediatric powder for oral solution, should not be taken within the two-hour period before or within the two-hour period after taking norfloxacin (see PRECAUTIONS, *Drug Interactions*).
- that norfloxacin can cause dizziness and lightheadedness and, therefore, patients should know how they react to norfloxacin before they operate an automobile or machinery or engage in activities requiring mental alertness and coordination.
- to discontinue treatment and inform their physician if they experience pain, inflammation, or rupture of a tendon, and to rest and refrain from exercise. The risk of serious tendon disorders with quinolones is higher in those over 65 years of age and those on corticosteroids.
- that norfloxacin may be associated with hypersensitivity reactions, even following the first dose, and to discontinue the drug at the first sign of a skin rash or other allergic reaction.
- that photosensitivity/phototoxicity has been reported in patients receiving quinolones. Patients should minimize or avoid exposure to natural or artificial sunlight (tanning beds or UVA/B treatment) while taking quinolones. If patients need to be outdoors while using quinolones, they should wear loose-fitting clothes that protect skin from sun exposure and discuss other sun protection measures with their physician. If a sunburn-like reaction or skin eruption occurs, patients should contact their physician.
- that some quinolones may increase the effects of theophylline and/or caffeine (see PRECAUTIONS, *Drug Interactions*).
- that convulsions have been reported in patients taking quinolones, including norfloxacin, and to notify their physician before taking this drug if there is a history of this condition.
- that diarrhea is a common problem caused by antibiotics, which usually ends when the antibiotic is discontinued. Sometimes after starting the treatment with antibiotics, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibiotic. If this occurs, patients should contact their physician as soon as possible.

Patients should be counseled that antibacterial drugs including NOROXIN should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When NOROXIN is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by NOROXIN or other antibacterial drugs in the future.

[†] Registered trademark of Bristol-Myers Squibb Company

Laboratory Tests

As with any potent antibacterial agent, periodic assessment of organ system functions, including renal, hepatic, and hematopoietic, is advisable during prolonged therapy.

Drug Interactions

Quinolones, including norfloxacin, have been shown *in vitro* to inhibit CYP1A2. Concomitant use with drugs metabolized by CYP1A2 (e.g., caffeine, clozapine, ropinirole, tacrine, theophylline, tizanidine) may result in increased substrate drug concentrations when given in usual doses. Patients taking any of these drugs concomitantly with norfloxacin should be carefully monitored.

Elevated plasma levels of theophylline have been reported with concomitant quinolone use. There have been reports of theophylline-related side effects in patients on concomitant therapy with norfloxacin and theophylline. Therefore, monitoring of theophylline plasma levels should be considered and dosage of theophylline adjusted as required.

Elevated serum levels of cyclosporine have been reported with concomitant use of cyclosporine with norfloxacin. Therefore, cyclosporine serum levels should be monitored and appropriate cyclosporine dosage adjustments made when these drugs are used concomitantly.

Quinolones, including norfloxacin, may enhance the effects of oral anticoagulants, including warfarin or its derivatives or similar agents. When these products are administered concomitantly, prothrombin time or other suitable coagulation tests should be closely monitored.

The concomitant administration of quinolones including norfloxacin with glyburide (a sulfonylurea agent) has, on rare occasions, resulted in severe hypoglycemia. Therefore, monitoring of blood glucose is recommended when these agents are co-administered.

Diminished urinary excretion of norfloxacin has been reported during the concomitant administration of probenecid and norfloxacin.

The concomitant use of nitrofurantoin is not recommended since nitrofurantoin may antagonize the antibacterial effect of NOROXIN in the urinary tract.

Multivitamins, or other products containing iron or zinc, antacids or sucralfate, should not be administered concomitantly with, or within 2 hours of, the administration of norfloxacin, because they may interfere with absorption resulting in lower serum and urine levels of norfloxacin.

Videx® (Didanosine) chewable/buffered tablets or the pediatric powder for oral solution should not be administered concomitantly with, or within 2 hours of, the administration of norfloxacin, because these products may interfere with absorption resulting in lower serum and urine levels of norfloxacin.

Some quinolones have also been shown to interfere with the metabolism of caffeine. This may lead to reduced clearance of caffeine and a prolongation of the plasma half-life that may lead to accumulation of caffeine in plasma when products containing caffeine are consumed while taking norfloxacin.

The concomitant administration of a non-steroidal anti-inflammatory drug (NSAID) with a quinolone, including norfloxacin, may increase the risk of CNS stimulation and convulsive seizures. Therefore, NOROXIN should be used with caution in individuals receiving NSAIDs concomitantly.

Carcinogenesis, Mutagenesis, Impairment of Fertility

No increase in neoplastic changes was observed with norfloxacin as compared to controls in a study in rats, lasting up to 96 weeks at doses 8-9 times*** the usual human dose (on a mg/kg basis).

Norfloxacin was tested for mutagenic activity in a number of *in vivo* and *in vitro* tests. Norfloxacin had no mutagenic effect in the dominant lethal test in mice and did not cause chromosomal aberrations in hamsters or rats at doses 30-60 times*** the usual human dose (on a mg/kg basis). Norfloxacin had no mutagenic activity *in vitro* in the Ames microbial mutagen test, Chinese hamster fibroblasts and V-79 mammalian cell assay. Although norfloxacin was weakly positive in the Rec-assay for DNA repair, all other mutagenic assays were negative including a more sensitive test (V-79).

Norfloxacin did not adversely affect the fertility of male and female mice at oral doses up to 30 times*** the usual human dose (on a mg/kg basis).

Pregnancy

Teratogenic Effects. Pregnancy Category C. Norfloxacin has been shown to produce embryonic loss in monkeys when given in doses 10 times*** the maximum daily total human dose (on a mg/kg basis). At this dose, peak plasma levels obtained in monkeys were approximately 2 times those obtained in humans. There has been no evidence of a teratogenic effect in any of the animal species tested (rat, rabbit, mouse, monkey) at 6-50 times*** the maximum daily human dose (on a mg/kg basis). There are, however, no adequate and well-controlled studies in pregnant women. Norfloxacin should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers

It is not known whether norfloxacin is excreted in human milk.

When a 200-mg dose of NOROXIN was administered to nursing mothers, norfloxacin was not detected in human milk. However, because the dose studied was low, because other drugs in this class are secreted in human milk, and because of the potential for serious adverse reactions from norfloxacin in nursing infants, a decision should be made to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

The safety and effectiveness of oral norfloxacin in pediatric patients and adolescents below the age of 18 years have not been established. Norfloxacin causes arthropathy in juvenile animals of several animal species. (See WARNINGS and ANIMAL PHARMACOLOGY.)

Geriatric Use

Of the 340 subjects in one large clinical study of NOROXIN for treatment of urinary tract infections, 103 patients were 65 and older, 77 of whom were 70 and older; no overall differences in safety and effectiveness were evident between these subjects and younger subjects. In clinical practice, no difference in the type of reported adverse experiences have been observed between the elderly and younger patients except for a possible increased risk of tendon rupture in elderly patients receiving concomitant corticosteroids (see WARNINGS). In addition, increased risk for other adverse experiences in some older individuals cannot be ruled out (see ADVERSE REACTIONS).

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function (see DOSAGE AND ADMINISTRATION).

A pharmacokinetic study of NOROXIN in elderly volunteers (65 to 75 years of age with normal renal function for their age) was carried out (see CLINICAL PHARMACOLOGY).

In general, elderly patients may be more susceptible to drug-associated effects of the QTc interval. Therefore, precaution should be taken when using NOROXIN concomitantly with drugs that can result in prolongation of the QTc interval (e.g., class IA or class III antiarrhythmics) or in patients with risk factors for torsades de pointes (e.g., known QTc prolongation, uncorrected hypokalemia).

Patients over 65 years of age are at increased risk for developing severe tendon disorders including tendon rupture when being treated with a fluoroquinolone antibiotic such as NOROXIN. This risk is further increased in patients receiving concomitant corticosteroid therapy. Tendon rupture usually involves the Achilles, hand, or shoulder tendons and can occur during therapy or up to a few months post completion of therapy. Caution should be used when prescribing NOROXIN to elderly patients, especially those on corticosteroids. Patients should be informed of this potential side effect and advised to discontinue therapy and inform their physicians if any tendon symptoms occur.

ADVERSE REACTIONS

Single-Dose Studies

In clinical trials involving 82 healthy subjects and 228 patients with gonorrhea, treated with a single dose of norfloxacin, 6.5% reported drug-related adverse experiences. However, the following incidence figures were calculated without reference to drug relationship.

The most common adverse experiences (>1.0%) were: dizziness (2.6%), nausea (2.6%), headache (2.0%), and abdominal cramping (1.6%).

Additional reactions (0.3%-1.0%) were: anorexia, diarrhea, hyperhidrosis, asthenia, anal/rectal pain, constipation, dyspepsia, flatulence, tingling of the fingers, and vomiting.

Laboratory adverse changes considered drug-related were reported in 4.5% of patients/subjects. These laboratory changes were: increased AST (SGOT) (1.6%), decreased WBC (1.3%), decreased platelet count (1.0%), increased urine protein (1.0%), decreased hematocrit and hemoglobin (0.6%), and increased eosinophils (0.6%).

Multiple-Dose Studies

In clinical trials involving 52 healthy subjects and 1980 patients with urinary tract infections or prostatitis treated with multiple doses of norfloxacin, 3.6% reported drug-related adverse experiences. However, the incidence figures below were calculated without reference to drug relationship.

The most common adverse experiences (>1.0%) were: nausea (4.2%), headache (2.8%), dizziness (1.7%), and asthenia (1.3%).

Additional reactions (0.3%-1.0%) were: abdominal pain, back pain, constipation, diarrhea, dry mouth, dyspepsia/heartburn, fever, flatulence, hyperhidrosis, loose stools, pruritus, rash, somnolence, and vomiting.

Less frequent reactions (0.1%-0.2%) included: abdominal swelling, allergies, anorexia, anxiety, bitter taste, blurred vision, bursitis, chest pain, chills, depression, dysmenorrhea, edema, erythema, foot or hand swelling, insomnia, mouth ulcer, myocardial infarction, palpitation, pruritus ani, renal colic, sleep disturbances, and urticaria.

Abnormal laboratory values observed in these patients/subjects were: eosinophilia (1.5%), elevation of ALT (SGPT) (1.4%), decreased WBC and/or neutrophil count (1.4%), elevation of AST (SGOT) (1.4%), and increased alkaline phosphatase (1.1%). Those occurring less frequently included increased BUN, increased LDH, increased serum creatinine, decreased hematocrit, and glycosuria.

Post-Marketing

The most frequently reported adverse reaction in post-marketing experience is rash.

CNS effects characterized as generalized seizures, myoclonus and tremors have been reported with NOROXIN (see WARNINGS). Visual disturbances have been reported with drugs in this class.

The following additional adverse reactions have been reported since the drug was marketed:

Hypersensitivity Reactions

Hypersensitivity reactions have been reported including anaphylactoid reactions, angioedema, dyspnea, vasculitis, urticaria, arthritis, arthralgia and myalgia (see WARNINGS).

Skin

Toxic epidermal necrolysis, Stevens-Johnson syndrome and erythema multiforme, exfoliative dermatitis, photosensitivity/phototoxicity reactions (see PRECAUTIONS).

Gastrointestinal

Pseudomembranous colitis, hepatitis, jaundice including cholestatic jaundice and elevated liver function tests, pancreatitis (rare), stomatitis. The onset of pseudomembranous colitis symptoms may occur during or after antibacterial treatment (see WARNINGS).

Hepatic

Hepatic failure, including fatal cases.

Cardiovascular

On rare occasions, prolonged QTc interval and ventricular arrhythmia including torsades de pointes.

Renal

Interstitial nephritis, renal failure.

Nervous System/Psychiatric

Peripheral neuropathy, Guillain-Barré syndrome, ataxia, paresthesia, hypoesthesia, psychic disturbances including psychotic reactions and confusion.

Musculoskeletal

Tendinitis, tendon rupture; exacerbation of myasthenia gravis (see PRECAUTIONS); elevated creatine kinase (CK).

Hematologic

Neutropenia; leukopenia; agranulocytosis; hemolytic anemia, sometimes associated with glucose-6-phosphate dehydrogenase deficiency; thrombocytopenia.

Special Senses

Hearing loss, tinnitus, diplopia, dysgeusia.

Other adverse events reported with quinolones include: agranulocytosis, albuminuria, candiduria, crystalluria, cylindruria, dysphagia, elevation of blood glucose, elevation of serum cholesterol, elevation of serum potassium, elevation of serum triglycerides, hematuria, hepatic necrosis, symptomatic hypoglycemia, nystagmus, postural hypotension, prolongation of prothrombin time, and vaginal candidiasis.

OVERDOSAGE

No significant lethality was observed in male and female mice and rats at single oral doses up to 4 g/kg.

In the event of acute overdosage, the stomach should be emptied by inducing vomiting or by gastric lavage, and the patient carefully observed and given symptomatic and supportive treatment. Adequate hydration must be maintained.

DOSAGE AND ADMINISTRATION

Tablets NOROXIN should be taken at least one hour before or at least two hours after a meal or ingestion of milk and/or other dairy products. Multivitamins, other products containing iron or zinc, antacids containing magnesium and aluminum, sucralfate, or Videx® (Didanosine), chewable/buffered tablets or the pediatric powder for oral solution, should not be taken within 2 hours of administration of norfloxacin. Tablets NOROXIN should be taken with a glass of water. Patients receiving NOROXIN should be well hydrated (see PRECAUTIONS).

Normal Renal Function

The recommended daily dose of NOROXIN is as described in the following chart:

| <u>Infection</u> | <u>Description</u> | <u>Unit Dose</u> | <u>Frequency</u> | <u>Duration</u> | <u>Daily Dose</u> |
|-------------------------------|--|------------------|------------------|-----------------|-------------------|
| Urinary Tract | Uncomplicated UTI's (cystitis) due to <i>E. coli</i> , <i>K. pneumoniae</i> , or <i>P. mirabilis</i> | 400 mg | q12h | 3 days | 800 mg |
| | Uncomplicated UTI's due to other indicated organisms | 400 mg | q12h | 7-10 days | 800 mg |
| | Complicated UTI's | 400 mg | q12h | 10-21 days | 800 mg |
| Sexually Transmitted Diseases | Uncomplicated Gonorrhea | 800 mg | single dose | 1 day | 800 mg |
| Prostatitis | Acute or Chronic | 400 mg | q12h | 28 days | 800 mg |

Renal Impairment

NOROXIN may be used for the treatment of urinary tract infections in patients with renal insufficiency. In patients with a creatinine clearance rate of 30 mL/min/1.73 m² or less, the recommended dosage is one 400-mg tablet once daily for the duration given above. At this dosage, the urinary concentration exceeds the MICs for most urinary pathogens susceptible to norfloxacin, even when the creatinine clearance is less than 10 mL/min/1.73 m².

When only the serum creatinine level is available, the following formula (based on sex, weight, and age of the patient) may be used to convert this value into creatinine clearance. The serum creatinine should represent a steady state of renal function.

$$\text{Males: } \frac{(\text{weight in kg}) \times (140 - \text{age})}{(72) \times \text{serum creatinine (mg/100 mL)}}$$

$$\text{Females: } (0.85) \times (\text{above value})$$

Elderly

Elderly patients being treated for urinary tract infections who have a creatinine clearance of greater than 30 mL/min/1.73 m² should receive the dosages recommended under *Normal Renal Function*.

Elderly patients being treated for urinary tract infections who have a creatinine clearance of 30 mL/min/1.73 m² or less should receive 400 mg once daily as recommended under *Renal Impairment*.

HOW SUPPLIED

No. 8338 — Tablets NOROXIN 400 mg are white to off-white, oval shaped, film-coated tablets, coded 705 on one side and plain on the other. They are supplied as follows:

NDC 0006-0705-68 bottles of 100

NDC 0006-0705-20 unit of use bottles of 20.

Storage

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature]. Keep container tightly closed.

ANIMAL PHARMACOLOGY

Norfloxacin and related drugs have been shown to cause arthropathy in immature animals of most species tested (see WARNINGS).

Crystalluria has occurred in laboratory animals tested with norfloxacin. In dogs, needle-shaped drug crystals were seen in the urine at doses of 50 mg/kg/day. In rats, crystals were reported following doses of 200 mg/kg/day.

Embryo lethality and slight maternotoxicity (vomiting and anorexia) were observed in cynomolgus monkeys at doses of 150 mg/kg/day or higher.

Ocular toxicity, seen with some related drugs, was not observed in any norfloxacin-treated animals.

REFERENCES

1. National Committee for Clinical Laboratory Standards, Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically - 3rd ed., Approved Standard NCCLS Document M7-A3, Vol. 13, No. 25, NCCLS, Villanova, PA, 1993.
 2. National Committee for Clinical Laboratory Standards, Performance standards for antimicrobial disk susceptibility tests - 5th ed., Approved Standard NCCLS Document M2-A5, Vol. 13, No. 24, NCCLS, Villanova, PA, 1993.
-

Manufactured for:

 **MERCK & CO., INC.**, Whitehouse Station, NJ 08889, USA

By: Merck Sharp & Dohme (Italia) S.p.A.
Via Emilia, 21
27100 Pavia, Italy

Issued April 2008
Printed in USA

Patient Information
NOROXIN[®]
(norfloxacin)

400 mg Tablets

This section contains important information about NOROXIN* (norfloxacin), and should be read completely before you begin treatment. This section does not take the place of discussions with your doctor or health care professional about your medical condition or your treatment. This section does not list all benefits and risks of NOROXIN. The medicine described here can be prescribed only by a licensed health care professional. If you have any questions about NOROXIN, talk with your health care professional. Only your health care professional can determine if NOROXIN is right for you.

What is NOROXIN?

NOROXIN is an antimicrobial used to treat urinary tract infections, kidney infections, prostatitis and some forms of sexually transmitted diseases (specifically gonorrhea), caused by certain germs called bacteria. NOROXIN kills many of the types of bacteria that can infect the bladder, kidneys, prostate gland and urethra and has been shown in clinical trials to be safe and effective for the treatment of these bacterial infections. NOROXIN is not effective in the treatment of syphilis.

You should contact your doctor if you think your condition is not improving while taking NOROXIN.

NOROXIN Tablets contain 400 mg of active drug.

How and when should I take NOROXIN?

NOROXIN is usually taken twice a day for 3 to 28 days depending on the condition for which you are being treated. For gonorrhea it is given as a single dose of 2 pills taken at one time. It should be swallowed whole and taken with a large glass of water at least 1 hour before or 2 hours after a meal. Try to take the tablet at the same time each day.

You may begin to feel better quickly; however, in order to make sure that all bacteria are killed, you should complete the full course of medication. Do not take more than the prescribed dose of NOROXIN even if you missed a dose by mistake. You should not take a double dose.

Who should not take NOROXIN?

You should not take NOROXIN if you have ever had a severe allergic reaction to any of the group of antimicrobials known as "quinolones" such as ciprofloxacin or levofloxacin. If you develop hives, difficulty breathing, or other symptoms of a severe allergic reaction, seek emergency treatment right away. If you develop a skin rash, you should stop taking NOROXIN and call your health care professional.

You should avoid NOROXIN if you have a rare condition known as congenital prolongation of the QTc interval. If you or any of your family members have this condition you should inform your health care professional. You should avoid NOROXIN if you are being treated for heart rhythm disturbances with certain medicines such as quinidine, procainamide, amiodarone or sotalol. Inform your health care professional if you are taking a heart rhythm drug or if you have any problems with your heart rhythm.

You should also avoid NOROXIN if the amount of potassium in your blood is low. Low potassium can

* Registered trademark of MERCK & CO., Inc.
COPYRIGHT © 1986, 1989, 1999, 2001 MERCK & CO., Inc.
All rights reserved

sometimes be caused by medicines called diuretics such as furosemide and hydrochlorothiazide. If you are taking a diuretic medicine you should speak with your health care professional.

If you are pregnant or planning to become pregnant while taking NOROXIN, talk to your doctor before taking this medication. NOROXIN is not recommended for use during pregnancy or nursing, as the effects on the unborn child or nursing infant are unknown.

NOROXIN is not recommended for children.

What are the possible side effects of NOROXIN?

NOROXIN is generally well tolerated. The most common side effects caused by NOROXIN, which are usually mild, include dizziness, nausea, heartburn, headache, and rash. If a rash occurs stop taking NOROXIN and call your health care provider. Less common side effects may include changes in hearing and pain, burning, tingling, numbness, and/or weakness in arms or legs.

If you should develop hives, trouble breathing, wheezing, or other signs of an allergic reaction seek help immediately.

You should be careful about driving or operating machinery until you are sure NOROXIN is not causing dizziness.

Pain, swelling, and tears of the Achilles, shoulder, or hand tendons have been reported in patients receiving fluoroquinolones, including NOROXIN. The risk for tendon effects is higher if you are over 65 years of age, or if you are taking corticosteroids. If you develop pain, swelling, or tear of a tendon you should stop taking NOROXIN, avoid exercise and strenuous use of the affected area, and contact your health care provider.

Convulsions have been reported in patients receiving quinolone antibiotics. Be sure to let your physician know if you have a history of convulsions. Quinolones, including NOROXIN, have been rarely associated with other central nervous system events including confusion, tremors, hallucinations, anxiety, and depression.

Quinolones, including NOROXIN, may worsen the signs of myasthenia gravis leading to difficulty breathing which may be life-threatening. If you develop this symptom contact your health care provider right away.

NOROXIN may cause a rare heart problem known as prolongation of the QTc interval. This condition can cause an abnormal heartbeat and can be very dangerous. The chances of this event are increased in those with a family history of prolonged QTc interval, low potassium (hypokalemia), and those who are taking drugs to control heart rhythm called class IA (quinidine, procainamide), or class III (amiodarone, sotalol) antiarrhythmic agents. You should call your healthcare provider right away if you have symptoms of prolongation of the QTc interval including prolonged heart palpitation (a change in the way your heart beats) or a loss of consciousness (fainting spells).

Diarrhea that usually ends after treatment is a common problem caused by antimicrobials. A more serious form of diarrhea can occur during or up to 2 months after the use of antibiotics. This has been reported with all antimicrobials, including NOROXIN. If you develop a watery or bloody stool, with or without stomach cramps and fever, contact your physician as soon as possible.

Sun sensitivity (photosensitivity), which can appear as skin eruption or severe sunburn, can occur in some patients taking quinolone antimicrobials after exposure to sunlight or artificial ultraviolet light (UV) (e.g., tanning beds). NOROXIN has been infrequently associated with photosensitivity. Avoid excessive exposure to sunlight or artificial UV light while taking NOROXIN. Use a sunscreen and wear protective clothing if you are exposed to the sun. If photosensitivity develops, contact your physician.

NOROXIN can cause crystals to form in the urine of patients taking it if they are dehydrated. Make sure you drink plenty of water while taking NOROXIN.

If you notice any side effects not mentioned in this section or you have any concerns about the side effects you are experiencing, please inform your health care professional.

What about other medicines I am taking?

Tell your doctor about all other prescription and non-prescription medicines, dietary supplements and herbals you are taking. You should avoid taking NOROXIN with certain medicines used to treat an abnormal heartbeat. These include quinidine, procainamide, amiodarone, and sotalol.

Some medicines also produce an effect on the electrocardiogram test, including cisapride, erythromycin, some antidepressants, and some antipsychotic drugs. These may increase the risk of heart beat problems when taken with NOROXIN.

Many antacids and multivitamins may interfere with the absorption of NOROXIN and may prevent it from working properly. You should take NOROXIN either 2 hours before or 2 hours after taking these products.

Remember

- Take your dose of NOROXIN every day.
- Complete the course of medication even if you are feeling better.
- Keep this medication out of the reach of children.
- This information does not take the place of discussions with your doctor or health care professional about your medical condition or your treatment.
- For more complete information about NOROXIN request full prescribing information from your health care professional, pharmacist, or visit our website at www.merck.com.

Manufactured for:

 MERCK & CO., INC., Whitehouse Station, NJ 08889, USA

Manufactured by:

Merck Sharp & Dohme (Italia) S.p.A.
Via Emilia, 21
27100 – Pavia, Italy

Issued April 2008

9868300