

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all of the information needed to use Durezol safely and effectively. See full prescribing information for Durezol.

Durezol (difluprednate ophthalmic emulsion) 0.05%

Initial U.S. approval: 2008

-----INDICATIONS AND USAGE-----

Durezol is a topical corticosteroid that is indicated for the treatment of inflammation and pain associated with ocular surgery. (1)

-----DOSAGE AND ADMINISTRATION-----

Instill one drop into the conjunctival sac of the affected eye(s) 4 times daily beginning 24 hours after surgery and continuing throughout the first 2 weeks of the postoperative period, followed by 2 times daily for a week and then a taper based on the response. (2)

-----DOSAGE FORMS AND STRENGTHS-----

Durezol contains 0.05% difluprednate, as a sterile preserved ophthalmic emulsion for topical ophthalmic use only. (3)

-----CONTRAINDICATIONS-----

Durezol, as with other ophthalmic corticosteroids, is contraindicated in most viral diseases of the cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella, and also in mycobacterial infection of the eye and fungal diseases of ocular structures. (4)

-----WARNINGS AND PRECAUTIONS-----

- Intraocular pressure (IOP) increase-Prolonged use of corticosteroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision. If this product is used for 10 days or longer, IOP should be monitored. (5.1).

- Cataracts- Use of corticosteroids may result in posterior subcapsular cataract formation. (5.2)
- Delayed healing- The use of steroids after cataract surgery may delay healing and increase the incidence of bleb formation. In those diseases causing thinning of the cornea or sclera, perforations have been known to occur with the use of topical steroids. The initial prescription and renewal of the medication order beyond 28 days should be made by a physician only after examination of the patient with the aid of magnification such as slit lamp biomicroscopy and, where appropriate, fluorescein staining. (5.3)
- Bacterial infections- Prolonged use of corticosteroids may suppress the host response and thus increase the hazard of secondary ocular infections. In acute purulent conditions, steroids may mask infection or enhance existing infection. If signs and symptoms fail to improve after 2 days, the patient should be re-evaluated. (5.4)
- Viral infections- Employment of a corticosteroid medication in the treatment of patients with a history of herpes simplex requires great caution. Use of ocular steroids may prolong the course and may exacerbate the severity of many viral infections of the eye (including herpes simplex). (5.5)
- Fungal infections- Fungal infections of the cornea are particularly prone to develop coincidentally with long-term local steroid application. Fungus invasion must be considered in any persistent corneal ulceration where a steroid has been used or is in use. (5.6)

To report SUSPECTED ADVERSE REACTIONS, contact Sirion Therapeutics at (TBD) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION.

Revised date: June 2008

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE
- 2 DOSAGE AND ADMINISTRATION
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
 - 5.1 IOP increase
 - 5.2 Cataracts
 - 5.3 Delayed Healing
 - 5.4 Bacterial infections
 - 5.5 Viral infections
 - 5.6 Fungal infections
 - 5.7 Topical ophthalmic use only
- 6 ADVERSE REACTIONS
- 8 USE IN SPECIFIC POPULATIONS
 - 8.1 Pregnancy
 - 8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

- 11 DESCRIPTION
- 12 CLINICAL PHARMACOLOGY
 - 12.1 Mechanism of Action
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
 - 13.2 Animal Toxicology and/or Pharmacology
- 14 CLINICAL STUDIES
 - 14.1 Postoperative Ocular Inflammation and Pain
- 16 HOW SUPPLIED/STORAGE AND HANDLING
- 17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 Indications and Usage

Durezol (difluprednate ophthalmic emulsion) 0.05%, a topical corticosteroid, is indicated for the treatment of inflammation and pain associated with ocular surgery.

2 Dosage and Administration

Instill one drop into the conjunctival sac of the affected eye(s) 4 times daily beginning 24 hours after surgery and continuing throughout the first 2 weeks of the postoperative period, followed by 2 times daily for a week and then a taper based on the response. (2)

3 Dosage Forms and Strengths

Durezol contains 0.05% difluprednate as a sterile preserved emulsion for topical ophthalmic administration.

4 Contraindications

The use of Durezol, as with other ophthalmic corticosteroids, is contraindicated in most active viral diseases of the cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella, and also in mycobacterial infection of the eye and fungal disease of ocular structures.

5 Warnings and Precautions

5.1 IOP Increase

Prolonged use of corticosteroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision. Steroids should be used with caution in the presence of glaucoma. If this product is used for 10 days or longer, intraocular pressure should be monitored.

5.2 Cataracts

Use of corticosteroids may result in posterior subcapsular cataract formation.

5.3 Delayed healing

The use of steroids after cataract surgery may delay healing and increase the incidence of bleb formation. In those diseases causing thinning of the cornea or sclera, perforations have been known to occur with the use of topical steroids. The initial prescription and renewal of the medication order beyond 28 days should be made by a physician only after examination of the patient with the aid of magnification such as slit lamp biomicroscopy and, where appropriate, fluorescein staining.

5.4 Bacterial infections

Prolonged use of corticosteroids may suppress the host response and thus increase the hazard of secondary ocular infections. In acute purulent conditions, steroids may mask infection or enhance existing infection. If signs and symptoms fail to improve after 2 days, the patient should be re-evaluated.

5.5 Viral infections

Employment of a corticosteroid medication in the treatment of patients with a history of herpes simplex requires great caution. Use of ocular steroids may prolong the course and may exacerbate the severity of many viral infections of the eye (including herpes simplex).

5.6 Fungal infections

Fungal infections of the cornea are particularly prone to develop coincidentally with long-term local steroid application. Fungus invasion must be considered in any persistent corneal ulceration where a steroid has been used or is in use. Fungal culture should be taken when appropriate.

5.7 Topical ophthalmic use only

Durezol is not indicated for intraocular administration.

6 Adverse Reactions

Adverse reactions associated with ophthalmic steroids include elevated intraocular pressure, which may be associated with optic nerve damage, visual acuity and field defects, posterior subcapsular cataract formation, secondary ocular infection from pathogens including herpes simplex, and perforation of the globe where there is thinning of the cornea or sclera.

Ocular adverse reactions occurring in 5–15% of subjects in clinical studies with Durezol included corneal edema, ciliary and conjunctival hyperemia, eye pain, photophobia, posterior capsule opacification, anterior chamber cells, anterior chamber flare, conjunctival edema, and blepharitis. Other ocular adverse reactions occurring in 1–5% of subjects included reduced visual acuity, punctate keratitis, eye inflammation, and iritis. Ocular adverse events occurring in < 1% of subjects included application site discomfort or irritation, corneal

pigmentation and striae, episcleritis, eye pruritis, eyelid irritation and crusting, foreign body sensation, increased lacrimation, macular edema, scleral hyperemia, and uveitis. Most of these events may have been the consequence of the surgical procedure.

8 Use in Specific Populations

8.1 Pregnancy

Teratogenic Effects

Pregnancy Category C. Difluprednate has been shown to be embryotoxic (decrease in embryonic body weight and a delay in embryonic ossification) and teratogenic (cleft palate and skeletal anomalies when administered subcutaneously to rabbits during organogenesis at a dose of 1–10 µg/kg/day. The no-observed-effect-level (NOEL) for these effects was 1 µg/kg/day, and 10 µg/kg/day was considered to be a teratogenic dose that was concurrently found in the toxic dose range for fetuses and pregnant females. Treatment of rats with 10 µg/kg/day subcutaneously during organogenesis did not result in any reproductive toxicity, nor was it maternally toxic. At 100 µg/kg/day after subcutaneous administration in rats, there was a decrease in fetal weights and delay in ossification, and effects on weight gain in the pregnant females. It is difficult to extrapolate these doses of difluprednate to maximum daily human doses of Durezol, since Durezol is administered topically with minimal systemic absorption, and difluprednate blood levels were not measured in the reproductive animal studies. However, since use of difluprednate during human pregnancy has not been evaluated and cannot rule out the possibility of harm, Durezol should be used during pregnancy only if the potential benefit justifies the potential risk to the embryo or fetus.

8.3 Nursing Mothers

It is not known whether topical ophthalmic administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. Caution should be exercised when Durezol is administered to a nursing woman.

8.4 Pediatric Use


Safety and effectiveness in pediatric patients has not been established.

8.5 Geriatric Use

No overall differences in safety or effectiveness have been observed between elderly and younger patients.

11 Description

Durezol (difluprednate ophthalmic emulsion) 0.05% is a sterile, topical anti-inflammatory corticosteroid for ophthalmic use. The chemical name is 6α,9-difluoro-11β,17,21-trihydroxypregna-1,4-diene-3,20-dione 21-acetate 17-butyrate (CAS number 23674-86-4). Difluprednate is represented by the following structural formula:



Difluprednate has a molecular weight of 508.56, and the empirical formula is C₂₇H₃₄F₂O₇.

Each mL contains: ACTIVE: difluprednate 0.5 mg (0.05%); INACTIVES: boric acid, castor oil, glycerin, polysorbate 80, purified water, sodium acetate, sodium EDTA, sodium hydroxide (to adjust the pH to 5.2 to 5.8). The emulsion is essentially isotonic with a tonicity of 304 to 411 mOsm/kg. PRESERVATIVE: sorbic acid 0.1%.

12 Clinical Pharmacology

12.1 Mechanism of Action

Corticosteroids inhibit the inflammatory response to a variety of inciting agents that may delay or slow healing. They inhibit edema, fibrin deposition, capillary dilation, leukocyte migration, capillary proliferation, fibroblast proliferation, deposition of collagen, and scar formation associated with inflammation. There is no generally accepted explanation for the mechanism of action of ocular corticosteroids. However, corticosteroids are thought to act by the induction of phospholipase A2 inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of

inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A2.

Difluprednate is structurally similar to other corticosteroids.

12.3 Pharmacokinetics

Difluprednate undergoes deacetylation *in vivo* to 6 α ,9-difluoroprednisolone 17-butyrate (DFB), an active metabolite of difluprednate.

Clinical pharmacokinetic studies of difluprednate after repeat ocular instillation of 2 drops of difluprednate (0.01% or 0.05%) QID for 7 days showed that DFB levels in blood were below the quantification limit (50 ng/mL) at all time points for all subjects, indicating the systemic absorption of difluprednate after ocular instillation of Durezol is limited.

13 Nonclinical Toxicology

13.1 Carcinogenesis, Mutagenesis, and Impairment of Fertility

Difluprednate was not genotoxic *in vitro* in the Ames test, and in cultured mammalian cells CHL/IU (a fibroblastic cell line derived from the lungs of newborn female Chinese hamsters). An *in vivo* micronucleus test of difluprednate in mice was also negative. Treatment of male and female rats with subcutaneous difluprednate up to 10 μ g/kg/day prior to and during mating did not impair fertility in either gender. Long term studies have not been conducted to evaluate the carcinogenic potential of difluprednate.

13.2 Animal Toxicology and/or Pharmacology

In multiple studies performed in rodents and non-rodents, subchronic and chronic toxicity tests of difluprednate showed systemic effects such as suppression of body weight gain; a decrease in lymphocyte count; atrophy of the lymphatic glands and adrenal gland; and for local effects, thinning of the skin; all of which were due to the pharmacologic action of the molecule and are well known glucocorticosteroid effects. Most, if not all of these effects were reversible after drug withdrawal. The NOEL for the subchronic and chronic toxicity tests were consistent between species and ranged from 1–1.25 μ g/kg per day.

14 Clinical Studies

14.1 Postoperative Ocular Inflammation and Pain

Clinical efficacy was evaluated in 2 randomized, double-masked, placebo-controlled trials in which subjects with an anterior chamber cell grade \geq "2" (a cell count of 10 or higher) after cataract surgery were assigned to Durezol or placebo (vehicle) following surgery. One drop of Durezol or vehicle was self instilled either 2 (BID) or 4 (QID) times per day for 14 days, beginning the day after surgery. The presence of complete clearing (a cell count of 0) was assessed 8 and 15 days post-surgery using a slit lamp binocular microscope. In the intent-to-treat analyses of both studies, a significant benefit was seen in the QID Durezol-treated group in ocular inflammation and reduction of pain when compared with placebo. The consolidated clinical trial results are provided below.

| Ocular Inflammation and Pain Endpoints (Studies Pooled) | | | | |
|---|------------------------|--------------|--------------------|-------------|
| | Durezol QID N = 107 | | Vehicle N = 220 | |
| Day | 8 | 15 | 8 | 15 |
| Anterior Chamber cell clearing (% subjects) | 24 (22%)* | 44 (41%)* | 17 (7%) | 25 (11%) |
| Pain free (% subjects) | 62 (58%)* | 67 (63%)* | 59 (27%) | 76 (35%) |

* Statistically significantly better than vehicle, p<0.01

16 How Supplied/Storage and Handling

Durezol (difluprednate ophthalmic emulsion) 0.05% is a sterile, aqueous topical ophthalmic emulsion supplied in an opaque plastic bottle with a controlled drop tip and a pink cap in the following sizes:

- 2.5 mL in a 5 mL bottle (NDC 42826-601-25)
- 5 mL in a 5 mL bottle (NDC 42826-601-05)

Storage

Store at 15-25°C (59-77°F). Do not freeze. Protect from light. When not in use keep the bottles in the protective carton and the unused vials in the protective foil pouch.

Revised: June 2008

17 Patient Counseling Information

This product is sterile when packaged. Patients should be advised not to allow the dropper tip to touch any surface, as this may contaminate the emulsion. If pain develops or if redness, itching, or inflammation becomes aggravated, the patient should be advised to consult a physician. As with all ophthalmic preparations containing a preservative, patients should be advised not to wear contact lenses when using Durezol.