

 **SALONPAS** *PAIN RELIEF PATCH*  
Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

**15** PATCHES

**SALONPAS** *PAIN RELIEF PATCH* \*  
Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

*Effectiveness confirmed in clinical trial*  
NDC#46581-670-15

**ULTRA THIN  
COMFORT  
STRETCH**


- STIFF NECK
- SORE SHOULDER
- BACKACHE
- JOINT PAIN

**FDA Approved Non Prescription  
Pain Relieving Patch**

✳️ **DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS**

Manufactured by  
**Hisamitsu Pharmaceutical Co., Inc.** JAPAN SAGA TOSU  
Questions or comments? Toll free 1-877-SALONPAS MON-FRI 9AM to 5PM (PST)  
[www.salonpas.us](http://www.salonpas.us) **15**  $2\frac{3}{4} \times 3\frac{15}{16}$  inch (7cm x 10cm) PATCHES  
Made in Japan

**Hisamitsu**®



 **SALONPAS** *PAIN RELIEF PATCH*  
Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

**15** PATCHES

LOT

EXP.

4M219T-0

# SALONPAS<sup>®</sup>

**PAIN RELIEF PATCH**  
PAIN RELIEVING PATCH

Press  
Open

minty scent

Package not child resistant.  
Keep out of reach of children.

## Drug Facts

### Active ingredients (in each patch)

Menthol 3%.....	Topical analgesic
Methyl salicylate 10%(NSAID*).....	Topical analgesic

\*nonsteroidal anti-inflammatory drug

### Purpose

### Uses

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:

- strains
- sprains
- simple backache
- arthritis
- bruises

### Warnings

#### For external use only

**Stomach bleeding warning:** This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### Do not use

- on the face or rashes
- on wounds or damaged skin
- if allergic to aspirin or other NSAIDs
- with a heating pad
- when sweating (such as from exercise or heat)
- any patch from a pouch that has been open for 14 or more days
- right before or after heart surgery

### Ask a doctor before use if

- you are allergic to topical products
- the stomach bleeding warning applies to you
- you have high blood pressure, heart disease, or kidney disease
- you are taking a diuretic

### When using this product

- wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

### Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- rash, itching or skin irritation develops
- condition worsens
- symptoms last for more than 3 days
- symptoms clear up and occur again within a few days
- stomach pain or upset gets worse or lasts

**If pregnant or breast-feeding,** ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant.

### Directions

#### Adults 18 years and older:

- clean and dry affected area
- remove patch from backing film and apply to skin (see illustration)
- apply one patch to the affected area and leave in place for up to 8 to 12 hours
- if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
- only use one patch at a time
- do not use more than 2 patches per day
- do not use for more than 3 days in a row

#### Children under 18 years of age:

- do not use

### Drug Facts (Continued)

#### Other information

- some individuals may not experience pain relief until several hours after applying the patch
- avoid storing product in direct sunlight
- protect product from excessive moisture
- store at 20-25°C (68-77°F)

### Inactive ingredients

atlocyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

 **SALONPAS** PAIN RELIEF PATCH **5** PATCHES  
Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

# SALONPAS<sup>®</sup>

## PAIN RELIEF PATCH

Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

*Effectiveness confirmed in clinical trial*

NDC#46581-670-05



**ULTRA THIN  
COMFORT  
STRETCH**

- STIFF NECK
- SORE SHOULDER
- BACKACHE
- JOINT PAIN

**FDA Approved Non Prescription  
Pain Relieving Patch**

⚠ **DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS**

Manufactured by  
**Hisamitsu Pharmaceutical Co., Inc.** JAPAN SAGA TOSU  
Questions or comments? Toll free 1-87-SALONPAS MON-FRI 9AM to 5PM (PST)  
[www.salonpas.us](http://www.salonpas.us) Made in Japan

**Hisamitsu** <sup>®</sup>

**5** 2 3/4 x 3 15/16 inch  
(7cm x 10cm)  
**PATCHES**

4M218T-0



LOT

EXP.

 **SALONPAS** PAIN RELIEF PATCH  
Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH  
**5** PATCHES

(b) (4)

**SALONPAS**  
**PAIN RELIEF PATCH**  
PAIN RELIEVING PATCH

Press  
Open

minty scent

Package not child resistant.  
Keep out of reach of children.

**Drug Facts**

**Active ingredients (in each patch)**

**Purpose**

Menthol 3%..... Topical analgesic  
Methyl salicylate 10%(NSAID\*)..... Topical analgesic  
\*nonsteroidal anti-inflammatory drug

**Uses**

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:  
■ strains ■ sprains ■ simple backache ■ arthritis ■ bruises

**Warnings**

**For external use only**

**Stomach bleeding warning:** This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:

- are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use**

- on the face or rashes ■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs ■ with a heating pad ■ when sweating (such as from exercise or heat) ■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

**Ask a doctor before use if**

- you are allergic to topical products ■ the stomach bleeding warning applies to you ■ you have high blood pressure, heart disease, or kidney disease ■ you are taking a diuretic

**When using this product**

- wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water. ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding. ■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days ■ symptoms clear up and occur again within a few days ■ stomach pain or upset gets worse or lasts

**If pregnant or breast-feeding**, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant.

**Directions**

**Adults 18 years and older:**

- clean and dry affected area ■ remove patch from backing film and apply to skin (see illustration) ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours ■ only use one patch at a time ■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row

**Children under 18 years of age:**

- do not use

■ some individuals may not experience pain relief until several hours after applying the patch ■ avoid storing product in direct sunlight ■ protect product from excessive moisture ■ store at 20-25°C (68-77°F)

material: oil, polyisobutylene, polyisobutylene, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

(b) (4)



> CUT OPEN HERE ----- CUT OPEN HERE <

FOLD TWICE AFTER OPENING


Discard unused patches 14 days after opening

FOLD TWICE AFTER OPENING

# SALONPAS® PAIN RELIEF PATCH

Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

*Effectiveness confirmed in clinical trial*



Questions or comments?  
Toll free 1-87-SALONPAS MON-FRI 9AM to 5PM (PST)

[www.salonpas.us](http://www.salonpas.us)

**※DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS**

**Hisamitsu®**

Manufactured by  
**Hisamitsu Pharmaceutical Co., Inc.**  
JAPAN SAGA TOSU Made in Japan

**5** 2 3/4 X 3 1/2 inches  
(7cm X 10cm)  
**PATCHES**

CUT OPEN HERE ----- CUT OPEN HERE

FOLD TWICE AFTER OPENING

Discard unused patches 14 days after opening

FOLD TWICE AFTER OPENING

SALONPAS PAIN RELIEF PATCH PAIN RELIEVING PATCH minty scent

Drug Facts

Active ingredients (in each patch)

Purpose

Menthol 3% Topical analgesic
Methyl salicylate 10%(NSAID)\* Topical analgesic
\*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
strains sprains simple backache arthritis bruises

Warnings

For external use only

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

Do not use

on the face or rashes on wounds or damaged skin if allergic to aspirin or other NSAIDs with a heating pad when sweating (such as from exercise or heat) any patch from a pouch that has been open for 14 or more days right before or after heart surgery

Ask a doctor before use if

you are allergic to topical products the stomach bleeding warning applies to you you have high blood pressure, heart disease, or kidney disease you are taking a diuretic

When using this product

wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water. the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding. rash, itching or skin irritation develops condition worsens symptoms last for more than 3 days symptoms clear up and occur again within a few days stomach pain or upset gets worse or lasts

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant.

Directions

Adults 18 years and older:

clean and dry affected area remove patch from backing film and apply to skin (see illustration) apply one patch to the affected area and leave in place for up to 8 to 12 hours if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours only use one patch at a time do not use more than 2 patches per day do not use for more than 3 days in a row

Children under 18 years of age:

do not use

Other information

some individuals may not experience pain relief until several hours after applying the patch avoid storing product in direct sunlight protect product from excessive moisture store at 20-25°C (68-77°F)

Inactive ingredients

alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Package not child resistant. Keep out of reach of children.

4L296T-0

LOT

EXP.

(b) (4)



✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

FOLD TWICE AFTER OPENING

**Discard unused patches 14 days after opening**

FOLD TWICE AFTER OPENING

# SALONPAS<sup>®</sup>

## PAIN RELIEF PATCH<sup>®</sup>


Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

*Effectiveness confirmed in clinical trial*

NDC#46581-670-03



### ULTRA THIN COMFORT STRETCH



**FREE  
SAMPLE**

**FDA Approved Non Prescription  
Pain Relieving Patch**

- STIFF NECK
- SORE SHOULDER
- BACKACHE
- JOINT PAIN

Questions or comments?  
Toll free 1-87-SALONPAS MON-FRI 9AM to 5PM (PST) [www.salonpas.us](http://www.salonpas.us)

**✂ DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS**

**Hisamitsu**

Manufactured by  
**Hisamitsu Pharmaceutical Co., Inc.**  
JAPAN SAGA TOSU Made in Japan

**3** 2 3/4" x 3 1/8" inch  
(7cm x 10cm)  
**PATCHES**

> CUT OPEN HERE ----- CUT OPEN HERE >

**SALONPAS<sup>®</sup> PAIN RELIEF PATCH** PAIN RELIEVING PATCH

minty scent

### Drug Facts

#### Active ingredients (in each patch)

#### Purpose

Menthol 3% ..... Topical analgesic  
Methyl salicylate 10%(NSAID)\* ..... Topical analgesic  
\*nonsteroidal anti-inflammatory drug

#### Uses

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:  
■ strains ■ sprains ■ simple backache ■ arthritis ■ bruises

#### Warnings

##### For external use only

**Stomach bleeding warning:** This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:

■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ on the face or rashes ■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs ■ with a heating pad ■ when sweating (such as from exercise or heat) ■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

**Ask a doctor before use if** ■ you are allergic to topical products ■ the stomach bleeding warning applies to you ■ you have high blood pressure, heart disease, or kidney disease ■ you are taking a diuretic

**When using this product** ■ wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water. ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if** ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding. ■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days ■ symptoms clear up and occur again within a few days ■ stomach pain or upset gets worse or lasts

**If pregnant or breast-feeding,** ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant.

#### Directions

##### Adults 18 years and older:

■ clean and dry affected area ■ remove patch from backing film and apply to skin (see illustration) ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours ■ only use one patch at a time ■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row

**Children under 18 years of age:** ■ do not use

#### Other information

■ some individuals may not experience pain relief until several hours after applying the patch ■ avoid storing product in direct sunlight ■ protect product from excessive moisture ■ store at 20-25°C (68-77°F)

LOT

#### Inactive ingredients

alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

EXP.

Package not child resistant. Keep out of reach of children.

4L297T-0

 **SALONPAS** *ARTHRITIS PAIN*  
Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH


**15** PATCHES

**SALONPAS** *ARTHRITIS PAIN*  
Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

*Effectiveness confirmed in clinical trial*  
NDC#46581-680-15

**ULTRA THIN  
COMFORT  
STRETCH**

**FDA Approved Non Prescription  
Pain Relieving Patch**



**DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS**

Manufactured by **Hisamitsu Pharmaceutical Co., Inc.** JAPAN SAGA TOSU  
Questions or comments? Toll free 1-877-SALONPAS MON-FRI 9AM to 5PM (PST) **15**  $2\frac{3}{4} \times 3\frac{15}{16}$  inch (7cm x 10cm) **PATCHES**  
**Hisamitsu** [www.salonpas.us](http://www.salonpas.us) Made in Japan

 **SALONPAS** *ARTHRITIS PAIN*  
Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

**15** PATCHES

LOT  
EXP.

3M232T-0

# SALONPAS<sup>®</sup> ARTHRITIS PAIN PAIN RELIEVING PATCH

Press  
Open

minty scent

Package not child resistant.  
Keep out of reach of children.

## Drug Facts

### Active ingredients (in each patch)

	Purpose
Menthol 3%.....	Topical analgesic
Methyl salicylate 10%(NSAID*).....	Topical analgesic

\*nonsteroidal anti-inflammatory drug

### Uses

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:

- arthritis
- sprains
- strains
- bruises
- simple backache

### Warnings

#### For external use only

**Stomach bleeding warning:** This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### Do not use

- on the face or rashes
- on wounds or damaged skin
- if allergic to aspirin or other NSAIDs
- with a heating pad
- when sweating (such as from exercise or heat)
- any patch from a pouch that has been open for 14 or more days
- right before or after heart surgery

#### Ask a doctor before use if

- you are allergic to topical products
- the stomach bleeding warning applies to you
- you have high blood pressure, heart disease, or kidney disease
- you are taking a diuretic

#### When using this product

- wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

#### Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- rash, itching or skin irritation develops
- condition worsens
- symptoms last for more than 3 days
- symptoms clear up and occur again within a few days
- stomach pain or upset gets worse or lasts

**If pregnant or breast-feeding,** ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant.

### Directions

#### Adults 18 years and older:

- clean and dry affected area
- remove patch from backing film and apply to skin (see illustration)
- apply one patch to the affected area and leave in place for up to 8 to 12 hours
- if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
- only use one patch at a time
- do not use more than 2 patches per day
- do not use for more than 3 days in a row

#### Children under 18 years of age:

- do not use

### Drug Facts (Continued)

**Other information**

- some individuals may not experience pain relief until several hours after applying the patch
- avoid storing product in direct sunlight
- protect product from excessive moisture
- store at 20-25°C (68-77°F)

### Inactive ingredients

alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate



**SALONPAS** <sup>ARTHRITIS PAIN</sup>  
Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

**5**  
PATCHES

# SALONPAS<sup>®</sup>

## ARTHRITIS PAIN<sup>※</sup>

Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

*Effectiveness confirmed in clinical trial*

NDC#46581-680-05

**ULTRA THIN**  
**COMFORT**  
**STRETCH**

**FDA Approved Non Prescription**  
**Pain Relieving Patch**

**※ DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS**

**Hisamitsu** <sup>®</sup>

Manufactured by  
**Hisamitsu Pharmaceutical Co., Inc.** JAPAN SAGA TOSU  
Questions or comments? Toll free 1-877-SALONPAS MON-FRI 9AM to 5PM (PST)  
[www.salonpas.us](http://www.salonpas.us)

Made in Japan

**5** <sup>2 3/4" X 3 1/8" (7cm X 10cm)</sup>  
**PATCHES**

3M231T-0



LOT

EXP.



**SALONPAS** <sup>ARTHRITIS PAIN</sup>  
Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

**5** PATCHES

Drug Facts (Continued)

Inactive ingredients

(b) (4)

**SALONPAS**  
**ARTHRITIS PAIN**  
PAIN RELIEVING PATCH

Press  
Open

minty scent

Package not child resistant.  
Keep out of reach of children.

### Drug Facts

#### Active ingredients (in each patch)

#### Purpose

Menthol 3%.....	Topical analgesic
Methyl salicylate 10%(NSAID)*.....	Topical analgesic

\*nonsteroidal anti-inflammatory drug

#### Uses

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:

- arthritis
- sprains
- strains
- bruises
- simple backache

#### Warnings

##### For external use only

**Stomach bleeding warning:** This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

##### Do not use

- on the face or rashes
- on wounds or damaged skin
- if allergic to aspirin or other NSAIDs
- with a heating pad
- when sweating (such as from exercise or heat)
- any patch from a pouch that has been open for 14 or more days
- right before or after heart surgery

##### Ask a doctor before use if

- you are allergic to topical products
- the stomach bleeding warning applies to you
- you have high blood pressure, heart disease, or kidney disease
- you are taking a diuretic

##### When using this product

- wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

##### Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- rash, itching or skin irritation develops
- condition worsens
- symptoms last for more than 3 days
- symptoms clear up and occur again within a few days
- stomach pain or upset gets worse or lasts

**If pregnant or breast-feeding,** ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant.

#### Directions

##### Adults 18 years and older:

- clean and dry affected area
- remove patch from backing film and apply to skin (see illustration)
- apply one patch to the affected area and leave in place for up to 8 to 12 hours
- if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
- only use one patch at a time
- do not use more than 2 patches per day
- do not use for more than 3 days in a row

##### Children under 18 years of age:

- do not use

■ some individuals may not experience pain relief until several hours after applying the patch

■ avoid storing product in direct sunlight

■ protect product from excessive moisture

■ store at 20-25°C (68-77°F)

inactive: styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

(b) (4)



✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

FOLD TWICE AFTER OPENING

Discard unused patches 14 days after opening


FOLD TWICE AFTER OPENING

# SALONPAS<sup>®</sup>

## ARTHRITIS PAIN<sup>※</sup>

Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

*Effectiveness confirmed in clinical trial*



Questions or comments?  
Toll free 1-87-SALONPAS MON-FRI 9AM to 5PM (PST)

[www.salonpas.us](http://www.salonpas.us)

**※ DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS**

**Hisamitsu<sup>®</sup>**

Manufactured by  
**Hisamitsu Pharmaceutical Co., Inc.**  
JAPAN SAGA TOSU Made In Japan

**5** 2 3/4" x 3 15/16" (7cm x 10cm)  
**PATCHES**

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

FOLD TWICE AFTER OPENING

**Discard unused patches 14 days after opening**

FOLD TWICE AFTER OPENING

**SALONPAS<sup>®</sup> ARTHRITIS PAIN PAIN RELIEVING PATCH** minty scent

**Drug Facts**

Active ingredients (in each patch)	Purpose
Menthol 3%	Topical analgesic
Methyl salicylate 10%(NSAID)*	Topical analgesic
*nonsteroidal anti-inflammatory drug	

**Uses**  
Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:  
■ arthritis ■ sprains ■ strains ■ bruises ■ simple backache

**Warnings**  
**For external use only**  
**Stomach bleeding warning:** This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:  
■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use**  
■ on the face or rashes ■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs ■ with a heating pad ■ when sweating (such as from exercise or heat) ■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

**Ask a doctor before use if**  
■ you are allergic to topical products ■ the stomach bleeding warning applies to you ■ you have high blood pressure, heart disease, or kidney disease ■ you are taking a diuretic

**When using this product**  
■ wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water. ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**  
■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding. ■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days ■ symptoms clear up and occur again within a few days ■ stomach pain or upset gets worse or lasts

**If pregnant or breast-feeding**, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant.

**Directions**  
**Adults 18 years and older:**  
■ clean and dry affected area ■ remove patch from backing film and apply to skin (see illustration) ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours ■ only use one patch at a time ■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row

**Children under 18 years of age:**  
■ do not use

**Other information**  
■ some individuals may not experience pain relief until several hours after applying the patch ■ avoid storing product in direct sunlight ■ protect product from excessive moisture ■ store at 20-25°C (68-77°F)

LOT

**Inactive ingredients**  
alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

EXP.

**Package not child resistant. Keep out of reach of children.**

2L710T-0

(b) (4)

> CUT OPEN HERE ----- CUT OPEN HERE >

FOLD TWICE AFTER OPENING

Discard unused patches 14 days after opening

FOLD TWICE AFTER OPENING

# SALONPAS<sup>®</sup> ARTHRITIS PAIN<sup>※</sup>


Menthol 3%, Methyl Salicylate 10% PAIN RELIEVING PATCH

*Effectiveness confirmed in clinical trial*

NDC#46581-680-03

**FREE  
SAMPLE**

## ULTRA THIN COMFORT STRETCH



**FDA Approved Non Prescription  
Pain Relieving Patch**

Questions or comments?  
Toll free 1-87-SALONPAS MON-FRI 9AM to 5PM (PST) [www.salonpas.us](http://www.salonpas.us)

**※ DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS**

**Hisamitsu**

Manufactured by  
**Hisamitsu Pharmaceutical Co., Inc.**  
JAPAN SAGA TOSU Made in Japan

**3** 2 3/4" X 3 1/2" (7cm X 10cm)  
**PATCHES**

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

**SALONPAS<sup>®</sup> ARTHRITIS PAIN PAIN RELIEVING PATCH**

minty scent

**Drug Facts**

**Active Ingredients (in each patch)**

**Purpose**

Menthol 3% ..... Topical analgesic  
Methyl salicylate 10%(NSAID)\* ..... Topical analgesic  
\*nonsteroidal anti-inflammatory drug

**Uses**

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:  
■ arthritis ■ sprains ■ strains ■ bruises ■ simple backache

**Warnings**

**For external use only**

**Stomach bleeding warning:** This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:

■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ on the face or rashes ■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs ■ with a heating pad ■ when sweating (such as from exercise or heat) ■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

**Ask a doctor before use if** ■ you are allergic to topical products ■ the stomach bleeding warning applies to you ■ you have high blood pressure, heart disease, or kidney disease ■ you are taking a diuretic

**When using this product** ■ wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water. ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if** ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding. ■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days ■ symptoms clear up and occur again within a few days ■ stomach pain or upset gets worse or lasts

**If pregnant or breast-feeding,** ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant.

**Directions**

**Adults 18 years and older:**

■ clean and dry affected area ■ remove patch from backing film and apply to skin (see illustration) ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours ■ only use one patch at a time ■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row

**Children under 18 years of age:** ■ do not use

**Other information**

■ some individuals may not experience pain relief until several hours after applying the patch ■ avoid storing product in direct sunlight ■ protect product from excessive moisture ■ store at 20-25°C (68-77°F)

LOT

**Inactive Ingredients**

alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

EXP.

**Package not child resistant. Keep out of reach of children.**

3L711T-0

-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/

-----  
Joel Schiffenbauer  
6/19/2009 11:34:23 AM